


PHYS20040: Statistical Mechanics

Nigel Wilding



Welcome!

Remember to record your attendance

1. Download the University of Bristol app
2. Sign in with your university user-name and password
3. Select 'Record attendance'
(or tap on the notification)
4. If prompted, enable location services
5. Tap 'Check in' for this class, then 'Finish'
6. Your attendance confirmation will be displayed

Or go to check-in.bristol.ac.uk for web check-in

For more information or help, search for 'record attendance' on www.bristol.ac.uk



Apple App Store



Google Play Store

Delivery and format



Statistical Mechanics

- Detailed e-notes (see Blackboard) can be viewed on a variety of devices. Pdf available.
- I will give traditional lectures (Tues, Wed, Fri) in which I use slides to summarise and explain the lecture content. Questions welcome (within reason..)
- Try to read ahead in the notes, then come to lectures, listen to my explanations and then reread the notes.
- Rewriting the notes or slides to express your own understanding, or annotating a pdf copy can help wire the material into your own way of thinking.
- There are group problem classes (Tues, Thurs) where you can try problem sheets and seek help. I will go over some problems with the class.





1. Introduction

What is Statistical Mechanics?



Statistical Mechanics

Statistical mechanics, together with **classical thermodynamics**, form two branches of **thermal physics**

Each branch represents a distinct approach to thermal physics:

Macroscopic Approach (Classical Thermodynamics)

- deals with **macroscopic** variables i.e. variables that do not refer to any microscopic details
- input is phenomenological laws e.g. an equation of state, $P(V)$
- output is general relations between macroscopic variables
- advantage is the generality of the approach

Microscopic Approach (Statistical Mechanics)

- starts from a **microscopic** description and seeks to explain macroscopic properties
- input is a microscopic model of a given system
- output is predictions for macroscopic properties and behaviour
- predictions can be compared to experiment thus allowing refinement of the microscopic model

What is Statistical Mechanics?



Statistical Mechanics

	Thermodynamics	Statistical Mechanics
Quantity of interest	Macroscopic properties (eg. P, V, T, C_P, C_V)	Microscopic properties (eg: molecular speeds)
Strategy	Avoid microscopic model	Build on microscopic model
Strengths	Generality of results	Provides way of refining microscopic understanding
Weaknesses	No understanding of system-specific features. Conceptually opaque.	Requires additional input (the model). Requires additional techniques (probability theory; classical and quantum mechanics)

What is Statistical Mechanics?



Statistical Mechanics

- Provides powerful concepts and tools that help us understand the properties of complex systems with very many constituents.
- Used in research of systems as diverse as earthquakes, traffic jams, superconductivity, economics, and many more.
- **Aim of this course:** Show how key concepts that you have met in thermodynamics, such as the Boltzmann factor, entropy, and the second law, can be formulated and find expression in statistical terms.
- Develop and illustrate fundamental concepts and methods via two prototype systems:
 - Gases (classical and quantum)
 - Classical magnets
- For more advanced Soft Matter systems (eg. polymers, liquid crystals, glasses, surfactants, active matter) see M-level unit: **Complex and Disordered Matter**

The microscopic approach



- In principle, can imagine solving Newton's equation for the atomic and molecular motions in a system of interest to determine its behaviour.
- But typical systems contain of order a mol, ie. 10^{23} particles - more than all the grains of sand on all the beaches in the world, or stars in the visible universe!
- So exact approach is impractical and therefore we instead appeal to ideas from statistics and seek to make probabilistic statements about a systems behaviour.
- This works well because as the number of particles becomes very large, things get simpler...



Exercise: Revise your probability and statistics notes from first year laboratory, particularly on combinatorics, probability distributions, and summary statistics; read the section on probability in the lecture notes (end of sec 1.3)



Simplicity at large N

Toss a fair coin N times. Call this a “trial”.

What is the probability p_n of getting n heads from a trial?


Denote by p the probability that a head results from a single toss; then $q = 1 - p$ is probability for a tail.

This is binomial statistics. Recall:

$$\begin{aligned} p_n &\equiv \text{number of distinct ways of obtaining } n \text{ heads} \\ &\times \text{probability of any specific way of getting } n \text{ heads} \\ &= \binom{N}{n} p^n q^{N-n} \end{aligned}$$

The distribution p_n has mean $\bar{n} \equiv \sum_{n=0}^N n p_n = Np$

and variance $\overline{\Delta n^2} \equiv \sum_{n=0}^N (n - \bar{n})^2 p_n = Npq$





Simplicity at large N

- For a fair coin $p = q = \frac{1}{2}$
- Define $f = n/N$, the fraction of N tosses resulting in a head
- Mean of f


$$\bar{f} = \frac{\bar{n}}{N} = p = \frac{1}{2}$$

and standard deviation


$$(\overline{\Delta f^2})^{1/2} \equiv \frac{(\overline{\Delta n^2})^{1/2}}{N} = \left(\frac{pq}{N} \right)^{1/2} = \frac{1}{2N^{1/2}}$$

- The typical deviation of f from its mean value is thus vanishingly small for large N

Simplicity at large N



(a) For $N = 50$ tosses, we can be reasonably sure that f will be close to 0.5



(b) For $N = 10^{26}$ tosses We can be absolutely sure that f will be indistinguishable from 0.5