



# 11. Systems with varying particle number

# Motivation



- We developed the Boltzmann distribution assuming  $N$  is fixed and  $E$  is a free macroscopic variable.
- Now consider case where both  $N$  and  $E$  fluctuate.
- Why?
  1. Some systems are free to exchange particles eg. coexisting gas and liquid phases where a molecule may either be part of the liquid phase or of the gas phase.
  2. If  $N$  is a free macro variable, for a large system it is sharply defined at some mean value and thus we get the same behavior as a system of fixed particle number.
  3. Useful means-to-an-end for simplifying study of quantum gases



# The chemical potential

- Consider a system where two halves are free to exchange particles.
- Since  $N = N_1 + N_2$  is conserved,  $dN = 0$  and hence:


$$dN_1 = -dN_2$$

- Now free energy is extensive ie.  $F = F_1(N_1) + F_2(N_2)$

$$dF = \frac{\partial F_1(N_1)}{\partial N_1} dN_1 + \frac{\partial F_2(N_2)}{\partial N_2} dN_2 = \left[ \frac{\partial F_1(N_1)}{\partial N_1} - \frac{\partial F_2(N_2)}{\partial N_2} \right] dN_1$$

- As system at equilibrium, free energy is at a minimum,  $dF = 0$

$$\Rightarrow \frac{\partial F_1(N_1)}{\partial N_1} = \frac{\partial F_2(N_2)}{\partial N_2}$$





- c.f. sec 3.5 where we used analogous arguments to deduce the existence of a common temperature

# The chemical potential



- Thus the quantity  $\mu \equiv \left( \frac{\partial F}{\partial N} \right)_{T,V}$

is common to both systems that can exchange particles.  
We call  $\mu$  the **chemical potential**.



- If a system is not in equilibrium so that there is a chemical potential gradient, then particles will diffuse down then gradient
- Similar to heat diffusing down a temperature gradient

# The chemical potential



- An equivalent definition of the chemical potential appropriate for an isolated system (fixed  $E$ ) is found as follows:

$$\begin{aligned}\mu &\equiv \left( \frac{\partial F}{\partial N} \right)_{T,V} \\ &= \frac{\partial}{\partial N} (E - TS(E, N))_{T,V} \\ &= -T \left( \frac{\partial S}{\partial N} \right)_{E,V}\end{aligned}$$

# Example



- Two connected volumes of gas at different heights in a gravitational field

$$Z(N_l, N_u) = Z_l(N_l) \times Z_u(N_u) = \left(\frac{V}{\lambda^3}\right)^{N_l} \frac{1}{N_l!} \times \left(\frac{V}{\lambda^3}\right)^{N_u} e^{-\beta M g H N_u} \frac{1}{N_u!}$$

$$\lambda = \left(\frac{h^2}{2\pi M k T}\right)^{1/2}$$

$$F_l = -kT \ln Z_l = N_l kT [\ln(N_l \lambda^3 / V) - 1]$$


$$F_u = N_u kT [\ln(N_u \lambda^3 / V) - 1] + N_u M g H$$

$$\mu_l = \frac{\partial F_l}{\partial N_l} = kT \ln \left(\frac{N_l}{V \lambda^3}\right)$$

$$\mu_u = \frac{\partial F_u}{\partial N_u} = kT \ln \left(\frac{N_u}{V \lambda^3}\right) + M g H$$


$$\mu_l = \mu_u \Rightarrow kT \ln \left(\frac{N_l}{N_u}\right) = M g H$$

$$\Rightarrow N_u = N_l e^{-M g H / k T}$$



- Consistent with the Boltzmann distribution

# The grand canonical distribution



- A microstate  $r$  of the system has energy  $E$  and  $N$  particles.

$$P_r \propto \Omega_b(E_{TOT} - E, N_{TOT} - N)$$

$$= \exp \left( \frac{S_b(E_{TOT} - E, N_{TOT} - N)}{k} \right)$$

$$S_b(E_{TOT} - E, N_{TOT} - N) = S_b(E_{TOT}, N_{TOT}) - E \frac{\partial S_b(E_{TOT}, N_{TOT})}{\partial E} - N \frac{\partial S_b(E_{TOT}, N_{TOT})}{\partial N} + \dots$$

$$= \text{const} - \frac{E}{T} + \frac{N\mu}{T}$$

$$P_r \propto \exp \left( \frac{1}{kT} (N\mu - E) \right)$$

• Key point 15:  $P_r = \frac{1}{\mathcal{Z}} \exp(-\beta E_r + \beta \mu N_r), \quad \mathcal{Z} = \sum_j \exp(-\beta E_j + \beta \mu N_j), \quad \beta = \frac{1}{kT}$



## 12. Quantum gases