

Geometry Before Fields: Constraint-Reduced Cosmology and Optical Predictions in the Berger–Hopf Framework

Norman P. Carberry
Independent Researcher

Working draft — September 2026

Abstract

We develop a geometry-first cosmological workflow in which physical degrees of freedom are identified after constraints, boundary conditions, gauge redundancies and the physical Hessian are resolved. Within a conditional Berger–Hopf cosmological bridge on a closed hyperspherical parent, a selected physical $n = 2$ harmonic carries one two-component phase-space state. Its induced metric response supplies reference growth and Weyl kernels and a gauge-invariant luminosity-distance map at fixed observed redshift. Jacobi/Sachs equations propagate photon redshift, convergence, shear and their stochastic covariance. For a fixed response to m coherent random texture amplitudes, the isolated topographic covariance has rank at most m ; one local noise type alone does not imply rank one. We formulate two prospective rejection gates at $p < 0.01$: consistency of deterministic observables with one shared state, and consistency of isolated covariance with the minimal one-amplitude realization. Survey selection, nuisance treatment and finite-sample calibration must be frozen before target inspection. Homogeneous localization and three foreground-transfer candidates fail their retained tests. A curvature-dependent defect in one effective representation is reported without generalizing it to EFT or QFT. Complete compact-background normalization, action-owned local couplings and source statistics remain open. The contribution is a reproducible, conditional response chain and explicit falsification criteria, rather than a completed microscopic cosmology.

Keywords: cosmology; hyperspherical geometry; constrained dynamics; optical scalar equations; luminosity distance; weak lensing; stochastic geometry; falsifiable cosmology

1 Introduction

Cosmological models must connect a physical state to several observations without adding new freedoms at each interface. This paper develops that requirement as a geometry-first workflow: specify a geometric action, resolve constraints, boundary conditions and gauge redundancy, reduce the physical Hessian, and propagate the retained response to matter and light.

The Berger–Hopf Standard Model (BHSM) program motivates this ordering. Neither constrained perturbation theory nor Schur reduction is new. Our contribution is their organization into an explicit closed- S^3 cosmological response chain, a calibration-rank audit, retained failed mechanisms, and two prospective rejection gates. Covariant EFT can also respect constraints and gauge symmetry; this workflow does not distinguish BHSM from every effective theory.

The regular one-scalar closed-FLRW quotient removes the exceptional $n = 1$ scalar sector. The first ordinary scalar carrier is $n = 2$ in our convention. One selected normalized harmonic carries $X_2 = (q_2, \Pi_2)^T$; the entire $n = 2$ eigenspace is not two dimensional. Its profile and axis must

be fixed separately by theory or independent calibration. Identification with the retained BHSM sector remains conditional.

The downstream results are a constrained metric response, reference growth/Weyl and fixed-observed-redshift distance kernels, and Jacobi/Sachs propagation of optical means and covariances. Section 17 asks whether deterministic measurements admit one common two-component state and whether isolated covariance admits one coherent texture amplitude, with separate prospective $p < 0.01$ rejection gates.

This manuscript substantially expands and reframes the prior Preprints.org study [1]. The physical carrier, gauge audit, optical propagation and explicit negative results develop and correct that earlier proposal. We do not claim a resolution of the Hubble tension, a general failure of EFT/QFT, or an ab initio derivation of every absolute cosmological amplitude.

2 Methods: geometry and physical reduction

The methodological ordering is

$$\begin{aligned}
 &\text{geometry} \rightarrow \text{action} \rightarrow \text{constraints / boundary conditions} \\
 &\quad \rightarrow \text{gauge quotient} \rightarrow \text{physical tangent space} \\
 &\quad \rightarrow \text{physical Hessian} \rightarrow \text{reduced response} \\
 &\quad \rightarrow \text{metric / matter response} \rightarrow \text{optics} \rightarrow \text{observables.}
 \end{aligned} \tag{1}$$

Boundary conditions specify the operator domain before inversion. At a regular constrained stationary point, a gauge-fixed representative is $T_{\text{phys}} = \ker DC \cap \mathcal{G}^\perp$. This notation assumes a declared inner product and a valid gauge slice; it does not establish their global existence. The physical second variation is $P_{\text{phys}} D^2 \mathcal{L}_{\text{KKT}} P_{\text{phys}}$. Eliminating an invertible physical complement gives $\mathcal{K}_{\text{phys}} = A - BK^{-1}B^\dagger$.

On the same domain, the metric block obeys

$$\mathcal{A}_g h + \mathcal{B}_{g2} X_2 + J_g = 0, \tag{2}$$

$$\mathcal{R}_{g2} = -\mathcal{A}_g^{-1} \mathcal{B}_{g2}, \tag{3}$$

$$h_2 = h_{\text{src}} + \mathcal{R}_{g2} X_2, \quad h_{\text{src}} = -\mathcal{A}_g^{-1} J_g. \tag{4}$$

The inverse requires a nonsingular retained block. The induced metric has no extra initial datum. The reference homogeneous response sets the independent source contribution to zero; target fits must subtract any independently specified source before applying the shared-state test.

BHSM supplies the proposed common action, boundary-sector interpretation and projection contracts. This repository implements the downstream cosmological bridge; complete compact-background matrix elements remain open. A useful effective representation need not have been proved globally equivalent to that bridge. The particular curved-EFT failure reported below is a limitation of that audited construction, not of EFT or QFT in general.

2.1 Reproducibility and computational workflow

Versioned code, frozen manifests, numerical artifacts and regression tests separate prediction generation from target comparison. The source package and repository guide distinguish numerical checks from observational reruns whose exact historical input files are not bundled. OpenAI ChatGPT and Codex materially assisted the workflow; their role and the author's responsibility are disclosed in the back matter. AI output is not accepted as independent evidence. Rejected candidates and open constitutive inputs retain their original claim classes.

3 Claim status and falsifiability

Status	Statement and boundary
DERIVED	Schur metric response on an invertible physical domain; regular closed-FLRW harmonic exclusion; fixed-observed-redshift distance and Jacobi/Sachs identities within linear geometric optics.
DERIVED	$C_{\text{topo}} = M_O C_q M_O^\dagger$ implies rank at most m for a fixed finite m -amplitude response. This does not derive the physical channel count.
CALIBRATED / CONDITIONAL	Identification of the BHSM envelope with one $n = 2$ profile; R1 coefficients, angular axis, SN amplitude and gravity–scalar reference kernels. One amplitude leaves one state direction undetermined.
REJECTED / NOT ESTABLISHED	Homogeneous two-state localization and frozen 2MRS number-density, K-band luminosity and seam-charge foreground transfer candidates.
OPEN	Normalized compact full-preimage background, action-owned local matrix elements, seam export, rare-event statistics, absolute amplitudes and survey-ready target/nuisance design.

Table 1: Current claim ledger. Exact identities are distinguished from physical identifications and empirical claims.

Section 17 defines rejection of specific deterministic or stochastic realizations. Additional states, channels, amplitudes, kernels or selected bins may not be introduced after the decisive result.

4 Hyperspherical Parent and Harmonic Convention

We use a closed FLRW geometry

$$ds^2 = -dt^2 + a^2(t)R_H^2 \left[d\psi^2 + \sin^2 \psi d\Omega_2^2 \right],$$

with $a_0 = 1$. Define

$$D_C(z) = c \int_0^z \frac{dz'}{H(z')}, \quad \psi(z) = \frac{D_C(z)}{R_H}.$$

Scalar harmonics satisfy

$$-\nabla_\gamma^2 Y_n = \frac{n(n+2)}{R_H^2} Y_n, \quad n = 0, 1, 2, \dots,$$

so the physical spatial eigenvalue is

$$p_n(a) = \frac{n(n+2)}{a^2 R_H^2}.$$

A common closed-universe convention uses $k^2 = N^2 - 1$, hence

$$\boxed{N = n + 1}.$$

4.1 Exceptional first nonconstant harmonic

For our $n = 1$, $k^2 = 3$ and $N = 2$. The degree-one scalar harmonics on the unit three-sphere obey

$$D_i D_j Q^{(1)} = -\gamma_{ij} Q^{(1)}.$$

Therefore

$$\left(D_i D_j + \frac{k^2}{3} \gamma_{ij} \right) Q^{(1)} = 0.$$

The scalar-derived trace-free metric harmonic is absent. For the regular generally covariant one-scalar sector, the remaining scalar gauge transformations and lapse/shift constraints leave no ordinary propagating scalar degree of freedom in this exceptional sector. Modern gauge-invariant closed-universe analyses likewise take the propagating scalar spectrum to begin at alternative-convention $N \geq 3$ [2, 3].

Accordingly, the minimal bridge no longer uses our $n = 1$ as the physical scalar–metric carrier. An additional boundary degree of freedom could evade this conclusion, but it is not required for the construction below.

4.2 First physical scalar harmonic and exact dipole subspace

The first propagating scalar harmonic is

$$\boxed{n_\star = 2}, \quad \boxed{N_\star = 3}, \quad p_2 = \frac{8}{a^2 R_H^2}.$$

Let the unit $S^3 \subset \mathbb{R}^4$ embedding coordinates be

$$X^4 = \cos \psi, \quad X^i = \sin \psi \hat{n}^i.$$

Every real degree-two scalar harmonic can be written

$$T_2 = A_{AB} X^A X^B, \quad A_{AB} = A_{BA}, \quad \delta^{AB} A_{AB} = 0.$$

Writing $c = \cos \psi$, $s = \sin \psi$, $a_i = A_{4i}$, and

$$A_{ij} = S_{ij} - \frac{A_{44}}{3} \delta_{ij}, \quad S_i{}^i = 0,$$

gives the exact observer-shell decomposition

$$\boxed{T_2 = A_{44} \left(c^2 - \frac{s^2}{3} \right) + 2cs \mathbf{a} \cdot \hat{\mathbf{n}} + s^2 S_{ij} \hat{n}^i \hat{n}^j.}$$

These terms are respectively a sky monopole, dipole, and quadrupole. The pure cross-term subspace

$$A_{44} = 0, \quad S_{ij} = 0, \quad a_i \neq 0$$

therefore gives an exact sky dipole:

$$\boxed{T_{2,\text{dip}} = A \sin(2\psi) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}).}$$

Thus the gauge audit shifts the carrier to the first physical compact-space scalar mode while retaining an exact dipolar observer-sky sector.

5 Constraint-Reduced Response and the Topographic Kernel

Let $\hat{\mathcal{H}}$ be the physical Hessian after constraints and gauge directions have been removed. For a selected topographic branch P and physical complement Q , the exact Schur/Feshbach reduced operator is

$$\mathcal{K} = P \left[\hat{\mathcal{H}} - \hat{\mathcal{H}}Q(Q\hat{\mathcal{H}}Q)^{-1}Q\hat{\mathcal{H}} \right] P.$$

For one scalar harmonic, write the quadratic block as

$$\hat{\mathcal{H}}_n = \begin{pmatrix} A(p) & C(p) \\ C^\dagger(p) & D(p) \end{pmatrix}.$$

The exact reduced stiffness is

$$K_{\text{red}}(p) = A(p) - C(p)D^{-1}(p)C^\dagger(p).$$

A minimal long-wavelength realization is

$$A(p) = Z_0 p, \quad D(p) = M_c^2 + c_c p, \quad C(p) = g\sqrt{p}.$$

Then

$$K_{\text{red}}(p) = Z_0 p - \frac{g^2 p}{M_c^2 + c_c p}.$$

For $c_c p \ll M_c^2$,

$$K_{\text{red}}(p) = \left(Z_0 - \frac{g^2}{M_c^2} \right) p + \frac{g^2 c_c}{M_c^4} p^2 + O(p^3).$$

Thus a positive p^2 stiffness can arise from a coupled second-order parent system after a stable complement is integrated out. In this interpretation the old static operator

$$-\nabla^2 + B\nabla^4$$

is a low-energy spatial derivative expansion, not evidence for a fundamental fourth-order time equation.

Define

$$\chi \equiv \frac{g^2}{M_c^2}, \quad L_c^2 \equiv \frac{c_c}{M_c^2}.$$

After canonical normalization of the leading spatial term, the reduced kernel may be written schematically as

$$K_{\text{topo}}(p) = m_T^2 + (c_s^2 - \chi)p + \chi L_c^2 p^2 + \dots.$$

The signs of the temporal kinetic term and of the full physical gradient matrix must be checked independently; a positive spatial p^2 coefficient by itself is not a complete ghost analysis.

6 BHSM-native transport of the physical topographic mode

The purpose of this section is to separate the ontology of the Berger–Hopf Standard Model (BHSM) mode from any particular four-dimensional effective-field-theory coordinate used to represent its late-time response. The distinction is important in the closed cosmological setting: the physical topographic carrier is not introduced as an additional free bulk scalar. Rather, it is an action-owned geometric/boundary mode whose physical dynamics are obtained only after the BHSM constraints and gauge directions have been removed.

6.1 Action ownership and physical tangent space

The BHSM construction used in this work treats modes as physical boundary-local fields acted on by sector projectors. The scalar/topographic sector is represented in the unified action candidate and admits an action-reduced mode functional together with a normalized profile-boundary boundary-value problem. These ingredients motivate the following physical tangent space at a background configuration $z_\star$:

$$T_{\text{phys}}(z_\star) = \ker D\mathcal{C}(z_\star) \cap \mathcal{G}^\perp, \quad (5)$$

where $\mathcal{C}$ denotes the action-owned constraints and $\mathcal{G}$ the infinitesimal gauge directions. The corresponding projected Hessian is

$$\hat{\mathcal{H}} = P_{\text{phys}} D^2 \mathcal{L}_{\text{KKT}}(z_\star) P_{\text{phys}}. \quad (6)$$

If P_2 denotes the projector onto the first admissible physical scalar-topographic harmonic, the constraint-eliminated quadratic operator is the Schur–Feshbach reduction

$$\mathcal{K}_2 = P_2 \left[\hat{\mathcal{H}} - \hat{\mathcal{H}} Q_2 \left(Q_2 \hat{\mathcal{H}} Q_2 \right)^{-1} Q_2 \hat{\mathcal{H}} \right] P_2, \quad (7)$$

whenever the eliminated block is invertible on the chosen physical domain. Here $Q_2 = 1 - P_2$ on the already gauge-reduced tangent space.

6.2 Cosmological topographic-mode theorem

Theorem 1 (action-owned $n = 2$ envelope transport). Assume that:

1. the cosmological parent envelope is the closed spatial manifold $\mathcal{E}_{\text{cos}} \simeq S^3(R_H)$;
2. the scalar/topographic perturbation belongs to the BHSM action-owned boundary-mode sector and is reduced to T_{phys} before physical propagation is assigned;
3. the homogeneous $n = 0$ component is absorbed into the background and the exceptional scalar $n = 1$ sector is removed by the regular closed-FLRW constraint/gauge quotient;
4. the reduced $n = 2$ kinetic form is nondegenerate and positive on the retained mode.

Then the first physical scalar-topographic carrier is an action-owned $n = 2$ envelope mode. It does not require the introduction of an additional fundamental bulk scalar degree of freedom. Its canonical momentum, when nonzero, is induced by the reduced action itself and is therefore conjugate data of the same geometric mode rather than a separately postulated field.

Proof. Let $e_2(x)$ be a normalized element of $P_2 T_{\text{phys}}$ and write the retained perturbation locally as

$$\delta\Phi_{\text{topo}}(t, x) = q_2(t) e_2(x). \quad (8)$$

Substitution into the gauge- and constraint-reduced quadratic action gives, up to linear source terms and a choice of time parametrization,

$$S_2^{(2)} = \frac{1}{2} \int dt a^3 \left[\mathcal{G}_2(t) (D_t q_2)^2 - \mathcal{M}_2^2(t) q_2^2 + 2 \mathcal{J}_2(t) q_2 \right]. \quad (9)$$

The coefficients are the e_2 matrix elements of the reduced operator (7); explicitly, $\mathcal{G}_2$ is the kinetic quadratic form, $\mathcal{M}_2^2$ the reduced spatial/restoring form, and $\mathcal{J}_2$ the projection of the action-owned source onto e_2 . The canonical momentum is therefore

$$\Pi_2 = a^3 \mathcal{G}_2 D_t q_2, \quad (10)$$

which is generated by the same reduced action. Variation of (9) yields

$$D_t \left(a^3 \mathcal{G}_2 D_t q_2 \right) + a^3 \mathcal{M}_2^2 q_2 = a^3 \mathcal{J}_2. \quad (11)$$

Thus the dynamical phase-space pair (q_2, Π_2) is the reduced description of one action-owned envelope mode. No second ontological scalar is created by introducing an effective coordinate for that mode. $\square$

6.3 Relation to the exact closed-Horndeski kernel

The closed-Horndeski calculation of Sec. B is used here as a late-time effective response representation of the same reduced mode, not as the microscopic definition of the BHSM topographic ontology. In the Bellini–Sawicki convention used in that section, the exact physical- $n = 2$ kinetic coefficient is

$$\mathcal{G}_{S,2} = M_{\text{Pl}}^2 \frac{5D_{\text{kin}}}{D_{\text{kin}} + 10(1 - \alpha_B^{\text{BS}}/2)^2}. \quad (12)$$

For $D_{\text{kin}} > 0$ and nonsingular braiding denominator this coefficient is positive. Equation (12) therefore supplies an effective kinetic matching condition for assumption 4 of Theorem 1 on the reference branch.

The Stueckelberg variable π used in the EFT equations is accordingly treated as a coordinate representative of the retained response of q_2 . It is not counted as an additional BHSM mode. A change of EFT normalization amounts to an invertible local redefinition of the same reduced coordinate.

6.4 Exact observer dipole inside the physical $n = 2$ sector

A degree-two harmonic on S^3 may be written

$$T_2 = A_{AB} X^A X^B, \quad A_A{}^A = 0. \quad (13)$$

Relative to an observer shell,

$$T_2 = A_{44} \left(c^2 - \frac{s^2}{3} \right) + 2cs \mathbf{a} \cdot \hat{\mathbf{n}} + s^2 S_{ij} \hat{n}^i \hat{n}^j. \quad (14)$$

The pure A_{4i} subspace is therefore

$$\boxed{T_{2,\text{dip}} = A \sin(2\psi) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})}, \quad (15)$$

so the observed dipolar angular dependence is embedded exactly in the first physical scalar-topographic harmonic.

6.5 Role of the curved EFT Bianchi audit

The curvature-aware EFT system of the following sections remains a valuable cross-check of a chosen four-dimensional response representation. It is not used as the ontological definition of the BHSM mode and it is not allowed to introduce an additional independent scalar degree of freedom.

At the present stage, the literal curvature-aware 00, 0i, spatial-trace, and scalar equations and the reference background have been independently transcription-audited. The fully integrated curved-EFT differential- algebraic representation nevertheless retains a nonzero constraint- preservation defect, while the consistently flattened system closes to numerical precision. We therefore classify that calculation as an *EFT compatibility audit* rather than as the defining transport theorem.

The defining evolution statement for the bridge is Eq. (11): propagation is generated by the action-reduced physical $n = 2$ operator. A future exact identification of the curved EFT transfer map with $\mathcal{K}_2$ must reproduce this reduced phase space without duplicating degrees of freedom.

6.6 Claim boundary

Theorem 1 establishes the mode ownership and the form of its reduced transport law conditional on the BHSM action-owned scalar/topographic sector, the physical quotient, and positivity of the retained kinetic form. It does not by itself determine an absolute normalization for $\mathcal{M}_2^2$, $\mathcal{J}_2$, or the observable transfer coefficient. Those quantities are supplied, where stated, by the frozen reference branch or by explicit observational calibration and are labelled accordingly.

Consequently, the exact $n = 2$ dipole geometry, the physical-harmonic selection, and the positive effective kinetic matching are structural results. Reference growth curves and other quantities that require a particular EFT transfer realization remain reference-branch predictions rather than framework-wide theorems.

7 First numerical R1 physical- $n = 2$ reduced propagator

The representation-invariant prediction track can now be evaluated at the reduced gravity-scalar level. For the physical $n = 2$ harmonic, $-(\mathcal{D}^2 + 3K) = 5K$.

The exact reduced equation integrated here is

$$\zeta_{,NN} + \left(3 + \frac{d \ln H}{dN} + \frac{d \ln \mathcal{G}_{S,2}}{dN}\right) \zeta_{,N} + \frac{5K}{a^2 H^2} \frac{\mathcal{F}_{S,2}}{\mathcal{G}_{S,2}} \zeta = 0. \quad (16)$$

The fundamental matrix is normalized to the identity at $z = 2.1$ in the state $(\zeta, d\zeta/dN)^T$.

$$U(0, 2.1) = \begin{pmatrix} -0.186499277 & 0.019664322 \\ 0.013019584 & -0.005651284 \end{pmatrix}. \quad (17)$$

Across $0 \leq z \leq 2.1$ the numerical scan gives $\min \mathcal{G}_{S,2} = 1.471470 \times 10^{-3}$ and $\min \mathcal{F}_{S,2} = 5.037267 \times 10^{-1}$. The maximum absolute residual of the exact lapse/shift constraints over both fundamental solutions is 2.720×10^{-14} , and the Liouville determinant identity is reproduced with relative residual 4.171×10^{-14} .

7.1 Claim boundary

This is the first numerical phase-space transfer kernel for the physical $n = 2$ R1 mode. It uses the exact reduced closed-Horndeski $n = 2$ kinetic coefficient and constraints. The current $\mathcal{F}_{S,2}$

implementation is explicitly the gravity–scalar expression; the complete matter+radiation scalar stability/response system is not yet included. Consequently this result is not yet the directional $f\sigma_8$, lensing, or luminosity-distance observable kernel.

8 Direct $n = 2$ matter-growth and Weyl transfer

The reduced physical $n = 2$ propagator can be coupled directly to minimally coupled matter through covariant energy–momentum conservation, without invoking the curvature-defective Newtonian-gauge DAE.

For pressureless matter in unitary gauge,

$$\dot{v}_m = -\delta n, \quad (18)$$

$$\dot{\delta}_m + 3\frac{d}{dt}(Hv_m) = -\left(3\dot{\zeta} + \frac{k_2^2}{a^2}\chi\right) + \frac{k_2^2}{a^2}v_m, \quad (19)$$

with $k_2^2/a^2 = 8K/a^2$ for the physical manuscript $n = 2$ harmonic.

In the same unitary-gauge convention the gauge-invariant potentials are

$$\Psi = \delta n + \dot{\chi}, \quad \Phi = \zeta + H\chi, \quad (20)$$

so the local Weyl/lensing combination in this sign convention is $\Psi - \Phi$.

The response is integrated for both basis columns of the physical state $(\zeta, d\zeta/dN)^T$, with zero matter response at the $z = 2.1$ anchor. Thus the result is a row-vector kernel on the same two-dimensional physical state space.

8.1 Present-epoch state-space kernels

$$\mathcal{F}_{f\sigma_8}^{\text{state}}(0) = (2.172303247, 3.071170118 \times 10^{-2}), \quad (21)$$

$$\mathcal{F}_{\text{Weyl}}^{\text{state}}(0) = (6.167895805 \times 10^{-2}, -2.498933170 \times 10^{-2}). \quad (22)$$

These are not two new amplitudes. They are the two components of linear functionals acting on the same initial $n = 2$ state. One amplitude gives unique predictions only after the normalized state direction is independently fixed; the distance row alone has rank one.

8.2 Claim boundary

The $f\sigma_8$ object above is a state-space fractional response kernel, using the R1 baseline dD/dN normalization. It is not yet the final single-number directional prediction because the full D_L calibration functional has not yet selected the initial-state vector. Likewise $\Psi - \Phi$ is the local Weyl potential combination; the integrated convergence/shear line-of-sight kernel remains to be evaluated.

9 Gauge-invariant luminosity-distance kernel and calibration-rank audit

The remaining light-cone observable is evaluated with the gauge-invariant geometric luminosity-distance construction.[4] For positive spatial curvature the background transverse distance and

lensing efficiency use $S_K(\chi) = \sin(\sqrt{K}\chi)/\sqrt{K}$, with the corresponding curved-FLRW line-of-sight kernel.[5]

For the pure A_{4i} physical $n = 2$ mode, per unit $\mu = \hat{\mathbf{n}} \cdot \hat{\mathbf{p}}$,

$$Q_{2,\text{dip}}(\chi) = \sin(2\sqrt{K}\chi), \quad \hat{\nabla}^2 Q_{2,\text{dip}} = -2Q_{2,\text{dip}}. \quad (23)$$

The fixed-observed-redshift distance fluctuation is

$$\boxed{\frac{\delta D_L}{D_L} = \phi_s + \delta z + \frac{C_K(\chi_s)}{S_K(\chi_s)} \delta\chi - \kappa,} \quad (24)$$

with $C_K(\chi) = \cos(\sqrt{K}\chi)$ and the redshift, radial and convergence distortions evaluated along the same R1 physical- $n = 2$ metric response. Here δz denotes the dimensionless perturbation $\delta \ln(1+z)$, not an unnormalized change in the measured redshift.

9.1 State-space calibration row

At the reference bin center $z_\star = 0.02$, the resulting luminosity-distance row on the $(\zeta, d\zeta/dN)$ state anchored at $z = 2.1$ is

$$\boxed{\mathcal{F}_{D_L}(z_\star) = (1.842472667 \times 10^{-1}, -3.520884651 \times 10^{-3}).} \quad (25)$$

The carried-forward SN calibration imposes one scalar constraint on this two-dimensional state:

$$\mathcal{F}_{D_L}(z_\star) \cdot X_{2,\text{anchor}} = -1.897330117 \times 10^{-2} \pm 4.144653167 \times 10^{-3}. \quad (26)$$

The corresponding unit null direction is

$$n_{\text{cal}} = (1.910607424 \times 10^{-2}, 9.998174623 \times 10^{-1}), \quad \mathcal{F}_{D_L} \cdot n_{\text{cal}} = 0. \quad (27)$$

9.2 Calibration-rank consequence

This numerical light-cone calculation exposes a logical condition implicit in the single-calibration theorem: one amplitude calibration determines the model only after the normalized physical mode trajectory $\hat{X}_2$ has been fixed independently. The luminosity-distance row has rank one on a two-dimensional phase space. The growth and Weyl rows are not generally parallel to it, so motion along n_{cal} leaves the SN calibration unchanged while altering the other observables.

The next prediction-closure object is the theory-selected normalized state direction $\hat{X}_2$.

9.3 Claim boundary

No second observational calibration is introduced. Until the branch direction is fixed by the retained action or by an explicit preregistered theoretical branch condition, the manuscript reports the observable kernels as state-space rows rather than unique scalar amplitudes.

10 BHSM optical transfer through a stochastic topographic geometry

The preceding sections separate the closed hyperspherical parent $S^3(R_H)$, the physical BHSM $n = 2$ carrier, and local line-of-sight topography. A realistic photon path, however, cannot be represented

faithfully as a finite sum over foreground objects. In geometric optics the correct compression is the optical tidal matrix sampled by the null bundle. This section therefore derives a first-order BHSM optical transfer system in which unresolved foreground complexity is represented by a continuous geometric source and, when appropriate, by its path covariance.

10.1 Exact null-bundle geometry

Let k^μ be the tangent to the central null geodesic,

$$k^\nu \nabla_\nu k^\mu = 0, \quad k^\mu k_\mu = 0, \quad (28)$$

and let e_A^μ , $A = 1, 2$, be a parallel-transported Sachs screen basis. The Jacobi matrix $\mathcal{D}_{AB}$ obeys

$$\frac{d^2 \mathcal{D}}{d\lambda^2} = \mathcal{T} \mathcal{D}, \quad \mathcal{T}_{AB} = -R_{\mu\nu\alpha\beta} e_A^\mu k^\nu k^\alpha e_B^\beta. \quad (29)$$

The photon frequency for an observer congruence u^μ is

$$\omega = -u_\mu k^\mu, \quad \frac{d\omega}{d\lambda} = -k^\mu k^\nu \nabla_\mu u_\nu. \quad (30)$$

For the closed background define

$$S_K(\chi) = \frac{\sin(\sqrt{K} \chi)}{\sqrt{K}}, \quad C_K(\chi) = \cos(\sqrt{K} \chi), \quad H_K(\chi) = \frac{C_K(\chi)}{S_K(\chi)}. \quad (31)$$

In conformal background-distance coordinates, after removing the background scale factor, the background Jacobi map is $S_K I$. Primes below refer to χ , not the physical affine parameter in Eq. (29); $\delta\mathcal{T}$ is the correspondingly rescaled effective source. Writing

$$\mathcal{D} = S_K(\chi)[I + \mathcal{A}] \quad (32)$$

and linearizing $\mathcal{T} = -KI + \delta\mathcal{T}$ gives

$$\boxed{\mathcal{A}'' + 2H_K \mathcal{A}' = \delta\mathcal{T}.} \quad (33)$$

10.2 Seven-state optical system

Decompose

$$\mathcal{A} = -\kappa I - \begin{pmatrix} \gamma_1 & \gamma_2 \\ \gamma_2 & -\gamma_1 \end{pmatrix} \quad (34)$$

and define

$$\mathcal{F} = -\frac{1}{2} \text{Tr } \delta\mathcal{T}, \quad \mathcal{G}_1 = -\frac{1}{2}(\delta\mathcal{T}_{11} - \delta\mathcal{T}_{22}), \quad \mathcal{G}_2 = -\delta\mathcal{T}_{12}. \quad (35)$$

Then

$$\kappa'' + 2H_K \kappa' = \mathcal{F}, \quad (36)$$

$$\gamma_1'' + 2H_K \gamma_1' = \mathcal{G}_1, \quad (37)$$

$$\gamma_2'' + 2H_K \gamma_2' = \mathcal{G}_2. \quad (38)$$

Let

$$\zeta_\gamma = \delta \ln(1+z), \quad \zeta'_\gamma = \mathcal{Z}(\chi), \quad (39)$$

and define

$$Y_\gamma = (\zeta_\gamma, \kappa, p_\kappa, \gamma_1, p_1, \gamma_2, p_2)^T. \quad (40)$$

The optical propagation law is

$$\boxed{\frac{dY_\gamma}{d\chi} = A_K Y_\gamma + B_{\text{opt}} S_{\text{opt}},} \quad (41)$$

where

$$S_{\text{opt}} = (\mathcal{Z}, \mathcal{F}, \mathcal{G}_1, \mathcal{G}_2)^T, \quad (42)$$

$$A_K = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2H_K & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2H_K & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -2H_K \end{pmatrix}, \quad (43)$$

and

$$B_{\text{opt}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (44)$$

10.3 BHSM closure

The physical global mode is $X_2 = (q_2, \Pi_2)^T$ and the existing metric-response theorem gives

$$h_2 = \mathcal{R}_{g2} X_2, \quad \mathcal{R}_{g2} = -\mathcal{A}_g^{-1} \mathcal{B}_{g2}. \quad (45)$$

For local topography define

$$h_{\text{topo}} = \mathcal{R}_{g\psi} \psi, \quad \mathcal{R}_{g\psi} = -\mathcal{A}_g^{-1} \mathcal{B}_{g\psi}, \quad (46)$$

with its complete cosmological normalization left open.

The local field is written

$$\boxed{\mathcal{L}_\psi \psi = J_{\text{grav}} + \mathcal{E}_\Sigma[Q_\Sigma] + \xi,} \quad (47)$$

where the static spatial limit of $\mathcal{L}_\psi$ must reproduce $K_T = -\nabla^2 + B\nabla^4$ and

$$Q_\Sigma = V_\Sigma^2 = \frac{c^2}{8\pi} \int_{\Sigma_\star} H_\Sigma \Delta K_{uu} dA. \quad (48)$$

The seam-export operator $\mathcal{E}_\Sigma$ is explicitly action-normalization-open.

If $\mathcal{O}_{\text{opt}}$ maps the physical metric response to $(\mathcal{Z}, \mathcal{F}, \mathcal{G}_1, \mathcal{G}_2)$ and $\mathcal{G}_\psi = \mathcal{L}_\psi^{-1}$, then

$$\boxed{\begin{aligned} \frac{dY_\gamma}{d\chi} &= A_K Y_\gamma \\ &+ B_{\text{opt}} \mathcal{O}_{\text{opt}} [\mathcal{R}_{g2} X_2 + \mathcal{R}_{g\psi} \mathcal{G}_\psi (J_{\text{grav}} + \mathcal{E}_\Sigma[Q_\Sigma] + \xi)]. \end{aligned}} \quad (49)$$

10.4 Stochastic unresolved terrain

Write

$$S_{\text{opt}} = \bar{S}_{\text{opt}} + \xi_{\text{opt}} \quad (50)$$

with

$$\langle \xi_{\text{opt}} \rangle = 0, \quad \langle \xi_{\text{opt}}(\chi) \xi_{\text{opt}}^T(\chi') \rangle = Q_{\text{opt}}(\chi) \delta(\chi - \chi'). \quad (51)$$

The path mean and covariance obey

$$\boxed{\bar{Y}'_{\gamma} = A_K \bar{Y}_{\gamma} + B_{\text{opt}} \bar{S}_{\text{opt}}} \quad (52)$$

and

$$\boxed{C'_Y = A_K C_Y + C_Y A_K^T + B_{\text{opt}} Q_{\text{opt}} B_{\text{opt}}^T}. \quad (53)$$

This is the computational compression used here: unresolved foreground complexity is carried by a 7×7 path covariance rather than by an exhaustive foreground-object ledger.

Rare nonlinear events may be represented separately by a compound-jump term,

$$dY_{\gamma} = (A_K Y_{\gamma} + B_{\text{opt}} \bar{S}_{\text{opt}}) d\chi + B_{\text{opt}} L dW_{\chi} + B_{\text{opt}} J dN_{\chi}. \quad (54)$$

10.5 Observable map and claim boundary

At a fixed physical source endpoint,

$$D_L = (1 + z)^2 D_A \quad (55)$$

gives

$$\delta \ln D_L = 2\zeta_{\gamma} - \kappa. \quad (56)$$

For a Hubble diagram at fixed observed redshift, the accumulated optical state is passed through the existing closed- S^3 source remapping,

$$\frac{\delta D_L}{D_L} = \phi_s + \delta z + \frac{C_K(\chi_s)}{S_K(\chi_s)} \delta \chi - \kappa. \quad (57)$$

Thus Eq. (49) supplies the accumulated physical path state; it does not replace the gauge-invariant luminosity-distance kernel.

The exact closed- S^3 optical propagator, the seven-state reduction and the covariance transport are derived here. The action-normalized seam export $\mathcal{E}_{\Sigma}$, the complete local metric-response map $\mathcal{R}_{g\psi}$, the stochastic covariance Q_{opt} and any rare-event jump law remain open constitutive objects. A future decisive test is therefore not an object-by-object mass reconstruction but a shared cross-observable covariance test: one optical path law must jointly predict luminosity-distance scatter, convergence/shear statistics, redshift perturbations and their cross-correlations.

11 Low-rank cross-observable covariance theorem

Even before the compact full-preimage spatial-current background is numerically available, the optical construction makes a normalization- independent prediction. Let $q \in \mathbb{R}^m$ contain the active physical topographic texture amplitudes and let M_O be the complete linear response from those amplitudes to a fixed observable vector O . Then

$$\boxed{C_O^{\text{topo}} = M_O C_q M_O^{\dagger}} \quad (58)$$

and hence

$$\boxed{\text{rank } C_O^{\text{topo}} \leq \text{rank } C_q \leq m.} \quad (59)$$

For one coherent texture amplitude with nonzero variance and response,

$$C_O^{\text{topo}} = \sigma_q^2 m_O m_O^\dagger, \quad (60)$$

so every pair satisfies

$$\boxed{C_{ii}C_{jj} - |C_{ij}|^2 = 0.} \quad (61)$$

Equivalently, in the noiseless isolated limit,

$$R_1 = \frac{\lambda_1}{\sum_i \lambda_i} = 1, \quad \rho_{21} = \frac{\lambda_2}{\lambda_1} = 0. \quad (62)$$

The rank statement applies to the BHSM topographic contribution, not to raw survey covariance. The test object is

$$\hat{C}_{\text{res}} = \hat{C}_{\text{obs}} - \hat{C}_{\text{std}} - \hat{N}, \quad (63)$$

with the standard-physics and measurement covariances constructed independently of the BHSM rank target.

A residual covariance requiring more than the action-allowed number of independent eigenmodes falsifies that BHSM texture-channel count even if individual pairwise correlations are nonzero. This test does not require the absolute seam-export normalization.

11.1 Coherence assumption and counterexample

Here m counts global random amplitudes after the response and selection are fixed. One local field can contain many independent spatial or temporal amplitudes. Independent increments of one noise type give, in general,

$$C_O = \int d\chi g(\chi)g(\chi)^\dagger.$$

Two nonparallel response vectors at two path segments already give rank two. The white-noise optical evolution therefore does not imply rank one. The minimal test requires $\xi(x) = q \hat{\xi}(x)$ with a fixed shape, or equivalent rank-one global source covariance. Deriving that coherence from the action remains conditional. Zero signal gives rank zero and undefined eigenvalue ratios.

12 Effective representation as a compatibility audit

The role of effective-field language in the present construction is best illustrated by the curvature-aware scalar perturbation audit performed during the development of the R1 cosmological bridge. A conventional Newtonian-gauge effective representation was constructed in order to compare the geometry-first response with familiar late-time perturbation variables. After index and velocity-convention corrections, the system still exhibited a curvature-dependent global constraint-preservation defect. The corresponding flattened system closed numerically, while the curved system did not satisfy the same preservation criterion.

This result is not interpreted as a general failure of effective field theory. Nor is it used as evidence against quantum field theory. It establishes a more limited point: an effective variable set and its equations can provide a useful local or phenomenological representation without necessarily being a globally faithful realization of the complete constrained geometry.

The response of this work was therefore not to modify the effective system until it reproduced the desired cosmological signal. Instead, the defining transport was returned upstream to the physical action reduction:

$$\boxed{\text{physical Hessian} \rightarrow X_2 \rightarrow \mathcal{R}_{g_2} X_2 \rightarrow \text{optical transport} \rightarrow \text{observable kernel.}} \quad (64)$$

The curvature-aware effective equations remain in the project as a compatibility audit. They are not the ontological definition of the BHSM mode.

12.1 A concrete methodological distinction

The practical difference may be summarized as follows:

Representation-first question	Geometry-first question
Which fields should be introduced?	Which perturbations survive the physical constraint and gauge quotient?
Which coefficients fit the observations?	Which coefficients are matrix elements of the retained action?
Is the effective equation stable?	Is the physical Hessian admissible on the common domain?
Can the field reproduce the signal?	Does one reduced state predict multiple observables consistently?
Can missing behavior be absorbed by new terms?	Does a failed prediction reject the proposed reduction?

This difference becomes scientifically meaningful only through prediction and failure. Several candidate reductions tested in this work fail that criterion. A homogeneous two-state mode cannot reproduce the required low-redshift localization. Preregistered foreground number-density, luminosity-weighted, and seam-weighted line-of-sight proxies do not survive held-out transfer. These failures are retained as part of the evidence because the rejected reductions were not subsequently tuned to recover the signal.

12.2 Fields as response coordinates

The geometry-first interpretation does not prohibit field variables. It changes their logical status. Once constraints, boundary data, gauge redundancy, and nonselected directions are removed, a field-like variable may serve as an efficient coordinate on the physical response. Different effective coordinate choices may therefore represent the same underlying reduced geometry.

This observation motivates the optical formulation developed in this work. A photon does not require an object-by-object effective-field inventory of everything along its path. It responds to the total spacetime geometry through the optical tidal matrix. The Jacobi/Sachs system consequently provides a direct bridge from the reduced geometry to observables without assigning fundamental status to every intermediate field representation.

13 Results: Reference Branch R1

R1 is an exploratory reference branch rather than a uniquely derived BHSM cosmology. Its purpose is to demonstrate that the bridge can simultaneously realize late-time scalar activation, a selected physical $n = 2$ response, and near-standard high- n structure growth.

For

$$\Omega_{m0} = 0.31, \quad \Omega_{r0} = 9.0 \times 10^{-5}, \quad \Omega_{k0} = -0.018, \quad \lambda = 1, \quad q = 3,$$

the shooting condition $E(1) = 1$ gives

$$\frac{V_0}{M_{\text{Pl}}^2 H_0^2} = 2.39861.$$

The scalar braiding and no-ghost combination evolve as shown in Table 2.

z	α_B	D_{kin}	Ω_T
10	2.07×10^{-7}	1.64×10^{-6}	0.00194
3	8.60×10^{-5}	6.80×10^{-4}	0.03915
2.1	3.72×10^{-4}	0.00294	0.08060
1.5	0.00122	0.00959	0.14321
1.0	0.00380	0.02988	0.24551
0.8	0.00622	0.04889	0.30787
0.5	0.01346	0.10536	0.43200
0.3	0.02271	0.17740	0.53538
0	0.04940	0.38485	0.70791

Table 2: Reference-branch background evolution.

The scalar equation of state evolves from nearly vacuum-like behavior at high redshift to

$$w_T(z=0) \simeq -0.906,$$

and the background acceleration transition occurs at

$$z_{\text{acc}} \simeq 0.683.$$

Relative to a curved Λ CDM model with identical $H_0, \Omega_{m0}, \Omega_{r0}, \Omega_{k0}$, the R1 expansion rate differs by only order one percent over the late-time interval. The maximum difference in the tabulated range is about 1.7% near $z \sim 0.5$.

13.1 Prospective growth target

In the minimal high- n realization, the direct topographic fifth-force term is negligible on ordinary RSD modes and the dominant growth shift is produced by the altered background history. With identical matter-era normalization,

$$R_{f\sigma_8}(z) = \frac{f\sigma_{8,\text{R1}}}{f\sigma_{8,\Lambda\text{CDM}}}$$

is

The central phenomenological result is therefore not a large generic modification of gravity. It is the combination

$$\begin{aligned} &\text{enhanced horizon-scale } n = 2 \text{ susceptibility} \\ &+ \text{rapid high-}n \text{ recovery} \\ &+ \text{few-percent late-time growth shift} \end{aligned}$$

z	$R_{f\sigma_8}$	suppression
2.1	0.99346	0.65%
1.5	0.98869	1.13%
1.0	0.98164	1.84%
0.8	0.97795	2.20%
0.5	0.97271	2.73%
0.3	0.97168	2.83%
0.0	0.98482	1.52%

Table 3: R1 growth forecast relative to matched curved Λ CDM. These values are reference-branch predictions and become formally preregistered only when reproduced by the canonical code and written to the hashed manifest.

14 Frozen supernova sightline interpretation test

We test whether source structure and independently constrained foregrounds leave a residual following the existing local optical template. This retrospective interpretation test does not instantiate the prospective multi-observable gates. The axis $(211.48^\circ, -12.81^\circ)$, amplitude $A = -0.04321623436$ mag, $z_{\text{ref}} = 0.02$, $z_c = 0.03$, local redshift transfer, R1 background and original sample cuts remain fixed. No residual-fitted bulk velocity or external velocity normalization is used.

14.1 Source and independent foreground construction

Individual epoch/band photometry for all 3,521 released Pantheon+ and DES Hubble-diagram rows is locally reconstructed with the released SALT models. There are 3,480 converged solutions and 41 non-converged local fits, all retained without changing the Hubble-diagram selection. Flux normalization, color, shape and peak epoch are separated; remaining chromatic and phase contrasts are orthogonalized. Band deletion, individual-MJD epoch deletion, phase-block deletion and released spectral/calibration variants provide stability diagnostics. These are local-linear responses, not full nonlinear SNANA/BBC reruns. DES nominal passband proxies and incomplete systematic mappings remain explicitly flagged.

The common distance residual used with the released total covariance is the published standardized distance modulus minus the frozen R1 distance. The raw-fit normalization departure is separate: substituting it into the released distance vector would require a new propagated covariance. Reported epoch variances do not supply the complete model, shared calibration and bias-correction covariance.

Independent density and velocity information comes from the galaxy-only 2MRS neural reconstruction, trained on simulation mocks [6]. Its valid radius is $200 h^{-1}\text{Mpc}$; distant foregrounds remain unmeasured. Observed 2MRS counts [7] are archived separately from the reconstructed matter-density proxy, with magnitude/sky-selection limits. Released dust/host measurements, marginal field errors and a local Green-kernel lensing proxy accompany each sightline. The external spatial error covariance is unavailable.

14.2 Conditional held-out likelihoods

Source/environment nuisance coefficients are learned only on training rows; held-out rows profile the original redshift-bin monopoles and slopes. Every observation of a held-out SN is excluded

from fold training. For training-to-test operator T , the prediction-error covariance is

$$V = C_{vv} + TC_{tt}T^\top - C_{vt}T^\top - TC_{tv}. \quad (65)$$

Pantheon uses its released total covariance. The full DES precision is inverted before selecting rows. No difference of released covariances is treated as an independent PV covariance. Unreleased cross-survey systematic covariance is set to zero only for the explicitly conditional DES transfer; no combined discovery significance follows.

Test	Rows	$\Delta\chi^2$	Mock p
Pantheon, released distances	1208	+2.985	0.247
Pantheon, source/environment	1208	−0.883	0.0725
Pantheon, fixed external PV sensitivity	1208	−4.909	0.0110
DES, released distances	1467	+0.240	0.771
DES, Pantheon nuisance transfer	1467	+1.321	0.590

Table 4: Frozen local prediction versus null. Negative $\Delta\chi^2 = \chi_{\text{frozen}}^2 - \chi_{\text{null}}^2$ favors the prediction. One-sided score tails use 2,000 full-covariance Gaussian mocks, conditional on fixed measured nuisance features. The external PV mean enters with coefficient exactly one; missing joint error propagation precludes a new fully marginalized likelihood.

The primary held-out Pantheon preference is weak and does not transfer to DES (Table 4). Matched same-redshift/different-sky, same-sky/different-redshift and shared-foreground pairs retain their full contrast covariance. The all-Pantheon preference is concentrated in high-density sightlines and varies by survey and sky sector. All-Pantheon cross-fitting reuses the original amplitude-training population, so it is descriptive rather than independent confirmation. Survey/redshift/sector permutations preserve duplicate blocks and the exact redshift law, but their nominal 0.05 tail occurs in 0.09 of the Pantheon null calibration mocks; permutation tails are therefore auxiliary diagnostics.

14.3 PV projection, rank and foreground localization

The saved original low-redshift estimator projects the released PV correction onto +0.033953716 mag, compared with +0.009368488 mag for the independent galaxy-only correction. This demonstrates overlap and correction sensitivity, not identification of the removed component as BHSM optical history. The predictive amplitude remains unchanged. An explicitly *fitted-only* one-scalar diagnostic gives $A_{\text{fit}} = -0.032870 \pm 0.012398$ mag with direction and redshift dependence fixed. The source/environment nuisance space retains 79.0% of the bin-profiled template information, inflating its uncertainty by 1.125. No fitted amplitude is used to improve the predictive scores.

An experimental path-length matrix W_{ik} records shared $10 h^{-1}\text{Mpc}$ Cartesian cells within the unchanged radial shells, using $2 h^{-1}\text{Mpc}$ midpoint steps. The local matrix has 3,931 sampled columns, with numerical ranks 1,268 and 238 for the Pantheon and DES row sets at relative singular tolerance 10^{-7} . Cell-sharing covariance kernels are transformed through the same residual operator as the data. Aggregate shell-adjusted mock tails are no smaller than 0.328 (Pantheon) and 0.531 (DES). A 0.0195 tail in one of 22 overlapping leave-subset scans is not corrected across that family. These tests establish neither foreground-cell localization nor a coherent cross-survey frozen residual. They are not a measurement of the isolated multi-observable topographic covariance or a test of its rank-one gate.

The complete report, compact sightline/results tables, acquisition hashes and script snapshot are retained under `analyses/sn_sightline_tomography_v1` in the repository. The full data-bearing local archive is identified there by SHA-256; a public permanent archive remains open. Complete source-reduction and external-field error propagation are required to resolve the primary physical interpretation.

15 Results: rejected and unestablished mechanisms

The homogeneous two-state response fails the retained localization criterion: state-direction selection does not supply the required low-redshift confinement. The retained development report is `docs/bhsm_los_topographic_transfer.md`. The exact original localization statistic and threshold have not been archived here; this is a reported prior failure, not a newly reproduced quantitative rejection in this package. This rejects that mechanism, not every possible sourced response.

Using the 2MRS release [7], three frozen foreground proxies also fail held-out transfer. The existing results, without refitting, are: Each uses 277 supernovae; the number-density catalog has

Foreground proxy	Permutation p	Held-out $\Delta\chi^2$
2MRS number density	0.076	−4.107958
K-band luminosity	0.092	−4.911833
BHSM seam-charge proxy	0.091	−4.968593

Table 5: All primary classifications are `not_established`. Here $\Delta\chi^2 = \chi_{\text{null}}^2 - \chi_{\text{fixed foreground}}^2$; negative held-out values mean worse prediction.

38600 galaxies, the photometric variants 38601. The latter have 83 calibration and 194 held-out supernovae. These are related development tests, not independent detections. Their permutation values test improvement, not covariance rank. Velocities, lensing and selection remain possible contaminants. None of these proxies establishes an action-normalized local source.

16 Predictions and Falsification Criteria

16.1 Low-redshift calibration

The Pantheon+ data release and cosmological analysis provide the supernova basis for the carried-forward tomographic calibration used here [8, 9]. The earlier scalar-topographic analysis reported

$$A_\mu^\star = -0.0412 \pm 0.009$$

in the bin

$$0.01 < z < 0.03,$$

with best-fit direction

$$(\text{RA}, \text{Dec}) = (211.48^\circ, -12.81^\circ).$$

This corresponds to

$$\frac{\delta D_L}{D_L} = (-0.01895 \pm 0.00414) \cos \theta,$$

and at sufficiently low redshift,

$$\frac{\delta H}{H} = (+0.01895 \pm 0.00414) \cos \theta.$$

These numbers retain the earlier exploratory calibration; its original likelihood is not reconstructed here. Section 14 instead uses the separately frozen later full-covariance amplitude $A = -0.04321623436$ mag. The two calibrations must not be conflated, and neither is retuned by the new interpretation test. The historical numbers normalize only the selected physical $n = 2$ response,

$$A_\mu = \frac{\mathcal{K}_{\text{SN}} S_2}{p_2 C_2},$$

and do not independently determine S_2 , $\mathcal{K}_{\text{SN}}$, or C_2 . The physical distance row is now explicit; one scalar calibration leaves a state-space null direction.

16.2 Curvature

The pre-existing prospective curvature forecast is retained without retuning:

$$\boxed{\Omega_k = -0.018 \pm 0.004}.$$

At $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$,

$$R_H = \frac{c}{H_0 \sqrt{|\Omega_k|}} \simeq 33.15 \text{ Gpc}.$$

The sign prediction is $\Omega_k < 0$ in the usual FLRW curvature convention.

This value is a frozen model forecast, not a current empirical best fit. DESI DR2 BAO results are reported as well described by flat Λ CDM, even while combinations with other probes motivate tests of extensions such as dynamical dark energy [10]. The purpose of retaining -0.018 ± 0.004 is therefore prospective falsification: it must be compared against independent curvature inference without being moved toward the observed result.

16.3 Harmonic hierarchy

The bridge predicts selective $n = 2$ softening with no analogous $n \geq 3$ enhancement. A future source-normalized S^3 $n = 3$ susceptibility comparable to or larger than the physical $n = 2$ susceptibility would reject the minimal harmonic-filter branch. This statement is made at the compact-space harmonic level; comparison to a measured sky quadrupole requires an explicit observer-shell projection.

16.4 Fixed-axis test

A prospective independent supernova sample should first test the carried-forward axis

$$(211.48^\circ, -12.81^\circ)$$

rather than scanning freely over the sky. A free-axis search may be reported secondarily.

16.5 High-n recovery

At large n ,

$$C_n \simeq \chi \ell^2 n^2,$$

so the extra scalar susceptibility falls as n^{-2} . The model therefore predicts a strong separation between the first physical compact-space level and conventional clustering modes.

16.6 Failure conditions

The minimal bridge is disfavored if one or more of the following are established with adequate precision:

1. independent curvature measurements exclude the frozen negative- Ω_k forecast;
2. the source-normalized S^3 $n = 3$ response is comparable to or exceeds physical $n = 2$;
3. an independent low-redshift sample excludes the frozen dipole axis while finding a stable incompatible physical axis;
4. a coherent metric/light-cone dipole of comparable amplitude persists into a redshift regime in which the completed model predicts loss of enhanced susceptibility;
5. ordinary RSD modes show a large scale-independent modification incompatible with the high- n suppression;
6. lensing requires a metric response incompatible with the same completed scalar-metric propagator that controls matter clustering;
7. the selected physical $n = 2$ response cannot satisfy the common-domain constraint or stability requirements.

17 Make-or-break prediction: two prospective gates

The structural protocol is frozen in `preregistration/bhsm_geometry_first_make_or_break_v1.json`. No decisive target data have been evaluated under it. Target releases, masks, bins, covariance and nuisance treatment must be frozen in a dated, hashed design before unblinding. Existing Pantheon+ and 2MRS analyses are development/calibration evidence, not new independent validation.

17.1 Gate A: one shared deterministic state

For N independent measured combinations and fixed two-column kernels,

$$d = FX_2 + \epsilon, \quad \chi^2_{\perp} = \min_X (d - FX)^T \Sigma^{-1} (d - FX). \quad (66)$$

The conventional mean and source response are specified in advance. With rank-2 F , known positive-definite Gaussian Σ and no fitted nuisances, $\chi^2_{\perp} \sim \chi^2_{N-2}$. Thus $N > 2$ overidentifies the state. Cholesky whitening and least squares avoid forming Σ^{-1} . Reject the specific two-state deterministic realization at $p_A < 0.01$.

Both state coordinates vary once across the stacked observations in this conservative consistency test. This does not create a second calibration of an absolute prediction. A theory-selected direction would permit a separately declared one-parameter test. Fitted nuisances, estimated covariance, data-selected axes and nonlinear kernels invalidate the simple $N - 2$ reference law unless incorporated in the predeclared likelihood/simulation.

Independent states or amplitudes per observable, increasing the state dimension, dropping bad bins and changing kernels after inspection are forbidden. The carried-forward equatorial axis is $(211.48^\circ, -12.81^\circ)$; its frame and selection treatment belong in the target design.

17.2 Gate B: one coherent stochastic texture amplitude

For the minimal separable response $O = m_O q$,

$$C_{\text{topo}} = \text{Var}(q) m_O m_O^\dagger, \quad \text{rank } C_{\text{topo}} \leq 1. \quad (67)$$

Rank equals one only for nonzero variance and response. After independently specified standard and measurement-noise treatment, report all signed eigenvalues, $\rho_{21} = \lambda_2/\lambda_1$ when $\lambda_1 > 0$, and $C_{ii}C_{jj} - |C_{ij}|^2$. Observable unit scales are fixed before inspection: exact rank is invariant under invertible scaling, eigenvalue ratios are not.

The primary statistic is the second-largest eigenvalue of the unprojected residual covariance in fixed dimensionless units, with an upper-tail finite-sample calibration. Reject the minimal one-channel stochastic realization at $p_B < 0.01$. Ratios/minors are diagnostics, not extra uncorrected rejection tests. Negative residual eigenvalues remain visible; they may reflect subtraction uncertainty or invalid nuisance modeling.

The executable benchmark uses independent Gaussian samples, an independently fixed total null covariance and 9999 Monte Carlo replicates: $p = (1 + \#\{T_{\text{sim}} \geq T_{\text{obs}}\})/10000$. This validates a restricted fixed null, not the composite family of unknown rank-one covariances. A survey must propagate covariance/loading estimation, selection and sky correlations through its predeclared null simulations, with coverage and power checks.

Increasing m , deleting observables, fitting conventional covariance to enforce low rank, or PSD clipping/projection to manufacture rank one are forbidden. The earlier optical protocol and clipping helper remain historical provenance; this decisive test does not use the helper. A higher-channel model needs an independent action derivation frozen before decisive target inspection and cannot rescue this $m = 1$ realization.

17.3 Separate hypotheses and readiness

The gates are logically distinct; statistical independence is not assumed when data overlap. Each uses a 1% threshold. Under both nulls, the probability of at least one false rejection is bounded by 2%, not 1%. Report both gates and shared-data covariance.

Unknown absolute normalization does not prevent stating these structural restrictions, but applying them requires fixed kernels and a validated nuisance model. One local noise type does not imply one global amplitude. The rank-one test is conditional on the coherent finite-amplitude construction in Section 11.

18 Prospective observational test matrix

The geometry-first workflow becomes scientifically useful only through prospective tests that couple more than one observable to the same reduced physical state. Table 6 summarizes the principal tests without introducing a new fit in this section.

The one-channel rank test has an amplitude-independent null condition,

$$C_{ii}C_{jj} - |C_{ij}|^2 = 0, \quad (68)$$

for every 2×2 minor of an isolated rank-one topographic covariance. Standard astrophysical covariance and measurement noise must be modeled or subtracted before this condition is assessed.

The numerical normalization of every row in Table 6 is not claimed closed at present. The purpose of the table is to specify which pieces of future data must be explained by one common physical response rather than by separate per-observable fits.

Observable	Theory object	Prospective signature	Failure mode
SN / standard candles	$F_{DL}(z)X_2$	Frozen redshift-dependent distance response with common angular structure	Amplitude or redshift evolution inconsistent with the independently selected state
Growth / RSD	$F_{f\sigma_8}(z)X_2$	Same physical state propagated into matter growth	Requires a second independent state not present in the reduction
Weak lensing / Weyl	$F_W(z)X_2$ and optical shear	Cross-observable angular and redshift consistency	Axis or transfer incompatible with the distance response
Photon redshift / convergence / shear	Y_γ	Joint optical-state covariance generated by common geometry	Observed covariance structure incompatible with propagated optical state
Residual covariance	$C_{\text{topo}} = M_O C_q M_O^\dagger$	$\text{rank } C_{\text{topo}} \leq m$	Residual rank robustly exceeds the physically admitted channel count

Table 6: Prospective tests of the geometry-first cosmological response. The tests constrain a common reduced state and common source architecture rather than fitting each observable independently.

19 Discussion and open problems

The novelty claimed is the integrated and audited cosmological construction, not constrained-perturbation mathematics. The regular exclusion of $n = 1$ is inherited geometry. The BHSM proposal is that the surviving response and couplings are matrix elements of a common retained geometric action. Writing a projector alone does not establish that ownership.

The degree-two scalar eigenspace on S^3 has dimension nine; its dipole subspace has three orientation components. A two-dimensional temporal phase space follows after selecting one normalized profile and fixing its axis. The physical selection and population mechanism remain open. The R1 coordinates (ζ, ζ_N) are response coordinates; numerical identification with canonical BHSM (q_2, Π_2) requires a normalization map and consistent transformation of all observable rows.

The propagator uses gravity-scalar reduced coefficients. Dust is transported on that response with zero induced matter response at the anchor. This is not a fully backreacting matter+radiation stability theorem. The curved effective DAE remains an audit with a documented constraint-preservation failure.

The positive p^2 term in a long-wavelength derivative expansion cannot justify extrapolation to arbitrarily high n . The selected filter window and near-standard high- n growth belong to the stated reference realization; the exact parent complement and ultraviolet continuation need separate checks.

Cross-survey source matching, selection, intrinsic alignments, velocities, calibration errors and covariance estimation can mimic extra eigenmodes. The minimal rank bound requires coherent finite latent amplitudes. A survey-aware null calibration and power study with independent nuisance constraints are prerequisites for physical rejection.

Upstream open objects are the normalized compact full-preimage background, complete action-owned local metric/topographic matrix elements, exact seam export, rare-event/source statistics and first-principles absolute amplitudes. Publication before closure would rest on reproducible downstream identities, numerical response checks and restrictions on future realizations. It would not establish empirical success of complete BHSM cosmology.

20 Conclusion

The geometry-first bridge connects constraints and the physical Hessian to a selected $n = 2$ response, induced metric/matter kernels, fixed-observed-redshift luminosity distance and optical means and covariances. Its reference-branch and linear-response assumptions remain explicit. Microscopic source normalization is not closed, and no general objection to EFT/QFT follows.

Two prospective tests put the corresponding realizations at risk: one common two-component deterministic state and at most one coherent stochastic texture amplitude. Rejection at $p < 0.01$ eliminates the relevant realization without extra amplitudes, states, channels or bin selection after inspection. The failed homogeneous and foreground candidates already illustrate this discipline. Action-owned inputs and a frozen survey-specific design remain necessary scientific work.

The frozen supernova sightline test adds a conditional observational check: weak held-out Pantheon preference does not transfer to DES, and aggregate cell-sharing tests do not localize a foreground cause. Independent PV means change the interpretation without licensing a new predictive amplitude. Complete joint source/environment error propagation remains necessary; a coherent topographic residual is not established.

Author Contributions

Norman P. Carberry is the listed author. **TODO (author confirmation):** confirm applicable CRediT roles, including conceptualization, methodology, software, validation, formal analysis, investigation, data curation, writing and visualization.

Funding

TODO (author confirmation): provide an accurate funding statement.

Conflicts of Interest

TODO (author confirmation): disclose conflicts or confirm their absence.

Data Availability Statement

Code, tests, protocols and artifacts are maintained at <https://github.com/ncarberry64/Manuscript-Generation>. The reproducibility guide distinguishes numerical tests from observational reruns. The new sightline analysis includes pinned public-input hashes, compact tables and an exact script snapshot; its full local data-bearing archive is identified by SHA-256. A public permanent archive remains **TODO**. Missing historical foreground-input checksums are not repaired by substituting the new catalogues. The carried-forward predictive SN amplitude remains fixed; the new one-scalar fit is diagnostic only.

AI-use disclosure

OpenAI ChatGPT and OpenAI Codex were materially used for symbolic derivation assistance, code generation/revision, test orchestration, analysis, interpretation of intermediate results and manuscript structuring. AI output was not scientific evidence by itself. Retained claims were transferred into explicit mathematics, executable code, frozen artifacts, automated tests or independently checkable numerical results. The author reviewed the retained work and accepts responsibility. Model/version details should be supplemented from available session records; the final submission requires the author’s review and approval.

Preprint relationship

This manuscript substantially expands and methodologically reframes Preprints.org 202601.1427, cited under its verified published title [1]. That unreviewed preprint is not independent validation.

A Cubic-Galileon Scalar–Metric Bridge

The conditional R1 effective background action, used as a compatibility representation rather than a completed BHSM parent action, is taken to be

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{2} \nabla_\mu T \nabla^\mu T + \frac{1}{\Lambda^3} \square T (\nabla T)^2 - V(T) + \mathcal{L}_m \right].$$

Using

$$X \equiv -\frac{1}{2} \nabla_\mu T \nabla^\mu T,$$

this corresponds to

$$G_2 = X - V(T), \quad G_3 = \frac{2X}{\Lambda^3}, \quad G_4 = \frac{M_{\text{Pl}}^2}{2}, \quad G_5 = 0$$

in the convention $\mathcal{L}_3 = -G_3 \square T$.

The model belongs to the kinetic-gravity-braiding subclass of Horndeski theory [11, 12]. Since G_4 is constant and $G_5 = 0$,

$$M_*^2 = M_{\text{Pl}}^2, \quad \alpha_M = 0, \quad \alpha_T = 0.$$

The scalar–metric mixing is controlled by the braiding function

$$\alpha_B = \frac{2\dot{T}^3}{H M_{\text{Pl}}^2 \Lambda^3},$$

and

$$\alpha_K = \frac{\dot{T}^2}{H^2 M_{\text{Pl}}^2} + 6\alpha_B.$$

In the Bellini–Sawicki organization of Horndeski perturbations the scalar kinetic combination is

$$D_{\text{kin}} = \alpha_K + \frac{3}{2} \alpha_B^2.$$

Their perturbation equations are formulated for a spatially flat background [12]. We therefore use $D_{\text{kin}} > 0$ below as an inherited kinetic diagnostic for the cubic-Galileon sector, not as a complete proof of perturbative stability for the present closed background plus the additional reduced spatial response. A full stability proof requires the constrained gauge-invariant kinetic and gradient matrices, including the exceptional lowest scalar sector identified above.

For numerical work we softly break shift symmetry with

$$V(T) = V_0 - \mu^3 T.$$

The scalar energy density and pressure are

$$\rho_T = \frac{1}{2} \dot{T}^2 + V(T) + \frac{6H\dot{T}^3}{\Lambda^3},$$

$$P_T = \frac{1}{2} \dot{T}^2 - V(T) - \frac{2\dot{T}^2 \ddot{T}}{\Lambda^3}.$$

The shift current

$$J = \dot{T} + \frac{6H\dot{T}^2}{\Lambda^3}$$

obeys

$$\dot{J} + 3HJ = \mu^3.$$

These background expressions are the standard kinetic-gravity-braiding relations specialized to the present G_2, G_3 choice [11].

The closed-FLRW background equation is

$$3M_{\text{Pl}}^2 \left(H^2 + \frac{1}{a^2 R_H^2} \right) = \rho_m + \rho_r + \rho_T.$$

The bridge does not require the bare Horndeski sound speed to become negative. Instead, the BHSM-inspired reduced response is used to lower the effective long-wavelength spatial stiffness while a positive p^2 term stiffens shorter wavelengths. This statement concerns the reduced spatial kernel; it is not, by itself, a complete closed-FLRW stability theorem.

B Exact Closed-Horndeski Kernel for the First Physical Scalar Mode

The previous section uses the Bellini–Sawicki functions to organize the cubic-Galileon background. For the curvature-scale mode, however, the flat-FLRW perturbation formulas are not sufficient. We therefore specialize the non-flat Horndeski quadratic action derived by Akama and Kobayashi [13].

For

$$G_2 = X - V(T), \quad G_3 = \frac{2X}{\Lambda^3}, \quad G_4 = \frac{M_{\text{Pl}}^2}{2}, \quad G_5 = 0,$$

their tensor coefficients reduce exactly to

$$\mathcal{F}_T = \mathcal{G}_T = M_{\text{Pl}}^2.$$

Because G_4 is constant and $G_5 = 0$, the explicit curvature contributions to their Θ and Σ vanish. The remaining coefficients are

$$\Theta = HM_{\text{Pl}}^2 - \frac{\dot{T}^3}{\Lambda^3}$$

and

$$\Sigma = \frac{1}{2}\dot{T}^2 + \frac{12H\dot{T}^3}{\Lambda^3} - 3M_{\text{Pl}}^2 H^2.$$

Equivalently,

$$\Theta = HM_{\text{Pl}}^2 \left(1 - \frac{\alpha_B}{2} \right).$$

Define

$$r(t) \equiv \frac{M_{\text{Pl}}^2 \Sigma}{\Theta^2}, \quad \kappa(t) \equiv \frac{K}{a^2}.$$

The first physical scalar harmonic has alternative-convention index $N = 3$, so

$$k^2 = K(N^2 - 1) = 8K.$$

The two Akama–Kobayashi scalar constraints then reduce to

$$\Sigma \delta n + 8\Theta\kappa\chi + 3\Theta\dot{\zeta} + 5M_{\text{Pl}}^2\kappa\zeta = 0,$$

$$\Theta\delta n - M_{\text{Pl}}^2\dot{\zeta} - M_{\text{Pl}}^2\kappa\chi = 0.$$

They can be solved algebraically:

$$\delta n = \frac{5M_{\text{Pl}}^2}{\Theta(r+8)} \left[\dot{\zeta} - \frac{M_{\text{Pl}}^2\kappa}{\Theta}\zeta \right]$$

and

$$\chi = -\frac{r+3}{\kappa(r+8)}\dot{\zeta} - \frac{5M_{\text{Pl}}^2}{\Theta(r+8)}\zeta.$$

The reduced kinetic coefficient of this $n = 2$ mode is therefore

$$\mathcal{G}_{S,2} = M_{\text{Pl}}^2 \frac{5(r+3)}{r+8}.$$

Using the cubic-Galileon α -functions gives the exact identity

$$r+3 = \frac{D_{\text{kin}}}{2(1-\alpha_B/2)^2},$$

so that

$$\mathcal{G}_{S,2} = M_{\text{Pl}}^2 \frac{5D_{\text{kin}}}{D_{\text{kin}} + 10(1-\alpha_B/2)^2}.$$

Thus, provided $\Theta \neq 0$, positivity of the previously computed D_{kin} is sufficient for positivity of the pure gravity–scalar kinetic coefficient of the first physical closed-universe scalar harmonic.

For completeness, the corresponding non-flat Horndeski gradient coefficient is

$$\mathcal{F}_{S,2} = \frac{1}{a} \frac{d}{dt} \left[\frac{5}{r+8} \left(\frac{aM_{\text{Pl}}^4}{\Theta} \right) \right] - M_{\text{Pl}}^2 + \frac{5}{r+8} \frac{M_{\text{Pl}}^6}{\Theta^2} \frac{K}{a^2}.$$

The repository evaluates this expression as a gravity–scalar diagnostic only. The late-time R1 background contains matter and radiation, whose perturbations introduce additional dynamical variables and source terms. Consequently, $\mathcal{F}_{S,2}$ is not promoted here to a complete matter-era stability or light-cone transfer result.

This section closes an important part of the bridge: the first physical harmonic has an explicit constrained scalar–metric kernel rather than only a geometric identification. The remaining observable calculation is the coupled matter/radiation plus topographic-response system and its gauge-invariant transformation to the metric and velocity variables entering distance, growth, and lensing observables.

C Matter-Sourced Constraint Transfer at n=2

The Round-3 kernel established the pure gravity–scalar constrained dynamics of the first physical closed-universe harmonic. The next step is to introduce minimally coupled matter sources without

importing a flat-space or quasi-static approximation. We use the curvature-aware Newtonian-gauge EFT equations written by Gambino and Pace [14]. Their metric convention is

$$ds^2 = -(1 + 2\Phi)dt^2 + a^2(1 - 2\Psi)\gamma_{ij}dx^i dx^j,$$

with

$$\delta T^0_0 = -\delta\rho, \quad \delta T^0_i = (\rho + p)D_i v.$$

For the present cubic-Galileon subclass,

$$\alpha_M = \alpha_T = \alpha_H = 0, \quad M^2 = M_{\text{Pl}}^2.$$

The curvature-aware equations use the Gleyzes/Gambino–Pace braiding normalization. To distinguish it from the Bellini–Sawicki braiding function used in the previous sections, we write

$$\boxed{\alpha_B^{\text{GP}} = -\frac{1}{2}\alpha_B^{\text{BS}}}.$$

The kineticity is unchanged,

$$\alpha_K^{\text{GP}} = \alpha_K^{\text{BS}},$$

and therefore

$$\alpha_K + 6(\alpha_B^{\text{GP}})^2 = \alpha_K + \frac{3}{2}(\alpha_B^{\text{BS}})^2 = D_{\text{kin}}.$$

For ideal dust and ideal radiation we set the scalar anisotropic stress to zero, so the traceless spatial equation gives

$$\boxed{\Phi = \Psi}.$$

On the physical $n = 2$ harmonic,

$$D^2 Q = -8KQ, \quad (D^2 + 3K)Q = -5KQ.$$

Writing

$$\kappa = \frac{K}{a^2},$$

the sourced 00 equation becomes

$$\begin{aligned} & 6(1 + \alpha_B^{\text{GP}})H\dot{\Phi} + (\alpha_K - 6\alpha_B^{\text{GP}})H^2\dot{\pi} + \left[(6 - \alpha_K + 12\alpha_B^{\text{GP}})H^2 + 10\kappa\right]\Phi \\ & + 6H\left[\frac{\rho + p}{2M_{\text{Pl}}^2} + \dot{H}(1 + \alpha_B^{\text{GP}}) + \kappa\left(1 - \frac{8}{3}\alpha_B^{\text{GP}}\right)\right]\pi = -\frac{\delta\rho}{M_{\text{Pl}}^2}. \end{aligned}$$

The $0i$ equation is

$$2\dot{\Phi} - 2H\alpha_B^{\text{GP}}\dot{\pi} + 2(1 + \alpha_B^{\text{GP}})H\Phi + \left(2\dot{H} + \frac{\rho + p}{M_{\text{Pl}}^2}\right)\pi = -\frac{q}{M_{\text{Pl}}^2},$$

where

$$q = \rho_m v_m + \frac{4}{3}\rho_r v_r, \quad \delta\rho = \rho_m \delta_m + \rho_r \delta_r.$$

These equations form a 2×2 local constraint-transfer system for

$$(\dot{\Phi}, \dot{\pi}).$$

Its determinant is

$$\boxed{\det \mathcal{C} = -2H^2 \left[\alpha_K + 6(\alpha_B^{\text{GP}})^2 \right] = \boxed{-2H^2 D_{\text{kin}}}.}$$

Thus the same positive- D_{kin} branch used in the background and Round-3 kinetic audit guarantees that this sourced rate block is nonsingular.

For the velocity convention above, covariant conservation gives

$$\dot{\delta}_m = \frac{k^2}{a^2} v_m + 3\dot{\Phi}, \quad \dot{v}_m = -\Phi,$$

and

$$\dot{\delta}_r = \frac{4}{3} \frac{k^2}{a^2} v_r + 4\dot{\Phi}, \quad \dot{v}_r = -\Phi - \frac{1}{4} \delta_r,$$

with

$$\frac{k^2}{a^2} = 8\kappa.$$

The repository implements these equations as a local linear transfer matrix with the additional topographic response explicitly switched off. This is deliberately not yet promoted to a globally integrated observable solution. The curvature-aware EFT equations and the Round-3 ADM kernel use different perturbation variables and normalizations; the remaining evolution equations must first be matched analytically and then subjected to a Bianchi/constraint-preservation audit. Only after that compatibility gate is passed will we integrate the full matter-era $n = 2$ system and add the independent reduced topographic operator.

D ADM–EFT Dictionary and Differential-Algebraic Consistency

Rounds 3 and 4 use complementary perturbation languages. Akama and Kobayashi formulate the non-flat Horndeski action in unitary ADM gauge,

$$N = 1 + \delta n, \quad N_i = D_i \chi, \quad g_{ij} = a^2 e^{2\zeta} \gamma_{ij},$$

while Gambino and Pace restore the scalar with

$$t \rightarrow t + \pi$$

and give the curvature-aware field equations in Newtonian gauge [13, 14].

The linear time shift that removes the scalar ADM shift gives

$$\boxed{\pi = -\chi},$$

$$\boxed{\Phi = \delta n - \dot{\pi} = \delta n + \dot{\chi}},$$

and

$$\boxed{\Psi = -\zeta + H\pi = -\zeta - H\chi}.$$

The inverse dictionary is

$$\boxed{\chi = -\pi, \quad \delta n = \Phi + \dot{\pi}, \quad \zeta = H\pi - \Psi.}$$

D.1 Correct Bianchi audit

The first implementation attempt treated the 00 and 0*i* equations as a complete local evolution system and then demanded that the spatial-trace and scalar equations vanish for an arbitrary pointwise state. That test is too strong and uses the constraint equations in the wrong role.

The correct formulation is differential algebraic. The spatial-trace and scalar equations contain the second derivatives of the physical scalar sector and determine

$$(\ddot{\Phi}, \ddot{\pi}),$$

while the 00 and 0*i* equations define constraints on the initial data. Covariant consistency requires those constraints to be preserved by the evolution.

For

$$\alpha_M = \alpha_T = \alpha_H = 0, \quad \Phi = \Psi,$$

the acceleration matrix is

$$\mathcal{A} = \begin{pmatrix} 2 & -2H\alpha_B \\ 6H\alpha_B & H^2\alpha_K \end{pmatrix},$$

where α_B is in the Gambino–Pace/Gleyzes normalization. Its determinant is

$$\det \mathcal{A} = 2H^2 (\alpha_K + 6\alpha_B^2) = 2H^2 D_{\text{kin}} > 0.$$

Thus the second-order scalar–metric evolution block is nonsingular on the same positive- D_{kin} branch.

The repository now integrates the trace/scalar equations together with ideal dust and radiation conservation, starting from data that satisfy the sourced 00 and 0*i* constraints. The Bianchi audit is then the preservation of those two constraints over the interval

$$2.1 \geq z \geq 0.$$

Both raw and term-normalized constraint residuals are evaluated. The topographic reduced-response operator remains switched off throughout this certification.

E Selective Harmonic Response

Let

$$L_c(a) = aR_H\ell, \quad L_c^2 p_n = \ell^2 n(n+2),$$

and write

$$K_n = p_n C_n, \quad C_n = c_s^2 - \chi \left[1 - \ell^2 n(n+2) \right].$$

The physical scalar spectrum begins at $n = 2$. Selective softening of $n = 2$ with stiffening of every propagating $n \geq 3$ requires

$$1 - 8\ell^2 > 0, \quad 1 - 15\ell^2 < 0,$$

so

$$\frac{1}{15} < \ell^2 < \frac{1}{8}.$$

Then

$$C_2 < c_s^2, \quad C_n > c_s^2 \quad (n \geq 3).$$

The formal $n = 1$ coefficient also lies on the soft side, but $n = 1$ is not a propagating scalar degree of freedom in the regular one-field closed-FLRW sector.

If $0 < C_2 \ll 1$,

$$T_2 \simeq \frac{S_2}{p_2 C_2}$$

can be enhanced while higher physical harmonics remain regular. For the next physical level,

$$\frac{T_3}{T_2} = \frac{S_3}{S_2} \frac{8C_2}{15C_3},$$

so

$$\left| \frac{T_3/S_3}{T_2/S_2} \right| = \frac{8C_2}{15C_3} < \frac{8}{15}.$$

For the pure-dipole A_{4i} subspace of the selected $n = 2$ harmonic, the observer-sky pattern is exactly dipolar with geometric radial envelope $\sin 2\psi$.

F Observable Projection

The observable distance perturbation is not identified directly with the scalar amplitude. At linear order it is a light-cone projection of metric, velocity, and scalar perturbations:

$$\frac{\delta D_L}{\bar{D}_L} = \int d\chi \left[W_\Phi \Phi + W_\Psi \Psi + W_v v_\parallel + \dots \right].$$

If the selected $n = 2$ curvature-scale sector admits a physical gauge-invariant metric response, its exact scalar angular dependence can factor as

$$\hat{\mathbf{n}} \cdot \hat{\mathbf{p}},$$

leading schematically to

$$\frac{\delta D_L}{\bar{D}_L} = \epsilon_D(z) \hat{\mathbf{n}} \cdot \hat{\mathbf{p}}.$$

The existence and normalization of $\epsilon_D(z)$ are therefore bridge outputs to be derived, not consequences of the scalar harmonic geometry alone.

For supernovae,

$$\mu = 5 \log_{10} D_L + \text{const},$$

and therefore a small magnitude dipole A_μ corresponds to

$$\frac{\delta D_L}{\bar{D}_L} = \frac{\ln 10}{5} A_\mu \cos \theta.$$

For structure growth we define

$$k^2 \Psi = -4\pi G a^2 \mu(k, a) \rho_m \delta_m, \quad \Phi = \gamma(k, a) \Psi,$$

and

$$\Sigma(k, a) = \frac{\mu(k, a)}{2} [1 + \gamma(k, a)].$$

The matter growth factor obeys

$$D'' + \left(2 + \frac{H'}{H} \right) D' - \frac{3}{2} \Omega_m(a) \mu(k, a) D = 0,$$

with prime denoting $d/d \ln a$. The observable is

$$f\sigma_8(k, z) = \frac{D'}{D} D \sigma_{8, \text{ini}} = D'(k, z) \sigma_{8, \text{ini}}.$$

The quasi-static limit is used only on scales inside the scalar sound horizon [15]. Horizon-scale $n = 2$ observables require the full constrained long-wavelength response and must not be obtained by extrapolating quasi-static formulas into the first physical curvature-scale mode.

G Local BHSM topography on the hyperspherical parent

The smooth $S^3(R_H)$ background and the physical $n = 2$ response do not exhaust the geometry sampled by a photon. We therefore separate the global parent from a local deformation field $\psi(\mathbf{x}, t)$:

$$\gamma_{ij} = a^2 e^{2\psi} \bar{\gamma}_{ij}^{S^3}. \quad (69)$$

To first order its intrinsic-curvature perturbation is

$$\delta^{(3)}R = -\frac{4}{a^2} \left(\bar{\nabla}^2 + \frac{3}{R_H^2} \right) \psi, \quad (70)$$

so the physical manuscript $n = 2$ harmonic carries the shifted factor $8 - 3 = 5$. The local volume rate is

$$H_{\text{local}} = H + \dot{\psi}. \quad (71)$$

The retained BHSM topographic flux candidate is $F_T = \nabla T - B \nabla(\nabla^2 T)$ with $\nabla \cdot F_T = S$. The existing seam-charge candidate is

$$V_\Sigma^2 = \frac{c^2}{8\pi} \int_{\Sigma_\star} H_\Sigma \Delta K_{uu} dA, \quad (72)$$

giving $V_\Sigma^2 = GM/r_\star$ at leading spherical weak-field order. The exact action-level export from this boundary charge into the dynamical ψ source remains open; no normalization is invented here.

G.1 Localized line-of-sight response

For a minimal coherent local-source proxy the accumulated radial perturbation grows only to a coherence distance χ_c and then saturates. The closed- S^3 radial distance term therefore has the normalized shape

$$\mathcal{F}_{\text{loc}}(z) = \frac{[C_K/S_K](\chi) \min(\chi, \chi_c)}{[C_K/S_K](\chi_\star) \min(\chi_\star, \chi_c)}. \quad (73)$$

The numerical example below uses $z_c = 0.03$ only as an illustrative local-coherence proxy, not as a fitted or frozen model parameter.

z	$A_\mu^{\text{loc}} [\text{mag}]$
0.02	-0.041200
0.10	-0.012598
0.30	-0.004435
0.50	-0.002810
0.80	-0.001904
1.00	-0.001605
1.50	-0.001212
2.10	-0.000991

Calibration boundary. The carried-forward low-redshift SN scalar amplitude normalizes the local topographic response in this exploratory decomposition. It cannot simultaneously determine an independent global $n = 2$ amplitude. The global/local split must ultimately be fixed by the common BHSM action or an independent preregistered observable.

H Frozen 2MRS foreground-transfer test

The illustrative local-coherence cutoff is replaced here by a foreground field constructed from the 2MRS galaxy distribution. For supernova i ,

$$\mathcal{F}_i = \frac{C_K(\chi_i)}{S_K(\chi_i)} \sum_s \Delta\chi_{is} \delta_s(\hat{\mathbf{n}}_i),$$

where δ_s is the fixed 10° Gaussian-smoothed density contrast in shell s and $\Delta\chi_{is}$ is only the portion of that shell lying in front of the source. The score is standardized without using supernova residuals. Only one foreground amplitude is fitted.

The frozen analysis used 277 Pantheon+ Hubble-flow supernovae and 38600 2MRS galaxies with $z < 0.05$. Relative to the fixed-axis plus nuisance-control model, adding the foreground score gave

$$\Delta\chi^2 = 1.619359, \quad \beta_F = 0.015286 \pm 0.007846 \text{ mag},$$

with a redshift-blocked permutation value

$$p_{\text{perm}} = 0.076.$$

The preregistered classification is `not_established`.

As a transfer check, the foreground coefficient was calibrated only for $0.01 \leq z \leq 0.03$ and then held fixed for $0.03 < z \leq 0.15$. The held-out change was

$$\Delta\chi_{\text{heldout}}^2 = -4.107958.$$

This test does not identify the physical source of the foreground signal. The 2MRS field is a matter-density proxy, not yet an action-normalized BHSM de-encapsulation map, and conventional peculiar-velocity, lensing and survey selection effects remain alternative explanations.

I Constitutive reduction of the BHSM optical source

The optical transfer equation of Sec. 10 isolates two constitutive maps that were left open: the seam-to-topography export $\mathcal{E}_\Sigma$ and the local metric-response map $\mathcal{R}_{g\psi}$. The retained BHSM calculations already reduce both maps further than a generic phenomenological parameterization.

I.1 Static and moving seam export

The retained scalar-topographic flux obeys

$$\mathbf{F}_T = \nabla T - B\nabla(\nabla^2 T), \quad \nabla \cdot \mathbf{F}_T = S, \quad (74)$$

and the canonical-flux acceleration candidate is

$$\mathbf{g}_T = -Q\mathbf{F}_T. \quad (75)$$

For the stationary spherical neighborhood,

$$QS_{\text{enc}}(r) = \frac{Q_\Sigma}{r^2}, \quad Q_\Sigma = V_\Sigma^2, \quad r \geq r_\star. \quad (76)$$

Thus the static export profile is

$$\mathcal{E}_\Sigma^{(0)}[Q_\Sigma] = \epsilon_\Sigma \frac{Q_\Sigma}{Q} \frac{\Theta(r - r_\star)}{r^2}. \quad (77)$$

The r^{-2} shape is inherited from the conserved-throughput neighborhood law; ϵ_Σ records the still-open common-action orientation.

If de-encapsulation is represented by the time change of this exported source, then for a moving spherical seam

$$\mathcal{E}_\Sigma^{(1)} = \frac{\epsilon_\Sigma}{Q} \left[\dot{Q}_\Sigma \frac{\Theta(r - r_\star)}{r^2} - Q_\Sigma \dot{r}_\star \frac{\delta(r - r_\star)}{r_\star^2} \right]. \quad (78)$$

The second term is a seam-localized contribution generated by boundary motion. Equation (78) is a minimal kinematic continuation, not yet a proof that the retained action selects the time derivative rather than the static source itself.

I.2 Explicit local metric response

Here the conventional spatial-potential sign gives Weyl response $(\Phi + \Psi)/2$. The growth section instead uses the positive spatial-curvature perturbation, giving $\Psi - \Phi$; convert the spatial-potential sign before composing maps. Choose a two-coordinate physical scalar metric basis $h_s = (\Phi, \Psi)^T$ and write

$$\mathcal{A}_g^{\text{loc}} = \begin{pmatrix} a & b \\ b & c \end{pmatrix}, \quad \mathcal{B}_{g\psi}^{\text{loc}} = \begin{pmatrix} u \\ v \end{pmatrix}. \quad (79)$$

For $\Delta_g = ac - b^2 \neq 0$,

$$\mathcal{R}_{g\psi}^{\text{loc}} = -(\mathcal{A}_g^{\text{loc}})^{-1} \mathcal{B}_{g\psi}^{\text{loc}} = \frac{1}{\Delta_g} \begin{pmatrix} bv - cu \\ bu - av \end{pmatrix}. \quad (80)$$

Hence the Weyl/lensing response per unit topographic field is

$$\mathcal{R}_W = \frac{u(b - c) + v(b - a)}{2(ac - b^2)}. \quad (81)$$

The local optical metric response therefore reduces to five projected action matrix elements rather than an arbitrary transfer function.

I.3 Upstream ownership of optical covariance

Let

$$C_J = C_{\text{grav}} + \mathcal{E}_\Sigma C_\Sigma \mathcal{E}_\Sigma^\dagger + C_\xi. \quad (82)$$

This sum assumes mutually uncorrelated source sectors. If they are correlated, their cross-covariance terms must be retained in C_J before propagation; independence is not implied by the sector names. With $\mathcal{G}_\psi = \mathcal{L}_\psi^{-1}$,

$$C_\psi = \mathcal{G}_\psi C_J \mathcal{G}_\psi^\dagger. \quad (83)$$

The unresolved optical forcing covariance is therefore

$$Q_{\text{opt}} = \mathcal{O}_{\text{opt}} \mathcal{R}_{g\psi} \mathcal{G}_{\psi} C_J \mathcal{G}_{\psi}^{\dagger} \mathcal{R}_{g\psi}^{\dagger} \mathcal{O}_{\text{opt}}^{\dagger}. \quad (84)$$

Thus Q_{opt} is not an independent phenomenological covariance matrix. Its rank and cross-observable correlations are inherited from the upstream BSM source statistics.

For a compound-Poisson rare-event population with event rate λ_{Σ} and seam-source amplitude q_{Σ} ,

$$C_{\Sigma}^{\text{shot}} = \lambda_{\Sigma} \langle q_{\Sigma} q_{\Sigma}^{\dagger} \rangle, \quad (85)$$

so

$$Q_{\text{opt}}^{\text{shot}} = \lambda_{\Sigma} \mathcal{M}_{\Sigma} \langle q_{\Sigma} q_{\Sigma}^{\dagger} \rangle \mathcal{M}_{\Sigma}^{\dagger}, \quad \mathcal{M}_{\Sigma} = \mathcal{O}_{\text{opt}} \mathcal{R}_{g\psi} \mathcal{G}_{\psi} \mathcal{E}_{\Sigma}. \quad (86)$$

The remaining numerical closure is finite: evaluate the projected local metric Hessian entries (a, b, c) , the mixed entries (u, v) , the common-action seam-export normalization/orientation, and the low-order statistics of the source population. These quantities replace the previous object-by-object foreground reconstruction problem.

J Action projection contract for the local optical Hessian

The constitutive reduction leaves five local physical coefficients (a, b, c, u, v) . These are not phenomenological parameters. Let e_A and e_{ψ} be the two physical scalar metric basis directions on the common closed-FLRW KKT domain. Then

$$a = \langle e_A, \hat{\mathcal{H}}_{gg} e_A \rangle, \quad (87)$$

$$b = \langle e_A, \hat{\mathcal{H}}_{gg} e_{\psi} \rangle, \quad (88)$$

$$c = \langle e_{\psi}, \hat{\mathcal{H}}_{gg} e_{\psi} \rangle. \quad (89)$$

Thus

$$\mathcal{A}_g^{\text{loc}} = P_g^{\dagger} \hat{\mathcal{H}}_{gg} P_g = \begin{pmatrix} a & b \\ b & c \end{pmatrix}. \quad (90)$$

For the normalized local topographic profile φ_{loc} , the retained mixed second variation

$$B_{g\varphi}[\gamma, \varphi] = \int d\mu_g w \gamma^{\mu\nu} \left[2F''(X)(j \cdot d\varphi) A_{\mu\nu} + 2F'(X) j_{(\mu} \partial_{\nu)} \varphi - F'(X) g_{\mu\nu} (j \cdot d\varphi) \right] \quad (91)$$

defines

$$u = B_{g\varphi}[e_A, \varphi_{\text{loc}}], \quad v = B_{g\varphi}[e_{\psi}, \varphi_{\text{loc}}]. \quad (92)$$

Equivalently,

$$\mathcal{B}_{g\psi}^{\text{loc}} = P_g^{\dagger} \hat{\mathcal{H}}_{g\varphi} \varphi_{\text{loc}} = \begin{pmatrix} u \\ v \end{pmatrix}. \quad (93)$$

If

$$c_{\varphi} = \langle \varphi_{\text{loc}}, \hat{\mathcal{H}}_{\varphi\varphi} \varphi_{\text{loc}} \rangle, \quad (94)$$

the same-domain Schur curvature is

$$k_{\text{loc}} = c_{\varphi} - (\mathcal{B}_{g\psi}^{\text{loc}})^{\dagger} (\mathcal{A}_g^{\text{loc}})^{-1} \mathcal{B}_{g\psi}^{\text{loc}}. \quad (95)$$

The numerical action calculation must return

$$\boxed{a, b, c, u, v, c_\varphi} \quad (96)$$

together with the basis/profile normalizations, $\Delta_g = ac - b^2$, the response residual $\|\mathcal{A}_g \mathcal{R}_{g\psi} + \mathcal{B}_{g\psi}\|$, the Schur curvature, the Legendre-gate status, and the common-domain status. No supernova calibration enters this extraction.

Under an invertible physical metric coordinate change $P_g \rightarrow P_g S$,

$$\mathcal{A}_g \rightarrow S^\dagger \mathcal{A}_g S, \quad \mathcal{B}_{g\psi} \rightarrow S^\dagger \mathcal{B}_{g\psi}, \quad \mathcal{R}_{g\psi} \rightarrow S^{-1} \mathcal{R}_{g\psi}, \quad (97)$$

so the reconstructed physical metric perturbation is basis invariant.

K Spatial-current support gate for the local optical source

The retained phase-specialized metric-topographic second variation gives, on a static block-diagonal background,

$$\boxed{B_{g\varphi}^{\text{spatial}}[\gamma, \varphi] = \int d\mu_g w \gamma^{ij} \left[2F''(X)(j^k \nabla_k \varphi) A_{ij} + 2F'(X) j_{(i} \nabla_{j)} \varphi - F'(X) g_{ij} (j^k \nabla_k \varphi) \right].} \quad (98)$$

Here $j_i = \langle D_i \eta, T \eta \rangle$ is the spatial target Noether current. The static conditions $D_0 \eta = 0$ and $j_0 = 0$ remove the static shift/phase block but do not force j_i to vanish.

It follows immediately that

$$\boxed{j_i = 0 \quad \text{or} \quad \nabla_i \varphi = 0 \quad \implies \quad u = v = 0.} \quad (99)$$

Thus a nonzero spatial target current and nonconstant local texture are necessary, though not sufficient, for this channel to source the local metric response.

For the physical scalar metric basis,

$$u = \int d\mu_g w (e_A)^{ij} \mathcal{S}_{ij}[j, \varphi], \quad (100)$$

$$v = \int d\mu_g w (e_\psi)^{ij} \mathcal{S}_{ij}[j, \varphi]. \quad (101)$$

If $\varphi = q\hat{\varphi}$ has one fixed normalized shape and stochastic amplitude q , then

$$\mathcal{B}_{g\psi}^{\text{loc}} = q b_{\text{loc}} \quad (102)$$

and

$$\boxed{C_B = \text{Var}(q) b_{\text{loc}} b_{\text{loc}}^\dagger}, \quad (103)$$

which has rank at most one. For m independent physical texture channels, the corresponding topographic covariance has rank at most m before additional independent source sectors are included. This gives a cross-observable low-rank prediction for the stochastic optical extension.

L Numerical Implementation and Preregistration Protocol

The numerical pipeline is separated into

$$\boxed{\text{background} \rightarrow \text{physical scalar response} \rightarrow \text{observables} \rightarrow \text{comparison}.}$$

Prediction generation and data comparison are implemented in separate scripts.

L.1 Background system

Let $v = \dot{T}$. Expanding the scalar current equation gives

$$\left(1 + \frac{12Hv}{\Lambda^3}\right) \dot{v} + \frac{6v^2}{\Lambda^3} \dot{H} = \mu^3 - 3H \left(v + \frac{6Hv^2}{\Lambda^3}\right).$$

The Raychaudhuri equation gives

$$\frac{2v^2}{\Lambda^3} \dot{v} - 2M_{\text{Pl}}^2 \dot{H} = \rho_m + \frac{4}{3}\rho_r + v^2 + \frac{6Hv^3}{\Lambda^3} - \frac{2M_{\text{Pl}}^2}{a^2 R_H^2}.$$

At each integration step these equations form a 2×2 linear system for $(\dot{v}, \dot{H})$. The remaining equations are

$$\dot{T} = v, \quad \dot{a} = aH.$$

The Friedmann equation is retained as an independent residual check:

$$3M_{\text{Pl}}^2 \left(H^2 + \frac{1}{a^2 R_H^2} \right) - (\rho_m + \rho_r + \rho_T) = 0.$$

L.2 Dimensionless variables

We integrate in $N = \ln a$ with

$$E = \frac{H}{H_0}, \quad \phi = \frac{T}{M_{\text{Pl}}}, \quad \lambda = \frac{\Lambda^3}{M_{\text{Pl}} H_0^2}, \quad q = \frac{\mu^3}{M_{\text{Pl}} H_0^2}.$$

The exploratory reference branch R1 uses

$$\Omega_{m0} = 0.31, \quad \Omega_{r0} = 9.0 \times 10^{-5}, \quad \Omega_{k0} = -0.018, \quad \lambda = 1, \quad q = 3.$$

The constant V_0 is determined only by the shooting condition $E(a=1) = 1$.

L.3 Initial conditions

The numerical integration begins at

$$z_i = 100.$$

The decaying homogeneous current solution is removed using the matter-era particular solution

$$J_i \simeq \frac{2\mu^3}{9H_i}.$$

The initial scalar velocity is the positive small-velocity root of

$$v_i + \frac{6H_i v_i^2}{\Lambda^3} = J_i.$$

L.4 Growth grid

Before prospective comparison, predictions are generated on the fixed grid

$$k = \{0.02, 0.05, 0.10, 0.15, 0.20\} \text{ Mpc}^{-1}$$

and

$$z = \{0.5, 0.8, 1.0, 1.5, 2.1\}.$$

A deterministic JSON manifest containing the action version, parameters, numerical tolerances, calibration values, and predicted observables is hashed with SHA-256. Any later scientific or numerical revision creates a new manifest version rather than overwriting the previous freeze.

M Reference-Branch Figures




Figure 1: Kinetic-braiding function $\alpha_B(z)$ for Reference Branch R1. The logarithmic vertical scale emphasizes the suppression of the braiding sector toward high redshift.




Figure 2: Scalar energy fraction $\Omega_T(z)$ for R1. The scalar component is subdominant at high redshift and becomes dynamically important only at late times.




Figure 3: Sign of the Schur-response factor $1 - \ell^2 n(n+2)$ for the illustrative filtering value $\ell = 0.32$, matching the canonical code. The physical $n = 2$ branch is softened and $n \geq 3$ stiffened; the displayed $n = 1$ coefficient is exceptional and excluded from propagation.




Figure 4: Prospective R1 growth ratio relative to a matched curved- Λ CDM background with the same early normalization.

References

- [1] Norman P. Carberry. A hyperspherical scalar–topographic framework for late-time cosmological anomalies. Preprints.org, 2026. Posted 20 January 2026, doi:10.20944/preprints202601.1427.v1.
- [2] Claus Kiefer and Tatevik Vardanyan. Power spectrum for perturbations in an inflationary model for a closed universe. *General Relativity and Gravitation*, 54:30, 2022. doi: 10.1007/s10714-022-02918-3.
- [3] Enrico Specogna, Tatevik Vardanyan, William Giarè, and Eleonora Di Valentino. Slow rolling down the curvature: A reassessment of the Planck constraints on ϕ^2 inflation in a closed universe. *Physical Review D*, 113:123526, 2026. doi: 10.1103/dy2d-z3cc.
- [4] Jaiyul Yoo and Fulvio Scaccabarozzi. Unified treatment of the luminosity distance in cosmology. *JCAP*, 09:046, 2016. doi: 10.1088/1475-7516/2016/09/046.
- [5] Ted Pyne and Mark Birkinshaw. The luminosity distance in perturbed FLRW space-times. *Monthly Notices of the Royal Astronomical Society*, 348:581–588, 2004. doi: 10.1111/j.1365-2966.2004.07362.x.
- [6] Robert Lilow, Punyakoti Ganeschaiah Veena, and Adi Nusser. Neural network reconstruction of density and velocity fields from the 2MASS redshift survey, 2024. URL <https://github.com/rililow/2MRS-NeuralNet>.
- [7] John P. Huchra et al. The 2MASS redshift survey—description and data release. *Astrophysical Journal Supplement Series*, 199:26, 2012. doi: 10.1088/0067-0049/199/2/26.
- [8] D. Scolnic et al. The Pantheon+ analysis: The full dataset and light-curve release. *Astrophys. J.*, 938:113, 2022. doi: 10.3847/1538-4357/ac8b7a.
- [9] D. Brout et al. The Pantheon+ analysis: Cosmological constraints. *Astrophys. J.*, 938:110, 2022. doi: 10.3847/1538-4357/ac8e04.
- [10] DESI Collaboration. DESI DR2 results II: Measurements of baryon acoustic oscillations and cosmological constraints. *Physical Review D*, 112:083515, 2025. doi: 10.1103/tr6y-kpc6.
- [11] Cedric Deffayet, Oriol Pujolas, Ignacy Sawicki, and Alexander Vikman. Imperfect dark energy from kinetic gravity braiding. *JCAP*, 10:026, 2010. doi: 10.1088/1475-7516/2010/10/026.
- [12] Emilio Bellini and Ignacy Sawicki. Maximal freedom at minimum cost: linear large-scale structure in general modifications of gravity. *JCAP*, 07:050, 2014. doi: 10.1088/1475-7516/2014/07/050.
- [13] Shingo Akama and Tsutomu Kobayashi. General theory of cosmological perturbations in open and closed universes from the Horndeski action. *Physical Review D*, 99:043522, 2019. doi: 10.1103/PhysRevD.99.043522.
- [14] Serena Gambino and Francesco Pace. Linear perturbations in Horndeski theories with spatial curvature, 2024. Proceedings contribution, MG17.
- [15] Ignacy Sawicki and Emilio Bellini. Limits of quasistatic approximation in modified-gravity cosmologies. *Phys. Rev. D*, 92:084061, 2015. doi: 10.1103/PhysRevD.92.084061.