

Boundary-Mode Topographic Cosmology: A Reduced-Response Bridge from Berger–Hopf Geometry to Late-Time Structure Growth

Norman P. Carberry
Independent Researcher

Working draft — September 2026

Abstract

We develop a late-time cosmological bridge between a scalar topographic effective field theory on compact spatial sections $S^3(R_H)$ and the constraint-reduced response architecture of the Berger–Hopf program. A closed-FLRW gauge audit changes the identification of the relevant infrared carrier: the first nonconstant scalar harmonic in our convention, $n = 1$, maps to the exceptional $N = 2$ sector of standard closed-universe perturbation theory and is nondynamical or pure gauge in the regular single-field covariant scalar sector. The first propagating scalar harmonic is therefore our $n = 2$ ($N = 3$ in the alternative convention). This does not remove the sky-dipole mechanism. A general degree-two scalar harmonic on S^3 contains observer-sky monopole, dipole, and quadrupole pieces, and a pure cross-term subspace produces an exact sky dipole proportional to $\sin[2\psi(z)] \hat{\mathbf{n}} \cdot \hat{\mathbf{p}}$. The reduced Schur/Feshbach response can selectively soften this first physical harmonic while stiffening all propagating $n \geq 3$ modes when $1/15 < \ell^2 < 1/8$. Reference Branch R1 and its frozen numerical manifest are left unchanged because the background and high- n growth calculation do not depend on the identity of the curvature-scale carrier. The theoretical harmonic identification is amended prospectively before comparison with the targeted independent data. The resulting framework is falsifiable through curvature, the fixed dipole axis and radial envelope, compact-harmonic hierarchy, growth, and lensing tests.

1 Introduction

Late-time cosmological tensions and directional residuals motivate tests that are capable of separating local structure, survey geometry, and genuinely new large-scale dynamics. The scalar-topographic program introduced a compact hyperspherical parent $S^3(R_H)$ and a long-wavelength scalar response whose largest observational effect occurs at low redshift [1]. A later covariant formulation placed the topographic scalar in the cubic-Galileon sector of Horndeski theory, preserving second-order time evolution while allowing nontrivial late-time scalar–metric braiding.

The present work asks a narrower question: can the long-wavelength topographic response be represented as the large-scale branch of the same constraint-reduced response logic used in the Berger–Hopf program, without assuming that the full microscopic BHSM construction has already been compactified into cosmology? We answer this at the level of a controlled bridge model.

The central mathematical chain is

$$S[\Phi] \longrightarrow \hat{\mathcal{H}}_{\text{phys}} \longrightarrow \mathcal{K}_{\text{topo}} \longrightarrow \lambda_{\text{topo}} \longrightarrow T_n \longrightarrow \delta g_{\mu\nu} \longrightarrow \{D_L, H, f\sigma_8, \Phi + \Psi\}.$$

The distinction between established input and bridge extension is essential. The S^3 harmonic spectrum and cubic-Galileon background are treated as the cosmological starting point. The Schur/Feshbach reduction is an exact mathematical operation once a physical complement is specified. The identification of its low-energy coefficients with the topographic response is a bridge assumption to be tested, not a declaration that the complete microscopic BHSM action has already been reduced to four-dimensional cosmology.

The paper makes four main contributions. First, it standardizes the S^3 harmonic convention and shows how the observed sky dipole arises from the four-dimensional $n = 1$ eigenspace rather than by identifying a radial mode with an angular dipole. Second, it derives a reduced kernel in which a positive higher-spatial-gradient term can emerge after integrating out a stable complement. Third, it identifies a parameter interval in which the $n = 1$ response is selectively softened while all $n \geq 2$ harmonics are stiffened. Fourth, it gives an executable preregistration protocol for curvature, supernova anisotropy, growth, and lensing tests.

The claim boundary is therefore:

This work does not assume that the scalar-topographic cosmological field has already been derived from the complete microscopic Berger–Hopf action. Exact reductions are distinguished from cross-scale identifications and exploratory reference-branch assumptions. Prospective observables are frozen before comparison with designated independent data.

2 Hyperspherical Parent and Harmonic Convention

We use a closed FLRW geometry

$$ds^2 = -dt^2 + a^2(t)R_H^2 \left[d\psi^2 + \sin^2 \psi d\Omega_2^2 \right],$$

with $a_0 = 1$. Define

$$D_C(z) = c \int_0^z \frac{dz'}{H(z')}, \quad \psi(z) = \frac{D_C(z)}{R_H}.$$

Scalar harmonics satisfy

$$-\nabla_\gamma^2 Y_n = \frac{n(n+2)}{R_H^2} Y_n, \quad n = 0, 1, 2, \dots,$$

so the physical spatial eigenvalue is

$$p_n(a) = \frac{n(n+2)}{a^2 R_H^2}.$$

A common closed-universe convention uses $k^2 = N^2 - 1$, hence

$$\boxed{N = n + 1}.$$

2.1 Exceptional first nonconstant harmonic

For our $n = 1$, $k^2 = 3$ and $N = 2$. The degree-one scalar harmonics on the unit three-sphere obey

$$D_i D_j Q^{(1)} = -\gamma_{ij} Q^{(1)}.$$

Therefore

$$\left(D_i D_j + \frac{k^2}{3} \gamma_{ij} \right) Q^{(1)} = 0.$$

The scalar-derived trace-free metric harmonic is absent. For the regular generally covariant one-scalar sector, the remaining scalar gauge transformations and lapse/shift constraints leave no ordinary propagating scalar degree of freedom in this exceptional sector. Modern gauge-invariant closed-universe analyses likewise take the propagating scalar spectrum to begin at alternative-convention $N \geq 3$ [2, 3].

Accordingly, the minimal bridge no longer uses our $n = 1$ as the physical scalar-metric carrier. An additional boundary degree of freedom could evade this conclusion, but it is not required for the construction below.

2.2 First physical scalar harmonic and exact dipole subspace

The first propagating scalar harmonic is

$$\boxed{n_\star = 2}, \quad \boxed{N_\star = 3}, \quad p_2 = \frac{8}{a^2 R_H^2}.$$

Let the unit $S^3 \subset \mathbb{R}^4$ embedding coordinates be

$$X^A = (\cos \psi, \sin \psi \hat{n}^i).$$

Every real degree-two scalar harmonic can be written

$$T_2 = A_{AB} X^A X^B, \quad A_{AB} = A_{BA}, \quad \delta^{AB} A_{AB} = 0.$$

Writing $c = \cos \psi$, $s = \sin \psi$, $a_i = A_{4i}$, and

$$A_{ij} = S_{ij} - \frac{A_{44}}{3} \delta_{ij}, \quad S_i^i = 0,$$

gives the exact observer-shell decomposition

$$\boxed{T_2 = A_{44} \left(c^2 - \frac{s^2}{3} \right) + 2cs \mathbf{a} \cdot \hat{\mathbf{n}} + s^2 S_{ij} \hat{n}^i \hat{n}^j.}$$

These terms are respectively a sky monopole, dipole, and quadrupole. The pure cross-term subspace

$$A_{44} = 0, \quad S_{ij} = 0, \quad a_i \neq 0$$

therefore gives an exact sky dipole:

$$\boxed{T_{2,\text{dip}} = A \sin(2\psi) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}).}$$

Thus the gauge audit shifts the carrier to the first physical compact-space scalar mode while retaining an exact dipolar observer-sky sector.

3 Constraint-Reduced Response and the Topographic Kernel

Let $\hat{\mathcal{H}}$ be the physical Hessian after constraints and gauge directions have been removed. For a selected topographic branch P and physical complement Q , the exact Schur/Feshbach reduced operator is

$$\mathcal{K} = P \left[\hat{\mathcal{H}} - \hat{\mathcal{H}} Q (Q \hat{\mathcal{H}} Q)^{-1} Q \hat{\mathcal{H}} \right] P.$$

For one scalar harmonic, write the quadratic block as

$$\hat{\mathcal{H}}_n = \begin{pmatrix} A(p) & C(p) \\ C^\dagger(p) & D(p) \end{pmatrix}.$$

The exact reduced stiffness is

$$K_{\text{red}}(p) = A(p) - C(p)D^{-1}(p)C^\dagger(p).$$

A minimal long-wavelength realization is

$$A(p) = Z_0 p, \quad D(p) = M_c^2 + c_c p, \quad C(p) = g\sqrt{p}.$$

Then

$$K_{\text{red}}(p) = Z_0 p - \frac{g^2 p}{M_c^2 + c_c p}.$$

For $c_c p \ll M_c^2$,

$$K_{\text{red}}(p) = \left(Z_0 - \frac{g^2}{M_c^2} \right) p + \frac{g^2 c_c}{M_c^4} p^2 + O(p^3).$$

Thus a positive p^2 stiffness can arise from a coupled second-order parent system after a stable complement is integrated out. In this interpretation the old static operator

$$-\nabla^2 + B\nabla^4$$

is a low-energy spatial derivative expansion, not evidence for a fundamental fourth-order time equation.

Define

$$\chi \equiv \frac{g^2}{M_c^2}, \quad L_c^2 \equiv \frac{c_c}{M_c^2}.$$

After canonical normalization of the leading spatial term, the reduced kernel may be written schematically as

$$K_{\text{topo}}(p) = m_T^2 + (c_s^2 - \chi)p + \chi L_c^2 p^2 + \dots.$$

The signs of the temporal kinetic term and of the full physical gradient matrix must be checked independently; a positive spatial p^2 coefficient by itself is not a complete ghost analysis.

4 Cubic-Galileon Scalar–Metric Bridge

The cosmological parent action is taken to be

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{2} \nabla_\mu T \nabla^\mu T + \frac{1}{\Lambda^3} \square T (\nabla T)^2 - V(T) + \mathcal{L}_m \right].$$

Using

$$X \equiv -\frac{1}{2} \nabla_\mu T \nabla^\mu T,$$

this corresponds to

$$G_2 = X - V(T), \quad G_3 = \frac{2X}{\Lambda^3}, \quad G_4 = \frac{M_{\text{Pl}}^2}{2}, \quad G_5 = 0$$

in the convention $\mathcal{L}_3 = -G_3 \square T$.

The model belongs to the kinetic-gravity-braiding subclass of Horndeski theory [4, 5]. Since G_4 is constant and $G_5 = 0$,

$$M_*^2 = M_{\text{Pl}}^2, \quad \alpha_M = 0, \quad \alpha_T = 0.$$

The scalar–metric mixing is controlled by the braiding function

$$\alpha_B = \frac{2\dot{T}^3}{HM_{\text{Pl}}^2\Lambda^3},$$

and

$$\alpha_K = \frac{\dot{T}^2}{H^2 M_{\text{Pl}}^2} + 6\alpha_B.$$

In the Bellini–Sawicki organization of Horndeski perturbations the scalar kinetic combination is

$$D_{\text{kin}} = \alpha_K + \frac{3}{2}\alpha_B^2.$$

Their perturbation equations are formulated for a spatially flat background [5]. We therefore use $D_{\text{kin}} > 0$ below as an inherited kinetic diagnostic for the cubic-Galileon sector, not as a complete proof of perturbative stability for the present closed background plus the additional reduced spatial response. A full stability proof requires the constrained gauge-invariant kinetic and gradient matrices, including the exceptional lowest scalar sector identified above.

For numerical work we softly break shift symmetry with

$$V(T) = V_0 - \mu^3 T.$$

The scalar energy density and pressure are

$$\rho_T = \frac{1}{2}\dot{T}^2 + V(T) + \frac{6H\dot{T}^3}{\Lambda^3},$$

$$P_T = \frac{1}{2}\dot{T}^2 - V(T) - \frac{2\dot{T}^2\ddot{T}}{\Lambda^3}.$$

The shift current

$$J = \dot{T} + \frac{6H\dot{T}^2}{\Lambda^3}$$

obeys

$$\dot{J} + 3HJ = \mu^3.$$

These background expressions are the standard kinetic-gravity-braiding relations specialized to the present G_2, G_3 choice [4].

The closed-FLRW background equation is

$$3M_{\text{Pl}}^2 \left(H^2 + \frac{1}{a^2 R_H^2} \right) = \rho_m + \rho_r + \rho_T.$$

The bridge does not require the bare Horndeski sound speed to become negative. Instead, the BHSM-inspired reduced response is used to lower the effective long-wavelength spatial stiffness while a positive p^2 term stiffens shorter wavelengths. This statement concerns the reduced spatial kernel; it is not, by itself, a complete closed-FLRW stability theorem.

5 BISM-native transport of the physical topographic mode

The cosmological bridge is formulated first in the mode ontology of the Berger–Hopf Standard Model (BISM), and only secondarily in a four-dimensional effective-field-theory coordinate system. A topographic carrier is therefore not introduced as an additional free bulk scalar. It is an action-owned geometric/boundary mode whose physical dynamics are assigned after the constraint and gauge quotients have been taken.

5.1 Physical tangent space and reduced operator

At a background configuration $z_\star$, define

$$T_{\text{phys}}(z_\star) = \ker DC(z_\star) \cap \mathcal{G}^\perp, \quad (1)$$

where $\mathcal{C}$ is the action-owned constraint map and $\mathcal{G}$ is the space of infinitesimal gauge directions. Let

$$\hat{\mathcal{H}} = P_{\text{phys}} D^2 \mathcal{L}_{\text{KKT}}(z_\star) P_{\text{phys}} \quad (2)$$

be the physical Hessian. For the first admissible scalar-topographic harmonic projector P_2 , define the reduced operator

$$\mathcal{K}_2 = P_2 \left[\hat{\mathcal{H}} - \hat{\mathcal{H}} Q_2 \left(Q_2 \hat{\mathcal{H}} Q_2 \right)^{-1} Q_2 \hat{\mathcal{H}} \right] P_2, \quad (3)$$

where $Q_2 = 1 - P_2$ on the already physical tangent space and the inverse is taken only on the admissible eliminated block.

5.2 Action-owned $n = 2$ envelope transport theorem

Theorem 1. Assume that the cosmological parent envelope is $\mathcal{E}_{\text{cos}} \simeq S^3(R_H)$, that scalar/topographic perturbations are reduced to T_{phys} before physical propagation is assigned, that $n = 0$ is absorbed into the homogeneous background, and that the exceptional regular closed-FLRW scalar $n = 1$ sector is removed by the physical quotient. If the retained $n = 2$ kinetic form is positive and nondegenerate, then the first physical scalar-topographic carrier is an action-owned $n = 2$ envelope mode. Its canonical momentum is generated by the same reduced action and is not a separately postulated bulk field.

Proof. Let $e_2(x)$ be a normalized element of $P_2 T_{\text{phys}}$ and write

$$\delta \Phi_{\text{topo}}(t, x) = q_2(t) e_2(x). \quad (4)$$

The gauge- and constraint-reduced quadratic action has the one-mode form

$$S_2^{(2)} = \frac{1}{2} \int dt a^3 \left[\mathcal{G}_2(t) (D_t q_2)^2 - \mathcal{M}_2^2(t) q_2^2 + 2 \mathcal{J}_2(t) q_2 \right], \quad (5)$$

with coefficients obtained from matrix elements of the reduced physical operator. The conjugate momentum is

$$\Pi_2 = a^3 \mathcal{G}_2 D_t q_2, \quad (6)$$

and variation gives

$$D_t \left(a^3 \mathcal{G}_2 D_t q_2 \right) + a^3 \mathcal{M}_2^2 q_2 = a^3 \mathcal{J}_2. \quad (7)$$

Thus (q_2, Π_2) is the reduced phase-space pair of one action-owned geometric mode. $\square$

5.3 Effective closed-Horndeski matching

The exact closed-Horndeski $n = 2$ result derived elsewhere in this manuscript is used as an effective kinetic representation of the same retained mode, rather than as its microscopic ontology. In the Bellini–Sawicki convention,

$$\mathcal{G}_{S,2} = M_{\text{Pl}}^2 \frac{5D_{\text{kin}}}{D_{\text{kin}} + 10(1 - \alpha_B^{\text{BS}}/2)^2}. \quad (8)$$

Hence $D_{\text{kin}} > 0$ and a nonsingular denominator provide the effective positive-kinetic matching condition used on the reference branch. The EFT Stueckelberg coordinate π is consequently interpreted as a coordinate representative of the retained response of q_2 , not as an additional BHSM degree of freedom.

5.4 Exact observer dipole

A degree-two scalar harmonic on S^3 can be written

$$T_2 = A_{AB} X^A X^B, \quad A_A{}^A = 0. \quad (9)$$

On an observer shell,

$$T_2 = A_{44} \left(c^2 - \frac{s^2}{3} \right) + 2cs \mathbf{a} \cdot \hat{\mathbf{n}} + s^2 S_{ij} \hat{n}^i \hat{n}^j. \quad (10)$$

The pure A_{4i} subspace therefore gives the exact sky dipole

$$\boxed{T_{2,\text{dip}} = A \sin(2\psi) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})}. \quad (11)$$

5.5 Curved-EFT compatibility status

The curvature-aware EFT system remains an important representation-level cross-check. It is not used to define the existence or mode count of the BHSM carrier. The literal curved constraint/evolution equations and the reference background have been transcription-audited, while the global curved-EFT differential-algebraic implementation retains a nonzero constraint-preservation defect. That unresolved object is therefore classified as an EFT compatibility audit rather than as the foundational transport law. The defining bridge evolution is Eq. (7).

6 Metric-response theorem from the reduced BHSM operator

The preceding section identifies the physical $n = 2$ topographic carrier and its reduced phase-space pair $X_2 = (q_2, \Pi_2)^T$. The next required bridge is from that mode to the metric perturbation seen by late-time observables. This bridge follows directly from the mixed physical Hessian and does not require a second independently propagating scalar.

6.1 Coupled quadratic block

On the gauge-fixed physical tangent space, let h denote the retained scalar metric perturbation variables and let X_2 denote the physical topographic mode data. After first-order reduction of any derivative mixing, the linear metric equation can be written

$$\mathcal{A}_g h + \mathcal{B}_{g2} X_2 + J_g = 0, \quad (12)$$

where $\mathcal{A}_g$ is the physical metric Hessian block, $\mathcal{B}_{g2}$ is the metric–topographic mixed block, and J_g is any independent matter/environment source.

The BHSM eta–metric mixed second variation supplies a concrete microscopic source for such a mixed block. In the notation used earlier in the manuscript, that second variation contains terms of the schematic form

$$B_{g\phi}[\gamma, \phi] = \int d\mu_g w \gamma^{\mu\nu} \left[2F''(X)(j \cdot d\phi) A_{\mu\nu} + 2F'(X) j_{(\mu} \partial_{\nu)} \phi - F'(X) g_{\mu\nu} (j \cdot d\phi) \right]. \quad (13)$$

Equation (13) is not asserted to be the complete cosmological propagator. It establishes that the BHSM action already owns a metric–topographic mixing channel from which the reduced response map can be constructed.

6.2 Induced-metric theorem

Theorem 2 (induced metric response). Assume that $\mathcal{A}_g$ is invertible on the chosen physical metric domain. Then the metric response induced by the physical $n = 2$ mode is unique and is given by

$$\boxed{h_2 = h_{\text{src}} + \mathcal{R}_{g2} X_2, \quad \mathcal{R}_{g2} = -\mathcal{A}_g^{-1} \mathcal{B}_{g2},} \quad (14)$$

with

$$h_{\text{src}} = -\mathcal{A}_g^{-1} J_g. \quad (15)$$

No additional physical scalar degree of freedom is introduced.

Proof. Equation (12) is linear on the physical tangent space. Invertibility of $\mathcal{A}_g$ therefore gives

$$h = -\mathcal{A}_g^{-1} J_g - \mathcal{A}_g^{-1} \mathcal{B}_{g2} X_2.$$

The second term is the unique response sourced by the retained topographic mode. Since X_2 already contains the complete reduced canonical data of that mode, the induced metric response carries no additional independent initial datum. $\square$

6.3 Schur-complement consistency

In second-order notation, before first-order phase-space reduction, the quadratic action can be represented by a block operator

$$\mathbb{H}_2 = \begin{pmatrix} \mathcal{A}_g & \mathcal{B}_2 \\ \mathcal{B}_2^\dagger & \mathcal{C}_2 \end{pmatrix}. \quad (16)$$

Eliminating the metric block gives

$$\boxed{\mathcal{K}_2 = \mathcal{C}_2 - \mathcal{B}_2^\dagger \mathcal{A}_g^{-1} \mathcal{B}_2,} \quad (17)$$

which is the same reduced operator that governs the action-owned mode. Thus the metric-response map and the reduced topographic evolution are two faces of the same Schur/Feshbach reduction, rather than independent model assumptions.

6.4 Closed- S^3 harmonic preservation

For an isotropic closed background, $\mathcal{A}_g$ and the action-owned mixed block commute with the background isometry action. Consequently the response operator $\mathcal{R}_{g2}$ preserves the $n = 2$ representation unless an additional anisotropic source is inserted.

Corollary 2.1. For the pure A_{4i} component of the physical $n = 2$ mode, every scalar component of the induced metric response has the observer-sky factor

$$\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}. \quad (18)$$

Hence one may write

$$\Phi_2(z, \hat{\mathbf{n}}) = \Phi_{2,\text{amp}}(z)(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (19)$$

$$\Psi_2(z, \hat{\mathbf{n}}) = \Psi_{2,\text{amp}}(z)(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (20)$$

with amplitudes determined by Eq. (14).

6.5 Observable response operator

Let $\mathcal{L}_{\mathcal{O},z}$ denote the standard linear observation functional for a scalar observable $\mathcal{O}$ at redshift z . Combining the induced metric response with any matter response generated by the same reduced mode gives

$$\delta\mathcal{O}_2 = \mathcal{L}_{\mathcal{O},z} [\mathcal{R}_{g2}X_2 + \mathcal{R}_{m2}X_2]. \quad (21)$$

The observable transfer amplitude is therefore

$$\boxed{\mathcal{A}_{\mathcal{O}}(z) = \mathcal{L}_{\mathcal{O},z} [\mathcal{R}_{g2}(z) + \mathcal{R}_{m2}(z)] X_2(z),} \quad (22)$$

up to independent environmental sources explicitly separated from the single-mode prediction.

Equation (22) is the direct BHSM-native bridge from the reduced $n = 2$ phase-space state to the measurable amplitudes used in the distance, expansion, growth and lensing sectors.

6.6 Scientific closure condition

The remaining numerical closure object is now sharply defined:

$$\boxed{\mathcal{R}_{g2}(z) = -\mathcal{A}_g^{-1}(z)\mathcal{B}_{g2}(z)} \quad (23)$$

together with the matter-response block $\mathcal{R}_{m2}(z)$ on the frozen reference branch. Once these operators are evaluated from the same reduced action, the redshift-dependent observable kernels cease to be independent phenomenological inputs.

A curved four-dimensional EFT realization is accepted as an equivalent representation only if its physical response map agrees with Eq. (14) on the two-dimensional reduced phase space and does not add an additional physical scalar mode.

7 Matter-response theorem driven by the physical $n = 2$ mode

The metric-response theorem fixes the scalar metric perturbation generated by the reduced BHSM phase-space state $X_2 = (q_2, \Pi_2)^T$. Matter and radiation then respond through covariant stress-energy conservation on that induced metric. No additional topographic degree of freedom is introduced at this stage.

7.1 Forced conservation system

Let

$$Y_m = \begin{pmatrix} \delta_m \\ v_m \\ \delta_r \\ v_r \end{pmatrix} \quad (24)$$

denote the linear dust/radiation response variables in a fixed scalar gauge. For a specified background and scalar metric response h_2 , covariant conservation has the linear form

$$\frac{dY_m}{dN} = \mathbb{A}_m(N)Y_m + \mathbb{S}_m(N)h_2(N), \quad (25)$$

where $N = \ln a$. Substituting the metric-response theorem,

$$h_2 = h_{\text{src}} + \mathcal{R}_{g2}X_2, \quad (26)$$

gives

$$\frac{dY_m}{dN} = \mathbb{A}_m Y_m + \mathbb{S}_m h_{\text{src}} + \mathbb{S}_m \mathcal{R}_{g2} X_2. \quad (27)$$

7.2 Matter-response theorem

Theorem 3 (unique matter response). Assume that the linear matter conservation problem is well posed on the chosen redshift interval, with fundamental matrix $\mathbb{U}_m(N, N_i)$ for the homogeneous system $Y'_m = \mathbb{A}_m Y_m$. Then the response sourced by the physical $n = 2$ BHSM mode is

$$Y_{m,2}(N) = \mathbb{U}_m(N, N_i)Y_{m,2}(N_i) + \int_{N_i}^N \mathbb{U}_m(N, N')\mathbb{S}_m(N')\mathcal{R}_{g2}(N')X_2(N')dN'. \quad (28)$$

For a fixed physical initial-data prescription, this defines a linear matter response operator

$$Y_{m,2}(N) = \mathcal{R}_{m2}(N)X_2 \quad (29)$$

in the enlarged sense that $\mathcal{R}_{m2}$ includes the causal history integral generated by the reduced mode.

Proof. Equation (27) is an inhomogeneous first-order linear system. Variation of constants yields Eq. (28). Once the initial-data prescription is fixed as part of the physical branch, every term is linear in the retained mode data and its causal history. Hence the response can be represented by the linear history operator $\mathcal{R}_{m2}$. $\square$

7.3 Adiabatic branch and absence of extra free data

For the single-mode cosmological branch, the matter initial conditions are not treated as arbitrary anisotropic fit functions. The reference branch uses one of the following explicitly declared choices:

1. an adiabatic matter/radiation seed fixed by the background branch;
2. a vanishing anisotropic matter seed, so that all directional response is induced by the BHSM topographic mode;

3. an independently measured environmental source, written separately from the one-mode prediction.

Under the second choice, the directional matter response is completely induced by X_2 . Under the first choice, the isotropic growing mode and the directional $n = 2$ response remain separately identifiable.

7.4 Harmonic preservation

Because $\mathbb{A}_m$, $\mathbb{S}_m$, and $\mathcal{R}_{g2}$ are rotationally covariant on the closed background, a pure physical $n = 2$ source cannot generate a different scalar harmonic at linear order in the absence of an additional anisotropic source.

Corollary 3.1. For the pure A_{4i} component,

$$\delta_m(z, \hat{\mathbf{n}}) = \delta_{m,2}(z)(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (30)$$

$$v_m(z, \hat{\mathbf{n}}) = v_{m,2}(z)(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (31)$$

$$\delta_r(z, \hat{\mathbf{n}}) = \delta_{r,2}(z)(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (32)$$

$$v_r(z, \hat{\mathbf{n}}) = v_{r,2}(z)(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}). \quad (33)$$

Thus the leading matter, metric, distance, growth, and lensing responses share the same one-mode axis.

7.5 Growth-response functional

Let $D(N)$ denote the isotropic reference growth factor and let $\delta D_2(N)$ denote the $n = 2$ directional growth response obtained from $\mathcal{R}_{m2}$. Then, to linear order,

$$\frac{\delta(f\sigma_8)}{f\sigma_8} = \left[\frac{\delta f_2}{f} + \frac{\delta D_2}{D} \right] (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (34)$$

with

$$\delta f_2 = \frac{d}{dN} \left(\frac{\delta D_2}{D} \right). \quad (35)$$

Equation (34) is therefore not an independent ansatz: once $\mathcal{R}_{m2}$ is evaluated, the directional growth kernel is fixed.

7.6 Combined physical transfer operator

Combining the metric and matter responses gives the physical linear response operator

$$\boxed{\mathcal{R}_2 = \begin{pmatrix} \mathcal{R}_{g2} \\ \mathcal{R}_{m2} \end{pmatrix}}. \quad (36)$$

For any scalar linear observable,

$$\boxed{\mathcal{A}_\mathcal{O}(z) = \mathcal{L}_{\mathcal{O},z} \mathcal{R}_2(z) X_2}. \quad (37)$$

This is the closed operator-level bridge from the BHSM action-owned mode to late-time observables.

7.7 Numerical closure object

After Theorems 1–3, the remaining numerical task is no longer the discovery of a missing degree of freedom. It is the evaluation of the already-defined operators

$$\mathcal{G}_2, \quad \mathcal{M}_2^2, \quad \mathcal{J}_2, \quad \mathcal{R}_{g2}, \quad \mathcal{R}_{m2} \quad (38)$$

on the frozen reference branch. Those quantities determine the observable amplitudes in Eq. (37).

The reference-branch curves quoted later in the manuscript should therefore be read as candidate numerical evaluations of this operator chain, pending exact microscopic normalization of every reduced block.

8 Action-provenance gate for a genuinely BHSM-derived cosmological mode

The preceding sections construct a mathematically coherent BHSM-native reduction and observable map. A separate question is whether the physical $n = 2$ cosmological mode has been selected by the presently retained BHSM parent action, rather than introduced as a compatible bridge degree of freedom. The distinction is necessary because later BHSM audits separate harmonic intuition from retained-action provenance.

8.1 Hindsight from the retained BHSM action

The compact closed- S^3 topographic effective theory is useful for harmonic selection, observer projection and late-time response. It is not by itself an action-level completion of the retained BHSM parent theory. In particular, the retained stationary reflection-symmetric branch does not supply a nonzero cap-relative transport momentum. Consequently the existence of the physical $n = 2$ response must be tied to a sourced dynamical branch and a complete physical Hessian before it can be called an unconditional BHSM prediction.

This point does not invalidate the bridge theorems. It fixes their logical status: they are conditional theorems of the BHSM-native reduction until the action-provenance conditions below are discharged.

8.2 Provenance theorem

Theorem 5 (retained-action provenance criterion). Let the BHSM-native cosmological bridge satisfy Theorems 1–4. Suppose in addition that the retained parent action admits a sourced relative-periodic full-preimage solution with all of the following properties:

1. a nonzero conserved cap-relative momentum difference;
2. a globally constraint-reduced physical $\ell = 2$ inertia;
3. a degree-one self-adjoint physical domain with vanishing Green-form leakage at the admissible boundary;
4. a complete gauge-reduced Hessian whose physical $\ell = 2$ block is positive in the kinetic direction and whose mixed metric/topographic block is action owned;
5. a common reduction map that carries the retained parent variables into the cosmological $n = 2$ response without introducing a duplicate scalar degree of freedom.

Then the reduced operator $\mathcal{K}_2$, the metric response $\mathcal{R}_{g2}$, the matter response $\mathcal{R}_{m2}$ and the calibrated observable kernels $\mathcal{F}_\mathcal{O}$ are retained-action BHSM quantities rather than merely compatible bridge constructions.

Proof. Items 1–4 provide the nontrivial dynamical background, physical inertia, self-adjoint domain and complete mixed second variation required to define the physical Hessian on the retained action. Item 5 identifies the bridge variables with that retained physical mode and excludes double counting. The Schur/Feshbach reduction therefore acts on an action-selected physical subspace, so $\mathcal{K}_2$ is action owned. The metric-response theorem then gives $\mathcal{R}_{g2}$ from the same Hessian; covariant conservation gives $\mathcal{R}_{m2}$; and the observation functionals produce $\mathcal{F}_\mathcal{O}$. No additional phenomenological mode selection is required. $\square$

8.3 Fail-closed converse

If any provenance item in Theorem 5 is not established, the manuscript does not promote the resulting cosmological amplitudes to unconditional action-derived BHSM predictions. Instead they are classified as:

$$\boxed{\text{BHSM-native conditional bridge predictions}} \tag{39}$$

on the stated background and transfer realization.

This fail-closed rule is stronger than simply requiring numerical agreement. A successful fit, a stable EFT kernel, or agreement with one cosmological dataset cannot substitute for action provenance.

8.4 Observable-transport precedent

Earlier BHSM work constructed a common observable transport functor for a different physical sector using one declared scheme, one common scale, explicit threshold conventions and one calibration. That construction is used here as a methodological precedent: one common transport map and one declared calibration are allowed, hidden retuning is not.

The cosmological counterpart is the operator chain

$$\mathcal{K}_2 \longrightarrow \mathcal{R}_{g2} \longrightarrow \mathcal{R}_{m2} \longrightarrow \mathcal{F}_\mathcal{O}, \tag{40}$$

with the single calibration of Sec. 17.

8.5 Scientific closure statement

The manuscript therefore separates two levels of closure:

1. *Bridge closure:* Theorems 1–4 define the physical $n = 2$ mode, induced metric and matter response, observable transfer, common axis and single-calibration prediction structure.
2. *Retained-action closure:* Theorem 5 additionally requires the sourced full-preimage branch, physical $\ell = 2$ inertia, self-adjoint domain and complete Hessian to be action selected.

The paper is scientifically complete as a conditional bridge theory once the numerical kernels are evaluated and the claim labels remain explicit. It becomes an unconditional retained-action BHSM cosmology only when the provenance criterion of Theorem 5 is discharged.

9 The single upstream closure object: the physical $n = 2$ mixed Hessian

The finite-kernel inventory shows that the five anisotropic observable kernels are not presently independent numerical objects. Their common upstream input is the physical metric–topographic Hessian in the first admissible $n = 2$ scalar sector. This section reduces that object as far as the retained BHSM action presently permits.

9.1 Retained scalar/topographic operator

The BHSM v5.11 quadratic-operator audit gives the formal scalar/topographic block

$$\mathcal{H}_{\text{ST}} = \begin{pmatrix} -\partial_\rho^2 + 5 & -\mathcal{I}_{\text{mix}} \\ -\mathcal{I}_{\text{mix}}^\dagger & -\Delta_B + 5 \end{pmatrix}, \quad (41)$$

where the collar field and Berger-boundary field require a trace/extension map $\mathcal{I}_{\text{mix}}$ whose full nonhomogeneous domain remains open.

For the physical scalar harmonic $n = 2$ on a unit S^3 section,

$$-\Delta_B e_2 = 8e_2. \quad (42)$$

Let $\lambda_\rho \geq 0$ denote the collar radial eigenvalue on the selected self-adjoint radial domain and let

$$I_2 = \langle u_\rho, \mathcal{I}_{\text{mix}} e_2 \rangle \quad (43)$$

be the projected trace/extension matrix element. Then the retained scalar/topographic $n = 2$ block has the conditional form

$$\boxed{\mathcal{C}_{2,\text{ST}} = \frac{1}{L^2} \begin{pmatrix} \lambda_\rho + 5 & -I_2 \\ -I_2^* & 13 \end{pmatrix}.} \quad (44)$$

Theorem 7 (scalar/topographic $n = 2$ positivity criterion). On a one-mode collar/boundary reduction with $\lambda_\rho \geq 0$, the block in Eq. (44) is positive definite if and only if

$$\boxed{|I_2|^2 < 13(\lambda_\rho + 5).} \quad (45)$$

Proof. The first principal minor is $\lambda_\rho + 5 > 0$. Positivity of the 2×2 Hermitian matrix is therefore equivalent to positivity of its determinant,

$$13(\lambda_\rho + 5) - |I_2|^2 > 0.$$

The common factor L^{-2} is positive. □

9.2 Physical metric–topographic block

Let $\mathcal{A}_g^{(2)}$ denote the gauge-reduced scalar metric Hessian projected onto the same physical $n = 2$ representation, and let $\mathcal{B}_{g2}$ denote the projected metric–topographic mixed block. The complete retained cosmological closure object is

$$\boxed{\mathbb{H}_2^{\text{phys}} = \begin{pmatrix} \mathcal{A}_g^{(2)} & \mathcal{B}_{g2} \\ \mathcal{B}_{g2}^\dagger & \mathcal{C}_{2,\text{ST}} \end{pmatrix}.} \quad (46)$$

The later BHSM action audit already classifies the metric–scalar/topographic mixing as nonzero in source structure but not yet fully evaluated on a common nonhomogeneous domain. Therefore neither $\mathcal{A}_g^{(2)}$ nor $\mathcal{B}_{g2}$ may be replaced by the positive diagnostic coefficients used in the finite gauge-bookkeeping model.

9.3 Single upstream closure theorem

Theorem 8 (one Hessian closes all five observable kernels). Assume that:

1. $\mathbb{H}_2^{\text{phys}}$ is evaluated from the retained action on a common self-adjoint physical domain;
2. $\mathcal{A}_g^{(2)}$ is invertible after the physical gauge quotient;
3. Eq. (45) and the full Schur-positivity conditions hold on the selected branch;
4. the initial-data prescription for the reduced $n = 2$ branch is fixed.

Then the five R1 observable kernels

$$\mathcal{F}_{DL}, \quad \mathcal{F}_H, \quad \mathcal{F}_{DA}, \quad \mathcal{F}_{f\sigma_8}, \quad \mathcal{F}_{\text{lens}} \quad (47)$$

are all determined by this single upstream object together with covariant matter conservation and the observation functionals of Sec. 16.

Proof. The physical Schur reduction gives

$$\mathcal{K}_2 = \mathcal{C}_{2,\text{ST}} - \mathcal{B}_{g2}^\dagger (\mathcal{A}_g^{(2)})^{-1} \mathcal{B}_{g2}, \quad (48)$$

and the induced metric response is

$$\mathcal{R}_{g2} = -(\mathcal{A}_g^{(2)})^{-1} \mathcal{B}_{g2}. \quad (49)$$

The matter-response theorem then fixes $\mathcal{R}_{m2}$ from $\mathcal{R}_{g2}$ and the same $n = 2$ state. Finally each observable kernel is the corresponding linear observation functional acting on the common response. No independent kernel dynamics remain. $\square$

9.4 Correction to the numerical inventory

The existing `code/growth.py` quantity

$$R_{f\sigma_8} = D'_{\text{R1}}/D'_{\Lambda\text{CDM}} \quad (50)$$

is a useful isotropic/reference growth-ratio diagnostic. It is not the directional BHSM-native kernel $\mathcal{F}_{f\sigma_8}$ defined in the present closure protocol. It must therefore not be promoted to that role without the common $n = 2$ response map.

Likewise, the current luminosity-distance, Hubble and lensing expressions in the manuscript specify observation functionals or phenomenological relationships, not action-evaluated anisotropic kernels.

9.5 Scientific consequence

The numerical problem is therefore smaller than “compute five unrelated kernels” but more fundamental than the existing background-growth calculation. The exact remaining action object is

$$\boxed{\left\{ \mathcal{A}_g^{(2)}, \mathcal{B}_{g2}, I_2, \lambda_\rho, \mathcal{D}_{\text{phys}}^{(2)} \right\}}, \quad (51)$$

where $\mathcal{D}_{\text{phys}}^{(2)}$ is the common self-adjoint physical domain.

Once Eq. (51) is action-derived, all five observable kernels follow through Theorems 1–8 without introducing new anisotropic fit parameters.

10 Closed-FLRW metric tangent and the canonical $n = 2$ scalar/topographic candidate

A later BHSM audit identified a normalization conflict in an older critical- fold construction that attempted to use a branch coordinate X as though it were a local curvature perturbation. That issue must not be imported into the cosmological bridge.

10.1 Do not identify the branch coordinate with a local curvature field

In the retained BHSM lineage, the quantity

$$X = H^2 + a^{-2} \quad (52)$$

is an on-shell branch coordinate. It is not a stored independent local scalar field $X(x)$. Moreover, the same value of X occurs in different geometric branches with different scalar-curvature normalization. Therefore an equation of the form

$$\delta R_4 = \delta X \quad (53)$$

is not a branch-independent metric-tangent law.

The cosmological bridge instead uses the direct Fréchet derivative of the four-dimensional scalar curvature with respect to the metric:

$$\boxed{DR_g[k] = \nabla_\mu \nabla_\nu k^{\mu\nu} - \square \text{tr}_g k - R_{\mu\nu} k^{\mu\nu}.} \quad (54)$$

After the scalar gauge quotient, Eq. (54) is projected directly onto the physical closed-FLRW $n = 2$ scalar metric sector. No intermediate $X \rightarrow R_4$ normalization is introduced.

Theorem 9 (cosmological metric-tangent normalization). Let $k_{\mu\nu}^{(2)}$ be a physical $n = 2$ scalar metric perturbation of the closed-FLRW background. Then the action-level curvature variation entering the cosmological metric Hessian is uniquely normalized by

$$\delta R_4^{(2)} = DR_{\bar{g}}[k^{(2)}], \quad (55)$$

modulo the already-quotiented scalar gauge directions. Any auxiliary branch coordinate may be used only after its map to $DR_{\bar{g}}[k^{(2)}]$ has been independently derived on that same branch.

Proof. Equation (54) is the geometric Fréchet derivative of scalar curvature. It depends only on the background metric and its tangent $k_{\mu\nu}^{(2)}$. A branch coordinate has no independent status in this variation unless its derivative with respect to the metric is specified. Therefore the direct metric derivative fixes the normalization without a branch-dependent scalar intermediary. $\square$

10.2 Lowest Robin collar candidate

The v5.7 scalar/topographic boundary problem selects the lowest admissible self-adjoint homogeneous collar mode with Robin zero-flux data. The corresponding constant collar profile has

$$\lambda_\rho = 0. \quad (56)$$

For the nonhomogeneous physical $n = 2$ extension, the minimal harmonic-preserving candidate is to retain this lowest radial profile and extend the S^3 harmonic without changing its normalized boundary amplitude. On that candidate,

$$I_2 = 1. \quad (57)$$

These two values are *candidate domain data*, not a theorem that the full retained-action trace/extension map must take this form.

Substitution into Eq. (44) gives

$$\mathcal{C}_{2,\text{ST}}^{\text{cand}} = \frac{1}{L^2} \begin{pmatrix} 5 & -1 \\ -1 & 13 \end{pmatrix}. \quad (58)$$

Its determinant is

$$\det(L^2 \mathcal{C}_{2,\text{ST}}^{\text{cand}}) = 64 \quad (59)$$

and its eigenvalues are

$$\lambda_{\pm} = \frac{9 \pm \sqrt{17}}{L^2}, \quad (60)$$

both strictly positive.

10.3 What this closes and what remains

Equation (58) closes the scalar/topographic side of the mixed-Hessian calculation on one explicit, self-adjoint, lowest-radial candidate extension. It does not prove that the retained action selects that extension.

The remaining cosmological action object is therefore reduced to

$$\left\{ \mathcal{A}_g^{(2)}, \mathcal{B}_{g2}, \mathcal{D}_{g,\text{phys}}^{(2)} \right\} \quad (61)$$

together with action-level confirmation or rejection of the candidate $(\lambda_\rho, I_2) = (0, 1)$.

The next derivation must evaluate $\mathcal{A}_g^{(2)}$ and $\mathcal{B}_{g2}$ directly from the retained action on the closed-FLRW physical metric tangent. It must not reuse the invalid coefficient-one critical-fold map between X and R_4 .

11 Physical $n = 2$ metric KKT projection

The remaining cosmological action object can be written explicitly in a two-variable scalar metric basis once the momentum constraint and scalar longitudinal gauge have been handled. Later BHSM fixed-boundary work supplies an exact action-derived domain architecture for this reduction, although its radial coefficients belong to a different background and are not imported numerically into cosmology.

11.1 Action-derived domain architecture

For a nonzero scalar harmonic, the induced metric trace before scalar gauge fixing has two components,

$$\Theta = \psi(1) + \frac{1}{4}\square_4 E(1), \quad \mathcal{E} = E(1). \quad (62)$$

The radial lapse A and scalar normal–tangential shift B are not induced metric trace data. The action-derived fixed- h construction proceeds in the following order:

1. vary before gauge fixing;
2. use the momentum equation to eliminate the shift sector;
3. choose the scalar longitudinal gauge $E = 0$;
4. impose the induced-metric Dirichlet trace $\psi(1) = 0$;
5. retain the matcher reaction $\eta_{\text{tr}} = P_\psi(1)$;
6. quotient endpoint-preserving radial diffeomorphisms.

The extended Green current contains the finite matcher pairing

$$\psi_1 \eta_2 - \psi_2 \eta_1, \quad (63)$$

which cancels the canonical-momentum endpoint current on the constrained domain. This gives equal operator and adjoint domains in the corresponding indefinite KKT saddle problem.

Domain proposition. The cosmological $n = 2$ metric calculation adopts this sequence as the *action-derived domain architecture candidate* $\mathcal{D}_{g,\text{phys}}^{(2)}$. The domain architecture is inherited; the background-dependent radial coefficients are not.

11.2 Two-variable physical metric block

After the above reduction, let

$$y_g = \begin{pmatrix} A \\ \psi \end{pmatrix} \quad (64)$$

denote the scalar metric coordinates. The physical metric Hessian projected onto the $n = 2$ harmonic is

$$\boxed{\mathcal{A}_g^{(2)} = \begin{pmatrix} a_{AA} & a_{A\psi} \\ a_{A\psi} & a_{\psi\psi} \end{pmatrix}}, \quad (65)$$

where each entry is an action matrix element evaluated on $\mathcal{D}_{g,\text{phys}}^{(2)}$.

The critical-fold principal operator previously derived in BHSM demonstrates that a lapse–Weyl saddle structure of this type is action generated, but its numerical coefficients are branch specific. They are not substituted into Eq. (65).

11.3 Projected metric–topographic source

Let

$$X_2 = \begin{pmatrix} q_2 \\ \Pi_2 \end{pmatrix}. \quad (66)$$

The mixed physical Hessian defines the source map

$$\boxed{\mathcal{B}_{g2} = \begin{pmatrix} b_{Aq} & b_{A\Pi} \\ b_{\psi q} & b_{\psi\Pi} \end{pmatrix}}. \quad (67)$$

The retained BHSM scalar/topographic action already supplies a nonzero metric–topographic stress-mixing second variation. Projecting that bilinear form onto the physical $n = 2$ mode and the two metric basis directions defines the entries in Eq. (67). In particular, for metric test variation γ and topographic test variation ϕ , the source contains terms of the form

$$B_{g\phi}[\gamma, \phi] = \int d\mu_g w \gamma^{\mu\nu} \left[2F''(X)(j \cdot d\phi) A_{\mu\nu} + 2F'(X) j_{(\mu} \partial_{\nu)} \phi - F'(X) g_{\mu\nu} (j \cdot d\phi) \right]. \quad (68)$$

Thus the remaining coefficients are matrix elements of a stored action variation, not independent phenomenological couplings.

11.4 Explicit response formula

Define

$$\Delta_g = a_{AA} a_{\psi\psi} - a_{A\psi}^2. \quad (69)$$

If $\Delta_g \neq 0$, then

$$(\mathcal{A}_g^{(2)})^{-1} = \frac{1}{\Delta_g} \begin{pmatrix} a_{\psi\psi} & -a_{A\psi} \\ -a_{A\psi} & a_{AA} \end{pmatrix}, \quad (70)$$

and the physical metric response is

$$\boxed{\mathcal{R}_{g2} = -(\mathcal{A}_g^{(2)})^{-1} \mathcal{B}_{g2}}. \quad (71)$$

Therefore all metric response amplitudes are fixed once the seven independent action matrix elements

$$\boxed{a_{AA}, a_{A\psi}, a_{\psi\psi}, b_{Aq}, b_{A\Pi}, b_{\psi q}, b_{\psi\Pi}} \quad (72)$$

are evaluated on the physical domain.

11.5 Schur consistency with the scalar/topographic block

With the candidate scalar/topographic block $\mathcal{C}_{2,\text{ST}}$ from Sec. 10, the reduced mode operator is

$$\boxed{\mathcal{K}_2 = \mathcal{C}_{2,\text{ST}} - \mathcal{B}_{g2}^\dagger (\mathcal{A}_g^{(2)})^{-1} \mathcal{B}_{g2}}. \quad (73)$$

Equations (71) and (73) show that the observable metric response and the mode self-energy correction are controlled by the same finite set of action matrix elements.

11.6 Updated closure count

The action-provenance problem is now reduced to two tasks:

1. confirm that the fixed- h matcher/Green-current architecture above is the correct physical domain for the closed-FLRW $n = 2$ branch;
2. evaluate the seven matrix elements in Eq. (72) from the retained action.

No additional observable-specific anisotropic coefficient is permitted.

12 Exact Closed-Horndeski Kernel for the First Physical Scalar Mode

The previous section uses the Bellini–Sawicki functions to organize the cubic-Galileon background. For the curvature-scale mode, however, the flat-FLRW perturbation formulas are not sufficient. We therefore specialize the non-flat Horndeski quadratic action derived by Akama and Kobayashi [6].

For

$$G_2 = X - V(T), \quad G_3 = \frac{2X}{\Lambda^3}, \quad G_4 = \frac{M_{\text{Pl}}^2}{2}, \quad G_5 = 0,$$

their tensor coefficients reduce exactly to

$$\mathcal{F}_T = \mathcal{G}_T = M_{\text{Pl}}^2.$$

Because G_4 is constant and $G_5 = 0$, the explicit curvature contributions to their Θ and Σ vanish. The remaining coefficients are

$$\Theta = HM_{\text{Pl}}^2 - \frac{\dot{T}^3}{\Lambda^3}$$

and

$$\Sigma = \frac{1}{2}\dot{T}^2 + \frac{12H\dot{T}^3}{\Lambda^3} - 3M_{\text{Pl}}^2H^2.$$

Equivalently,

$$\Theta = HM_{\text{Pl}}^2 \left(1 - \frac{\alpha_B}{2}\right).$$

Define

$$r(t) \equiv \frac{M_{\text{Pl}}^2 \Sigma}{\Theta^2}, \quad \kappa(t) \equiv \frac{K}{a^2}.$$

The first physical scalar harmonic has alternative-convention index $N = 3$, so

$$k^2 = K(N^2 - 1) = 8K.$$

The two Akama–Kobayashi scalar constraints then reduce to

$$\Sigma \delta n + 8\Theta \kappa \chi + 3\Theta \dot{\zeta} + 5M_{\text{Pl}}^2 \kappa \zeta = 0,$$

$$\Theta \delta n - M_{\text{Pl}}^2 \dot{\zeta} - M_{\text{Pl}}^2 \kappa \chi = 0.$$

They can be solved algebraically:

$$\delta n = \frac{5M_{\text{Pl}}^2}{\Theta(r+8)} \left[\dot{\zeta} - \frac{M_{\text{Pl}}^2 \kappa}{\Theta} \zeta \right]$$

and

$$\chi = -\frac{r+3}{\kappa(r+8)} \dot{\zeta} - \frac{5M_{\text{Pl}}^2}{\Theta(r+8)} \zeta.$$

The reduced kinetic coefficient of this $n = 2$ mode is therefore

$$\mathcal{G}_{S,2} = M_{\text{Pl}}^2 \frac{5(r+3)}{r+8}.$$

Using the cubic-Galileon α -functions gives the exact identity

$$r+3 = \frac{D_{\text{kin}}}{2(1-\alpha_B/2)^2},$$

so that

$$\mathcal{G}_{S,2} = M_{\text{Pl}}^2 \frac{5D_{\text{kin}}}{D_{\text{kin}} + 10(1-\alpha_B/2)^2}.$$

Thus, provided $\Theta \neq 0$, positivity of the previously computed D_{kin} is sufficient for positivity of the pure gravity-scalar kinetic coefficient of the first physical closed-universe scalar harmonic.

For completeness, the corresponding non-flat Horndeski gradient coefficient is

$$\mathcal{F}_{S,2} = \frac{1}{a} \frac{d}{dt} \left[\frac{5}{r+8} \left(\frac{aM_{\text{Pl}}^4}{\Theta} \right) \right] - M_{\text{Pl}}^2 + \frac{5}{r+8} \frac{M_{\text{Pl}}^6}{\Theta^2} \frac{K}{a^2}.$$

The repository evaluates this expression as a gravity-scalar diagnostic only. The late-time R1 background contains matter and radiation, whose perturbations introduce additional dynamical variables and source terms. Consequently, $\mathcal{F}_{S,2}$ is not promoted here to a complete matter-era stability or light-cone transfer result.

This section closes an important part of the bridge: the first physical harmonic has an explicit constrained scalar-metric kernel rather than only a geometric identification. The remaining observable calculation is the coupled matter/radiation plus topographic-response system and its gauge-invariant transformation to the metric and velocity variables entering distance, growth, and lensing observables.

13 Matter-Sourced Constraint Transfer at n=2

The Round-3 kernel established the pure gravity-scalar constrained dynamics of the first physical closed-universe harmonic. The next step is to introduce minimally coupled matter sources without importing a flat-space or quasi-static approximation. We use the curvature-aware Newtonian-gauge EFT equations written by Gambino and Pace [7]. Their metric convention is

$$ds^2 = -(1+2\Phi)dt^2 + a^2(1-2\Psi)\gamma_{ij}dx^i dx^j,$$

with

$$\delta T^0_0 = -\delta\rho, \quad \delta T^0_i = (\rho+p)D_i v.$$

For the present cubic-Galileon subclass,

$$\alpha_M = \alpha_T = \alpha_H = 0, \quad M^2 = M_{\text{Pl}}^2.$$

For ideal dust and ideal radiation we set the scalar anisotropic stress to zero, so the traceless spatial equation gives

$$\boxed{\Phi = \Psi}.$$

On the physical $n = 2$ harmonic,

$$D^2 Q = -8KQ, \quad (D^2 + 3K)Q = -5KQ.$$

Writing

$$\kappa = \frac{K}{a^2},$$

the sourced 00 equation becomes

$$\begin{aligned} 6(1 + \alpha_B)H\dot{\Phi} + (\alpha_K - 6\alpha_B)H^2\dot{\pi} + \left[(6 - \alpha_K + 12\alpha_B)H^2 + 10\kappa\right]\Phi \\ + 6H\left[\frac{\rho + p}{2M_{\text{Pl}}^2} + \dot{H}(1 + \alpha_B) + \kappa\left(1 - \frac{8}{3}\alpha_B\right)\right]\pi = -\frac{\delta\rho}{M_{\text{Pl}}^2}. \end{aligned}$$

The $0i$ equation is

$$2\dot{\Phi} - 2H\alpha_B\dot{\pi} + 2(1 + \alpha_B)H\Phi + \left(2\dot{H} + \frac{\rho + p}{M_{\text{Pl}}^2}\right)\pi = -\frac{q}{M_{\text{Pl}}^2},$$

where

$$q = \rho_m v_m + \frac{4}{3}\rho_r v_r, \quad \delta\rho = \rho_m \delta_m + \rho_r \delta_r.$$

These equations form a 2×2 local constraint-transfer system for

$$(\dot{\Phi}, \dot{\pi}).$$

Its determinant is

$$\boxed{\det \mathcal{C} = -2H^2(\alpha_K + 6\alpha_B^2) = -2H^2\left(D_{\text{kin}} + \frac{9}{2}\alpha_B^2\right)}.$$

Thus the same positive- D_{kin} branch used in the background and Round-3 kinetic audit guarantees that this sourced rate block is nonsingular.

For the velocity convention above, covariant conservation gives

$$\dot{\delta}_m = \frac{k^2}{a^2}v_m + 3\dot{\Phi}, \quad \dot{v}_m = -\Phi,$$

and

$$\dot{\delta}_r = \frac{4}{3}\frac{k^2}{a^2}v_r + 4\dot{\Phi}, \quad \dot{v}_r = Hv_r - \Phi - \frac{1}{4}\delta_r,$$

with

$$\frac{k^2}{a^2} = 8\kappa.$$

The repository implements these equations as a local linear transfer matrix with the additional topographic response explicitly switched off. This is deliberately not yet promoted to a globally integrated observable solution. The curvature-aware EFT equations and the Round-3 ADM kernel use different perturbation variables and normalizations; the remaining evolution equations must first be matched analytically and then subjected to a Bianchi/constraint-preservation audit. Only after that compatibility gate is passed will we integrate the full matter-era $n = 2$ system and add the independent reduced topographic operator.

14 Selective Harmonic Response

Let

$$L_c(a) = aR_H\ell, \quad L_c^2 p_n = \ell^2 n(n+2),$$

and write

$$K_n = p_n C_n, \quad C_n = c_s^2 - \chi \left[1 - \ell^2 n(n+2) \right].$$

The physical scalar spectrum begins at $n = 2$. Selective softening of $n = 2$ with stiffening of every propagating $n \geq 3$ requires

$$1 - 8\ell^2 > 0, \quad 1 - 15\ell^2 < 0,$$

so

$$\boxed{\frac{1}{15} < \ell^2 < \frac{1}{8}}.$$

Then

$$C_2 < c_s^2, \quad C_n > c_s^2 \quad (n \geq 3).$$

The formal $n = 1$ coefficient also lies on the soft side, but $n = 1$ is not a propagating scalar degree of freedom in the regular one-field closed-FLRW sector.

If $0 < C_2 \ll 1$,

$$T_2 \simeq \frac{S_2}{p_2 C_2}$$

can be enhanced while higher physical harmonics remain regular. For the next physical level,

$$\frac{T_3}{T_2} = \frac{S_3}{S_2} \frac{8C_2}{15C_3},$$

so

$$\boxed{\left| \frac{T_3/S_3}{T_2/S_2} \right| = \frac{8C_2}{15C_3} < \frac{8}{15}}.$$

For the pure-dipole A_{4i} subspace of the selected $n = 2$ harmonic, the observer-sky pattern is exactly dipolar with geometric radial envelope $\sin 2\psi$.

15 Observable Projection

The observable distance perturbation is not identified directly with the scalar amplitude. At linear order it is a light-cone projection of metric, velocity, and scalar perturbations:

$$\frac{\delta D_L}{\bar{D}_L} = \int d\chi \left[W_\Phi \Phi + W_\Psi \Psi + W_v v_\parallel + \dots \right].$$

If the selected $n = 2$ curvature-scale sector admits a physical gauge-invariant metric response, its exact scalar angular dependence can factor as

$$\hat{\mathbf{n}} \cdot \hat{\mathbf{p}},$$

leading schematically to

$$\frac{\delta D_L}{\bar{D}_L} = \epsilon_D(z) \hat{\mathbf{n}} \cdot \hat{\mathbf{p}}.$$

The existence and normalization of $\epsilon_D(z)$ are therefore bridge outputs to be derived, not consequences of the scalar harmonic geometry alone.

For supernovae,

$$\mu = 5 \log_{10} D_L + \text{const},$$

and therefore a small magnitude dipole A_μ corresponds to

$$\frac{\delta D_L}{D_L} = \frac{\ln 10}{5} A_\mu \cos \theta.$$

For structure growth we define

$$k^2 \Psi = -4\pi G a^2 \mu(k, a) \rho_m \delta_m, \quad \Phi = \gamma(k, a) \Psi,$$

and

$$\Sigma(k, a) = \frac{\mu(k, a)}{2} [1 + \gamma(k, a)].$$

The matter growth factor obeys

$$D'' + \left(2 + \frac{H'}{H}\right) D' - \frac{3}{2} \Omega_m(a) \mu(k, a) D = 0,$$

with prime denoting $d/d \ln a$. The observable is

$$f \sigma_8(k, z) = \frac{D'}{D} D \sigma_{8, \text{ini}} = D'(k, z) \sigma_{8, \text{ini}}.$$

The quasi-static limit is used only on scales inside the scalar sound horizon [8]. Horizon-scale $n = 1$ observables require the full constrained long-wavelength response and must not be obtained by extrapolating quasi-static formulas into the first physical curvature-scale mode.

16 Observable transfer theorem for the physical $n = 2$ mode

The reduced BHSM mode becomes observationally useful only after its action-generated phase-space data are mapped to metric and matter response variables. This section defines that map without introducing an additional degree of freedom.

Let

$$X_2(t) = \begin{pmatrix} q_2(t) \\ \Pi_2(t) \end{pmatrix} \tag{74}$$

denote the physical $n = 2$ phase-space state. Linear response of the retained metric/matter variables can be represented abstractly as

$$Y_2(t) = \mathbb{T}_2(t, t_i) X_2(t_i) + \int_{t_i}^t \mathbb{G}_2(t, t') S_2(t') dt', \tag{75}$$

where Y_2 may contain the two scalar metric potentials, matter density and velocity response, and any auxiliary response coordinate used by the chosen four-dimensional representation. The operators $\mathbb{T}_2$ and $\mathbb{G}_2$ are determined by the reduced action and its source projection; they are not independent phenomenological fields.

16.1 Angular-factorization theorem

Theorem 2 (single-mode observable factorization). Assume (i) the retained perturbation is the pure A_{4i} component of the physical $n = 2$ scalar-topographic mode, (ii) the response is linear, and (iii) the transfer operator is rotationally covariant on the closed background. Then every scalar linear observable $\mathcal{O}$ sourced by this single mode has the leading angular form

$$\boxed{\delta\mathcal{O}(z, \hat{\mathbf{n}}) = \mathcal{A}_{\mathcal{O}}(z) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})}, \quad (76)$$

where the entire observable dependence is contained in the redshift transfer amplitude $\mathcal{A}_{\mathcal{O}}(z)$ and the sky axis $\hat{\mathbf{p}}$ is shared by all observables at leading one-mode order.

Proof. The physical perturbation belongs to one $n = 2$ representation of the isometry group of S^3 . A rotationally covariant linear transfer operator cannot generate a different harmonic representation in the absence of an additional anisotropic source. Restriction of the retained A_{4i} subspace to the observer shell is exactly $T_{2,\text{dip}} \propto \hat{\mathbf{n}} \cdot \hat{\mathbf{p}}$ by Eq. (11). Applying any scalar linear observation functional changes only its radial/time coefficient. Therefore the angular factor is common and Eq. (76) follows. $\square$

The metric response of Sec. 6 and the matter response of Sec. 7 together define the physical response operator $\mathcal{R}_2$. Consequently all linear observable kernels below are functionals of one common reduced phase-space state rather than independent anisotropic fit functions.

16.2 Observable kernels

For each observable define a linear functional $\mathcal{L}_{\mathcal{O},z}$ acting on Eq. (75). Then

$$\mathcal{A}_{\mathcal{O}}(z) = \mathcal{L}_{\mathcal{O},z} \left[\mathbb{T}_2 X_2(t_i) + \int \mathbb{G}_2 S_2 dt' \right]. \quad (77)$$

The principal late-time observables used in this work are represented by

$$\frac{\delta D_L}{D_L} = \mathcal{A}_{D_L}(z) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (78)$$

$$\frac{\delta H}{H} = \mathcal{A}_H(z) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (79)$$

$$\frac{\delta D_A}{D_A} = \mathcal{A}_{D_A}(z) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (80)$$

$$\delta(\Phi + \Psi) = \mathcal{A}_{\text{lens}}(z) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (81)$$

$$\frac{\delta(f\sigma_8)}{f\sigma_8} = \mathcal{A}_{f\sigma_8}(z) (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}), \quad (82)$$

whenever the corresponding observable is evaluated in the linear response regime.

For supernova distance modulus,

$$\delta\mu = \frac{5}{\ln 10} \frac{\delta D_L}{D_L}, \quad (83)$$

so that

$$\boxed{A_\mu(z) = \frac{5}{\ln 10} \mathcal{A}_{D_L}(z)}. \quad (84)$$

Equation (84) supplies the direct bridge between the geometric $n = 2$ carrier and the low-redshift supernova dipole amplitude.

16.3 Shared-axis and amplitude-ratio predictions

Once the transfer kernels are fixed by the reduced action, the amplitudes are not independent fit parameters. For any two linear observables,

$$\frac{\mathcal{A}_{\mathcal{O}_1}(z)}{\mathcal{A}_{\mathcal{O}_2}(z)} = \frac{\mathcal{L}_{\mathcal{O}_1,z}[\mathcal{R}_2]}{\mathcal{L}_{\mathcal{O}_2,z}[\mathcal{R}_2]}, \quad (85)$$

where $\mathcal{R}_2$ denotes the common physical response generated by X_2 and its projected sources. Thus the single-mode branch predicts a common leading axis and linked redshift-dependent amplitude ratios across distance, expansion, growth, and lensing observables.

A statistically persistent axis mismatch between independent observables, after survey-window and calibration effects are controlled, would falsify the pure one-mode realization. Likewise, a leading angular morphology that cannot be represented by the A_{4i} dipole subspace requires additional physical harmonics or rejection of the one-mode branch.

16.4 Higher-mode control

For the harmonic filter

$$C_n = c_s^2 - \chi \left[1 - \ell^2 n(n+2) \right], \quad (86)$$

the interval

$$\frac{1}{15} < \ell^2 < \frac{1}{8} \quad (87)$$

softens the physical $n = 2$ carrier while stiffening every $n \geq 3$ scalar harmonic. The next source-normalized response obeys

$$\left| \frac{T_3/S_3}{T_2/S_2} \right| = \frac{8C_2}{15C_3}. \quad (88)$$

This provides the explicit control parameter for departures from the single-axis theorem.

16.5 Closure criterion for the manuscript

Scientific closure of the observable bridge requires the reduced operator $\mathcal{K}_2$ to determine the transfer kernels entering Eq. (77). A four-dimensional EFT realization is acceptable only if it reproduces the same two-dimensional reduced phase space (q_2, Π_2) and the same observable transfer kernels without adding an independent physical scalar. Reference-branch numerical curves may be quoted before this exact identification only when they are explicitly labeled as reference predictions rather than framework-wide theorems.

17 Single-calibration closure and cross-observable predictions

The preceding theorems reduce the cosmological anisotropic response to one physical $n = 2$ phase-space state X_2 and one common response operator $\mathcal{R}_2$. The remaining overall normalization of the linear one-mode branch can therefore be fixed by one observational calibration without introducing an independent amplitude for each observable.

17.1 Amplitude factorization

For a fixed background branch and a fixed normalized physical mode shape, write

$$X_2(z) = A_2 \hat{X}_2(z), \quad (89)$$

where $\hat{X}_2$ is the unit-normalized reduced solution and A_2 is the single branch amplitude. For any scalar linear observable, Eq. (37) becomes

$$\mathcal{A}_{\mathcal{O}}(z) = A_2 \mathcal{F}_{\mathcal{O}}(z), \quad (90)$$

with

$$\mathcal{F}_{\mathcal{O}}(z) = \mathcal{L}_{\mathcal{O},z} \mathcal{R}_2(z) \hat{X}_2(z). \quad (91)$$

The functions $\mathcal{F}_{\mathcal{O}}$ are fixed by the reduced action and the chosen background branch.

17.2 Single-calibration theorem

Theorem 4 (one calibration fixes all linear amplitudes). Suppose there exists one calibration observable $\mathcal{C}$ at redshift z_* for which $\mathcal{F}_{\mathcal{C}}(z_*) \neq 0$. Then the single-mode amplitude is

$$A_2 = \frac{\mathcal{A}_{\mathcal{C}}^{\text{obs}}(z_*)}{\mathcal{F}_{\mathcal{C}}(z_*)} \quad (92)$$

and every other linear observable is predicted by

$$\mathcal{A}_{\mathcal{O}}(z) = \mathcal{A}_{\mathcal{C}}^{\text{obs}}(z_*) \frac{\mathcal{F}_{\mathcal{O}}(z)}{\mathcal{F}_{\mathcal{C}}(z_*)}. \quad (93)$$

Thus no second anisotropic amplitude parameter is available once the branch, mode normalization and one calibration are fixed.

Proof. Equation (90) is linear in the single branch amplitude A_2 . Solving the calibration equation for A_2 gives Eq. (92). Substitution into the response of any other observable gives Eq. (93). $\square$

17.3 Reference supernova calibration

The current reference branch carries forward the low-redshift supernova distance-modulus calibration

$$A_{\mu}^* = -0.0412 \pm 0.009, \quad 0.01 < z < 0.03, \quad (94)$$

with reference sky axis

$$\text{RA} = 211.48^\circ, \quad \text{Dec} = -12.81^\circ. \quad (95)$$

Because

$$\delta\mu = \frac{5}{\ln 10} \frac{\delta D_L}{D_L}, \quad (96)$$

this corresponds to

$$\left(\frac{\delta D_L}{D_L} \right)_* = -0.01897 \pm 0.00414 \quad (97)$$

for the reference calibration bin.

Equations (94)–(97) are calibration inputs to the reference branch. They are not derived by Theorem 4 and they are not treated as independent calibrations of the other observables.

17.4 Parameter-free amplitude ratios after calibration

For any two observables on the same frozen branch,

$$\boxed{\frac{\mathcal{A}_{\mathcal{O}_1}(z_1)}{\mathcal{A}_{\mathcal{O}_2}(z_2)} = \frac{\mathcal{F}_{\mathcal{O}_1}(z_1)}{\mathcal{F}_{\mathcal{O}_2}(z_2)}} \quad (98)$$

provided the denominator is nonzero. The overall mode amplitude cancels.

This is the principal route from a one-observable calibration to genuine cross-observable prediction. In particular, once the numerical kernels are evaluated, the supernova calibration fixes the absolute anisotropic normalization for

$$\delta H/H, \quad \delta D_A/D_A, \quad \delta(f\sigma_8)/(f\sigma_8), \quad \delta(\Phi + \Psi), \quad (99)$$

without fitting those amplitudes independently.

17.5 Shared-axis theorem as a falsification test

The pure one-mode branch also carries one common axis $\hat{\mathbf{p}}$. Therefore

$$\delta \mathcal{O}_i(z, \hat{\mathbf{n}}) = \mathcal{A}_{\mathcal{O}_i}(z)(\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}) \quad (100)$$

for every scalar linear observable in the branch.

A persistent, statistically resolved disagreement between the axes inferred from independent observables, after survey masks, calibration systematics, and local-structure contamination are controlled, falsifies the pure single- $n = 2$ realization even if individual amplitudes agree.

17.6 Reference-branch versus framework-wide statements

Theorem 4 is structural. The numerical functions $\mathcal{F}_{\mathcal{O}}(z)$ are branch-dependent. Accordingly:

- the existence of one common amplitude A_2 and one common axis is a consequence of the single-mode linear branch;
- the particular numerical values of $\mathcal{F}_{\mathcal{O}}(z)$ on R1 are reference-branch predictions;
- the supernova amplitude in Eq. (94) is an observational calibration input;
- no cross-observable amplitude may be promoted to a framework-wide prediction until its kernel has been evaluated from the reduced operator chain.

17.7 Closure target

The final numerical closure task for the manuscript is now finite and explicit:

$$\boxed{\{\mathcal{F}_{D_L}(z), \mathcal{F}_H(z), \mathcal{F}_{D_A}(z), \mathcal{F}_{f\sigma_8}(z), \mathcal{F}_{\text{lens}}(z)\}_{\text{R1}}}. \quad (101)$$

Once the five frozen-branch kernels in Eq. (101) are evaluated from the already-defined operators, one supernova calibration determines all corresponding anisotropic amplitudes through Eq. (93).

18 Predictions and Falsification Criteria

18.1 Low-redshift calibration

The Pantheon+ data release and cosmological analysis provide the supernova basis for the carried-forward tomographic calibration used here [9, 10]. The earlier scalar-topographic analysis reported

$$A_\mu^* = -0.0412 \pm 0.009$$

in the bin

$$0.01 < z < 0.03,$$

with best-fit direction

$$(\text{RA}, \text{Dec}) = (211.48^\circ, -12.81^\circ).$$

This corresponds to

$$\frac{\delta D_L}{D_L} = (-0.01895 \pm 0.00414) \cos \theta,$$

and at sufficiently low redshift,

$$\frac{\delta H}{H} = (+0.01895 \pm 0.00414) \cos \theta.$$

These numbers are a carried-forward calibration input; the present repository does not independently rerun the Pantheon+ tomographic likelihood. They normalize only the observed $n = 1$ response,

$$A_\mu = \frac{\mathcal{K}_{\text{SN}} S_2}{p_2 C_2},$$

and do not independently determine S_2 , $\mathcal{K}_{\text{SN}}$, or C_2 . Their interpretation as a metric/light-cone response is additionally conditional on resolving the exceptional closed-FLRW $n = 1$ sector.

18.2 Curvature

The pre-existing prospective curvature forecast is retained without retuning:

$$\boxed{\Omega_k = -0.018 \pm 0.004}.$$

At $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$,

$$R_H = \frac{c}{H_0 \sqrt{|\Omega_k|}} \simeq 33.15 \text{ Gpc}.$$

The sign prediction is $\Omega_k < 0$ in the usual FLRW curvature convention.

This value is a frozen model forecast, not a current empirical best fit. DESI DR2 BAO results are reported as well described by flat Λ CDM, even while combinations with other probes motivate tests of extensions such as dynamical dark energy [11]. The purpose of retaining -0.018 ± 0.004 is therefore prospective falsification: it must be compared against independent curvature inference without being moved toward the observed result.

18.3 Harmonic hierarchy

The bridge predicts selective $n = 2$ softening with no analogous $n \geq 3$ enhancement. A future source-normalized S^3 $n = 2$ susceptibility comparable to or larger than the $n = 1$ susceptibility would reject the minimal harmonic-filter branch. This statement is made at the compact-space harmonic level; comparison to a measured sky quadrupole requires an explicit observer-shell projection.

18.4 Fixed-axis test

A prospective independent supernova sample should first test the carried-forward axis

$$(211.48^\circ, -12.81^\circ)$$

rather than scanning freely over the sky. A free-axis search may be reported secondarily.

18.5 High- n recovery

At large n ,

$$C_n \simeq \chi \ell^2 n^2,$$

so the extra scalar susceptibility falls as n^{-2} . The model therefore predicts a strong separation between the first nonconstant compact-space level and conventional clustering modes.

18.6 Failure conditions

The minimal bridge is disfavored if one or more of the following are established with adequate precision:

1. independent curvature measurements exclude the frozen negative- Ω_k forecast;
2. the source-normalized S^3 $n = 2$ response is comparable to or exceeds $n = 1$;
3. an independent low-redshift sample excludes the frozen dipole axis while finding a stable incompatible physical axis;
4. a coherent metric/light-cone dipole of comparable amplitude persists into a redshift regime in which the completed model predicts loss of enhanced susceptibility;
5. ordinary RSD modes show a large scale-independent modification incompatible with the high- n suppression;
6. lensing requires a metric response incompatible with the same completed scalar-metric propagator that controls matter clustering;
7. no gauge-invariant physical metric response exists for the exceptional closed-FLRW harmonic corresponding to the proposed $n = 1$ topographic mode.

19 Numerical Implementation and Preregistration Protocol

The numerical pipeline is separated into

$$\text{background} \rightarrow \text{physical scalar response} \rightarrow \text{observables} \rightarrow \text{comparison}.$$

Prediction generation and data comparison are implemented in separate scripts.

19.1 Background system

Let $v = \dot{T}$. Expanding the scalar current equation gives

$$\left(1 + \frac{12Hv}{\Lambda^3}\right) \dot{v} + \frac{6v^2}{\Lambda^3} \dot{H} = \mu^3 - 3H \left(v + \frac{6Hv^2}{\Lambda^3}\right).$$

The Raychaudhuri equation gives

$$\frac{2v^2}{\Lambda^3} \dot{v} - 2M_{\text{Pl}}^2 \dot{H} = \rho_m + \frac{4}{3}\rho_r + v^2 + \frac{6Hv^3}{\Lambda^3} - \frac{2M_{\text{Pl}}^2}{a^2 R_H^2}.$$

At each integration step these equations form a 2×2 linear system for $(\dot{v}, \dot{H})$. The remaining equations are

$$\dot{T} = v, \quad \dot{a} = aH.$$

The Friedmann equation is retained as an independent residual check:

$$3M_{\text{Pl}}^2 \left(H^2 + \frac{1}{a^2 R_H^2} \right) - (\rho_m + \rho_r + \rho_T) = 0.$$

19.2 Dimensionless variables

We integrate in $N = \ln a$ with

$$E = \frac{H}{H_0}, \quad \phi = \frac{T}{M_{\text{Pl}}}, \quad \lambda = \frac{\Lambda^3}{M_{\text{Pl}} H_0^2}, \quad q = \frac{\mu^3}{M_{\text{Pl}} H_0^2}.$$

The exploratory reference branch R1 uses

$$\Omega_{m0} = 0.31, \quad \Omega_{r0} = 9.0 \times 10^{-5}, \quad \Omega_{k0} = -0.018, \quad \lambda = 1, \quad q = 3.$$

The constant V_0 is determined only by the shooting condition $E(a = 1) = 1$.

19.3 Initial conditions

The numerical integration begins at

$$z_i = 100.$$

The decaying homogeneous current solution is removed using the matter-era particular solution

$$J_i \simeq \frac{2\mu^3}{9H_i}.$$

The initial scalar velocity is the positive small-velocity root of

$$v_i + \frac{6H_i v_i^2}{\Lambda^3} = J_i.$$

19.4 Growth grid

Before prospective comparison, predictions are generated on the fixed grid

$$k = \{0.02, 0.05, 0.10, 0.15, 0.20\} \text{ Mpc}^{-1}$$

and

$$z = \{0.5, 0.8, 1.0, 1.5, 2.1\}.$$

A deterministic JSON manifest containing the action version, parameters, numerical tolerances, calibration values, and predicted observables is hashed with SHA-256. Any later scientific or numerical revision creates a new manifest version rather than overwriting the previous freeze.

20 Finite numerical closure protocol for the R1 observable kernels

After Theorems 1–5, the remaining cosmological calculation is finite. It is not a search for a new degree of freedom or a new phenomenological amplitude. It is the evaluation of the already-defined reduced response kernels on the frozen R1 branch.

20.1 Frozen closure set

Define

$$\mathfrak{K}_{\text{R1}} = \{\mathcal{F}_{DL}(z), \mathcal{F}_H(z), \mathcal{F}_{DA}(z), \mathcal{F}_{f\sigma_8}(z), \mathcal{F}_{\text{lens}}(z)\}. \quad (102)$$

The branch is numerically closed only when each member of $\mathfrak{K}_{\text{R1}}$ is generated from one common reduced state $X_2 = (q_2, \Pi_2)^T$, one common metric response $\mathcal{R}_{g2}$, one common matter response $\mathcal{R}_{m2}$ and one common background realization.

20.2 No-retuning theorem

Theorem 6 (frozen-kernel closure). Let the R1 background, physical-mode normalization, response operators and supernova calibration be fixed. Then no independent anisotropic nuisance amplitude may be introduced into any member of $\mathfrak{K}_{\text{R1}}$. Every observable amplitude is generated by

$$\mathcal{A}_{\mathcal{O}}(z) = A_2 \mathcal{F}_{\mathcal{O}}(z), \quad (103)$$

with the same A_2 fixed by the single calibration theorem.

If any observable requires a second directional amplitude to match data, that result is evidence against the pure one-mode R1 realization rather than permission to refit the branch.

Proof. By Theorem 4, all linear observable amplitudes depend on the same unique single-mode amplitude A_2 . The response operators are fixed functions of the frozen branch. Introducing a second directional amplitude would increase the physical mode count or alter the frozen transfer map. Either operation defines a different model and therefore cannot be part of the R1 closure calculation. $\square$

20.3 Required numerical outputs

A complete R1 closure record shall contain, for every requested redshift bin:

1. the background quantities $E(z)$, $\Omega_T(z)$ and the curvature scale K/a^2 ;
2. the normalized physical state $\hat{X}_2(z)$;
3. the metric response amplitudes $\Phi_{2,\text{amp}}(z)$ and $\Psi_{2,\text{amp}}(z)$;
4. the matter response amplitudes relevant to density and velocity growth;
5. the five observable kernels in $\mathfrak{K}_{\text{R1}}$;
6. the calibrated observable amplitudes obtained from the one supernova normalization;
7. a residual record for every defining reduced equation and transfer identity;
8. immutable hashes of the R1 manifest and the numerical output file.

20.4 Internal consistency identities

The following identities are mandatory checks rather than optional diagnostics:

$$\mathcal{R}_{g2} = -\mathcal{A}_g^{-1}\mathcal{B}_{g2}, \quad (104)$$

$$\mathcal{K}_2 = \mathcal{C}_2 - \mathcal{B}_2^\dagger \mathcal{A}_g^{-1} \mathcal{B}_2, \quad (105)$$

$$Y_{m,2} = \mathcal{R}_{m2} X_2, \quad (106)$$

$$\mathcal{A}_\mathcal{O} = \mathcal{L}_\mathcal{O} \mathcal{R}_2 X_2. \quad (107)$$

Where an effective four-dimensional representation is used, its reduced physical response must agree with these identities on the same two-dimensional phase space. Failure of that equivalence leaves the EFT representation open without invalidating the BHSM-native reduction.

20.5 Prospective-data protocol

The R1 kernels are to be evaluated before comparison with future high-redshift measurements. In particular, growth and lensing kernels must not be recalibrated to DESI, Euclid or Rubin data after those measurements are examined. Those datasets therefore test the forward transfer map rather than participate in its calibration.

20.6 Closure verdicts

The numerical campaign has exactly three allowed verdicts:

R1 kernel closure. All five kernels are produced from the common response chain and satisfy the declared residual and provenance checks.

R1 branch falsified. The common one-mode chain is internally inconsistent or prospective observations reject its linked amplitudes/axis.

Action provenance still conditional. The bridge kernels are numerically well defined, but Theorem 5 has not yet been discharged on the retained BHSM parent action. The outputs remain BHSM-native conditional bridge predictions.

No fourth verdict based on hidden retuning is permitted.

21 Results: Reference Branch R1

R1 is an exploratory reference branch rather than a uniquely derived BHSM cosmology. Its purpose is to demonstrate that the bridge can simultaneously realize late-time scalar activation, a selectively enhanced $n = 1$ response, and near-standard high- n structure growth.

For

$$\Omega_{m0} = 0.31, \quad \Omega_{r0} = 9.0 \times 10^{-5}, \quad \Omega_{k0} = -0.018, \quad \lambda = 1, \quad q = 3,$$

the shooting condition $E(1) = 1$ gives

$$\boxed{\frac{V_0}{M_{\text{Pl}}^2 H_0^2} = 2.39861.}$$

The scalar braiding and no-ghost combination evolve as shown in Table 1.

z	α_B	D_{kin}	Ω_T
10	2.07×10^{-7}	1.64×10^{-6}	0.00194
3	8.60×10^{-5}	6.80×10^{-4}	0.03915
2.1	3.72×10^{-4}	0.00294	0.08060
1.5	0.00122	0.00959	0.14321
1.0	0.00380	0.02988	0.24551
0.8	0.00622	0.04889	0.30787
0.5	0.01346	0.10536	0.43200
0.3	0.02271	0.17740	0.53538
0	0.04940	0.38485	0.70791

Table 1: Reference-branch background evolution.

The scalar equation of state evolves from nearly vacuum-like behavior at high redshift to

$$w_T(z=0) \simeq -0.906,$$

and the background acceleration transition occurs at

$$z_{\text{acc}} \simeq 0.683.$$

Relative to a curved Λ CDM model with identical $H_0, \Omega_{m0}, \Omega_{r0}, \Omega_{k0}$, the R1 expansion rate differs by only order one percent over the late-time interval. The maximum difference in the tabulated range is about 1.7% near $z \sim 0.5$.

21.1 Prospective growth target

In the minimal high- n realization, the direct topographic fifth-force term is negligible on ordinary RSD modes and the dominant growth shift is produced by the altered background history. With identical matter-era normalization,

$$R_{f\sigma_8}(z) = \frac{f\sigma_{8,\text{R1}}}{f\sigma_{8,\Lambda\text{CDM}}}$$

is

z	$R_{f\sigma_8}$	suppression
2.1	0.99346	0.65%
1.5	0.98869	1.13%
1.0	0.98164	1.84%
0.8	0.97795	2.20%
0.5	0.97271	2.73%
0.3	0.97168	2.83%
0.0	0.98482	1.52%

Table 2: R1 growth forecast relative to matched curved Λ CDM. These values are reference-branch predictions and become formally preregistered only when reproduced by the canonical code and written to the hashed manifest.

The central phenomenological result is therefore not a large generic modification of gravity. It is the combination

enhanced horizon-scale $n = 1$ susceptibility
 + rapid high- n recovery
 + few-percent late-time growth shift

22 Reference-Branch Figures




Figure 1: Kinetic-braiding function $\alpha_B(z)$ for Reference Branch R1. The logarithmic vertical scale emphasizes the suppression of the braiding sector toward high redshift.

23 Discussion

The closed-FLRW gauge audit produces a concrete correction. The first nonconstant scalar harmonic in our original notation, $n = 1$, maps to the exceptional $N = 2$ sector of standard closed-universe perturbation theory. Its scalar-derived trace-free tensor harmonic vanishes, and the regular generally covariant one-field scalar sector contains no propagating scalar degree of freedom there. Modern closed-inflation calculations likewise take physical scalar modes to begin at $N \geq 3$ [2, 3].

The natural response is to move the topographic carrier to the first physical scalar level: our $n = 2$, equivalent to $N = 3$. A degree-two harmonic on S^3 is a traceless quadratic form in the $\mathbb{R}^4$ embedding coordinates. Its cross components A_{4i} generate an exact observer-sky dipole

$$T_{2,\text{dip}} \propto \sin 2\psi (\hat{\mathbf{n}} \cdot \hat{\mathbf{p}})$$

without requiring a sky quadrupole.

The reduced-response filter also adapts naturally. The corrected window

$$\frac{1}{15} < \ell^2 < \frac{1}{8}$$




Figure 2: Scalar energy fraction $\Omega_T(z)$ for R1. The scalar component is subdominant at high redshift and becomes dynamically important only at late times.

softens $n = 2$ while stiffening every higher propagating scalar harmonic $n \geq 3$. The formal $n = 1$ coefficient is irrelevant to the regular physical scalar spectrum. The enhanced carrier is therefore the first physical compact-space scalar mode.

The role of BHSM remains structural rather than a completed microscopic derivation. The imported element is the constraint-reduced response logic. The specific mixed block and the source law S_2 remain exploratory until derived from the parent boundary action.

The complete $n = 2$ scalar–metric transfer kernel must still be derived in the closed cubic-Galileon background. $D_{\text{kin}} > 0$ remains an inherited kinetic diagnostic rather than a complete closed-background stability proof. R1 values $q = 3$ and $\lambda = 1$ remain reference choices, and the frozen curvature forecast must not be retuned to observational results.

The R1 numerical background and high- n growth forecast do not depend on this harmonic correction. Their frozen manifest is therefore preserved unchanged, while the theoretical amendment is recorded separately before comparison with the targeted independent datasets.

24 Conclusions

The closed-FLRW gauge analysis resolves the principal ambiguity in the original bridge construction. Our $n = 1$ scalar harmonic maps to the exceptional $N = 2$ sector and is not retained as the ordinary propagating scalar carrier. The first physical scalar level is

$$\boxed{n_\star = 2} \quad (\boxed{N_\star = 3}).$$

This first physical harmonic contains an exact sky-dipole subspace,

$$T_{2,\text{dip}} = A \sin[2\psi(z)](\hat{\mathbf{n}} \cdot \hat{\mathbf{p}}),$$

when only the cross components A_{4i} are populated.




Figure 3: Sign of the Schur-response factor $1 - \ell^2 n(n+2)$ for the illustrative filtering value $\ell = 0.40$. The $n = 1$ branch lies on the softening side while every displayed $n \geq 2$ mode lies on the stiffening side.

The corrected selective-response window is

$$\boxed{\frac{1}{15} < \ell^2 < \frac{1}{8}},$$

within which $n = 2$ is softened and every higher propagating mode $n \geq 3$ is stiffened. The next-level source-normalized response obeys

$$\left| \frac{T_3/S_3}{T_2/S_2} \right| = \frac{8C_2}{15C_3} < \frac{8}{15}.$$

The carried-forward supernova amplitude now calibrates the physical $n = 2$ dipole response. The R1 background and high- n growth predictions remain unchanged.

The revised signature is therefore

$$\boxed{\begin{aligned} &\text{first physical } S^3 \text{ scalar mode } (n = 2) \text{ enhanced} \\ &\oplus \text{ fixed-axis dipole subspace with } \sin 2\psi \text{ envelope} \\ &\quad \oplus \text{ suppressed } n \geq 3 \text{ susceptibility} \\ &\quad \oplus \text{ near-GR high-}k \text{ growth.} \end{aligned}}$$

The next calculation is the full gauge-invariant $n = 2$ scalar-metric transfer matrix in the closed cubic-Galileon background.

References

- [1] Norman P. Carberry. A hyperspherical scalar-topographic framework for late-time cosmological anomalies. Preprints.org, 2026. Posted 20 January 2026, doi:10.20944/preprints202601.1427.v1.




Figure 4: Prospective R1 growth ratio relative to a matched curved- Λ CDM background with the same early normalization.

- [2] Claus Kiefer and Tatevik Vardanyan. Power spectrum for perturbations in an inflationary model for a closed universe. *General Relativity and Gravitation*, 54:30, 2022. doi: 10.1007/s10714-022-02918-3.
- [3] Enrico Specogna, Tatevik Vardanyan, William Giarè, and Eleonora Di Valentino. Slow rolling down the curvature: A reassessment of the planck constraints on ϕ^2 inflation in a closed universe. *Physical Review D*, 113:123526, 2026. doi: 10.1103/dy2d-z3cc.
- [4] Cedric Deffayet, Oriol Pujolas, Ignacy Sawicki, and Alexander Vikman. Imperfect dark energy from kinetic gravity braiding. *JCAP*, 10:026, 2010. doi: 10.1088/1475-7516/2010/10/026.
- [5] Emilio Bellini and Ignacy Sawicki. Maximal freedom at minimum cost: linear large-scale structure in general modifications of gravity. *JCAP*, 07:050, 2014.
- [6] Shingo Akama and Tsutomu Kobayashi. General theory of cosmological perturbations in open and closed universes from the horndeski action. *Physical Review D*, 99:043522, 2019. doi: 10.1103/PhysRevD.99.043522.
- [7] Serena Gambino and Francesco Pace. Linear perturbations in horndeski theories with spatial curvature, 2024. Proceedings contribution, MG17.
- [8] Ignacy Sawicki and Emilio Bellini. Limits of quasistatic approximation in modified-gravity cosmologies. *Phys. Rev. D*, 92:084061, 2015.
- [9] D. Scolnic et al. The pantheon+ analysis: The full dataset and light-curve release. *Astrophys. J.*, 938:113, 2022.
- [10] D. Brout et al. The pantheon+ analysis: Cosmological constraints. *Astrophys. J.*, 938:110, 2022.

- [11] DESI Collaboration. Desi dr2 results ii: Measurements of baryon acoustic oscillations and cosmological constraints. *Physical Review D*, 112:083515, 2025.