

# Monte Carlo simulations of (quasi) constrained ensembles

Victor Martin-Mayor

**Universidad Complutense de Madrid**

in collaboration with

L.A. Fernández, A. Gordillo, J.J. Ruiz-Lorenzo, B. Seoane, P. Verrochio, and D. Yllanes,

Phys. Rev. Lett. **98**, 137207, (2007)

Phys. Rev. Lett. **100**, 057201, (2008)

Nucl. Phys. B **807**, 424-454, (2009)

Phys. Rev. E **80**, 015701(R), (2009)

Phys. Rev. E, *in press*. arXiv:0910.4924

Monte Carlo Algorithms, Melbourne, July 2010

- 1 Introduction
- 2 Quasi constrained statistical ensembles
- 3 Local simulation algorithm
- 4 Cluster methods
- 5 First order phase transitions
- 6 Metastability in disordered systems
- 7 Conclusions

# Outline

- 1 Introduction
- 2 Quasi constrained statistical ensembles
- 3 Local simulation algorithm
- 4 Cluster methods
- 5 First order phase transitions
- 6 Metastability in disordered systems
- 7 Conclusions

# The problem

- Metastability is **ubiquitous** in Physics. In particular, **first-order** phase transitions and/or **glassy** behaviour.

# The problem

- Metastability is **ubiquitous** in Physics. In particular, **first-order** phase transitions and/or **glassy** behaviour.
- Even more **challenging** than critical behaviour  $\tau \propto L^z$ ,  
( $\tau$ : characteristic time,  $L$ : system size,  $z \sim 2$ .)

# The problem

- Metastability is **ubiquitous** in Physics. In particular, **first-order** phase transitions and/or **glassy** behaviour.
- Even more **challenging** than critical behaviour  $\tau \propto L^z$ , ( $\tau$ : characteristic time,  $L$ : system size,  $z \sim 2$ .)
- For metastable systems (and maybe structural glasses),  
**Exponential Critical Slowing Down**

$$\tau \propto e^{\Sigma L^{D-1}}, \quad (\Sigma : \text{surface tension, } D: \text{space dimension}).$$

Spin glasses in 3D:  $z \approx 6.7 \frac{T_c}{T}$ .

# The problem

- Metastability is **ubiquitous** in Physics. In particular, **first-order** phase transitions and/or **glassy** behaviour.
- Even more **challenging** than critical behaviour  $\tau \propto L^z$ , ( $\tau$ : characteristic time,  $L$ : system size,  $z \sim 2$ .)
- For metastable systems (and maybe structural glasses),  
**Exponential Critical Slowing Down**

$$\tau \propto e^{\Sigma L^{D-1}}, \quad (\Sigma : \text{surface tension, } D: \text{space dimension}).$$


Spin glasses in 3D:  $z \approx 6.7 \frac{T_c}{T}$ .

- **Random-walks** on reaction coordinate **alleviate** but do not **cure**.  
Reaction coordinate: temperature (Simulated Annealing, Parallel Tempering), energy (Wang-Landau, multicanonical), spin overlap (multioverlap), etc.

# A different approach

## Metastability

**Discontinuity** in intensive quantity (*reaction coordinate*) as a function of temperature, pressure, ... in large  $N$  limit  
( $N = L^D$ : number of degrees of freedom)






# A different approach

## Metastability

Discontinuity  $\rightarrow$  two-peaked pdf. Minimum  $\sim \exp[-2\Sigma L^{D-1}]$ .





# A different approach

## Metastability

**Idea:** go microcanonical, and reconstruct the **gentle function**  $\langle T \rangle_e$  from **Fluctuation-Dissipation Theorem**. Without random-walk.

Microcanonical MC: Lustig 98', FDT: Martin-Mayor 07' (see also de Pablo 03')

Advantages of micro *analysis* (even with canonical data!): Janke 98'.




# A different approach

## Metastability

**Idea:** go microcanonical, and reconstruct the **gentle function**  $\langle T \rangle_e$  from **Fluctuation-Dissipation Theorem**. Without random-walk.

Microcanonical MC: Lustig 98', FDT: Martin-Mayor 07' (see also de Pablo 03')

Advantages of micro *analysis* (even with canonical data!): Janke 98'.



Sometimes not enough: feed extra information **Tethered Monte Carlo**.

# Outline

- 1 Introduction
- 2 Quasi constrained statistical ensembles**
- 3 Local simulation algorithm
- 4 Cluster methods
- 5 First order phase transitions
- 6 Metastability in disordered systems
- 7 Conclusions

# This talk' toy models

## Ising and Potts models

- Standard benchmark for MC simulation methods.
- Square or cubic lattice, periodic boundary conditions.
- Partition function and main observables ( $N = L^D$ ):

$$Z = \sum_{\{\sigma_{\mathbf{x}}\}} \exp \left[ \beta \sum_{\langle \mathbf{x}, \mathbf{y} \rangle} \sigma_{\mathbf{x}} \sigma_{\mathbf{y}} + h \sum_{\mathbf{x}} \sigma_{\mathbf{x}} \right], \quad \sigma_{\mathbf{x}} = \pm 1,$$

$$U = Nu = - \sum_{\langle \mathbf{x}, \mathbf{y} \rangle} \sigma_{\mathbf{x}} \sigma_{\mathbf{y}}, \quad M = Nm = \sum_{\mathbf{x}} \sigma_{\mathbf{x}}.$$

- We denote canonical averages by  $\langle \dots \rangle_{\beta}$ :

$$C = N[\langle u^2 \rangle_{\beta} - \langle u \rangle_{\beta}^2], \quad \chi = N[\langle m^2 \rangle_{\beta} - \langle m \rangle_{\beta}^2].$$

# This talk' toy models

## Ising and Potts models

- Standard benchmark for MC simulation methods.
- Square or cubic lattice, periodic boundary conditions.
- Partition function and main observables ( $N = L^D$ ):

$$Z = \sum_{\{\sigma_{\mathbf{x}}\}} \exp \left[ \beta \sum_{\langle \mathbf{x}, \mathbf{y} \rangle} \sigma_{\mathbf{x}} \sigma_{\mathbf{y}} + h \sum_{\mathbf{x}} \sigma_{\mathbf{x}} \right], \quad \sigma_{\mathbf{x}} = \pm 1,$$

$$U = Nu = - \sum_{\langle \mathbf{x}, \mathbf{y} \rangle} \sigma_{\mathbf{x}} \sigma_{\mathbf{y}}, \quad M = Nm = \sum_{\mathbf{x}} \sigma_{\mathbf{x}}.$$

- We denote canonical averages by  $\langle \dots \rangle_{\beta}$ :

$$C = N[\langle u^2 \rangle_{\beta} - \langle u \rangle_{\beta}^2], \quad \chi = N[\langle m^2 \rangle_{\beta} - \langle m \rangle_{\beta}^2].$$

- Variation:  $Q$ -states Potts model,  $\sigma_{\mathbf{x}} = 1, 2, \dots, Q$ .

## The problem

- Standard thermodynamic hand-waving:

$$Z(\beta) = \int du \, e^{N[s(u) - \beta u]} \approx e^{N[s(u^*) - \beta u^*]} \quad , \quad \beta = \left. \frac{ds}{du} \right|_{u=u^*}$$

$$Z(\beta, h) = \int dm \, e^{N[\Omega_\beta(m) - hm]} \approx e^{N[\Omega_\beta(m^*) - hm^*]} \quad , \quad h = \left. \frac{\partial \Omega_\beta}{\partial m} \right|_{m=m^*}$$

## The problem

- Standard thermodynamic hand-waving:

$$Z(\beta) = \int du e^{N[s(u) - \beta u]} \approx e^{N[s(u^*) - \beta u^*]}, \quad \beta = \left. \frac{ds}{du} \right|_{u=u^*}$$

$$Z(\beta, h) = \int dm e^{N[\Omega_\beta(m) - hm]} \approx e^{N[\Omega_\beta(m^*) - hm^*]}, \quad h = \left. \frac{\partial \Omega_\beta}{\partial m} \right|_{m=m^*}$$

- Yet,  $m$  and  $u$  discretized variables for finite  $N$ :

$$\Delta m, \Delta u \sim \frac{1}{N}.$$

$s(u), \Omega_\beta(m)$  : comb-like sums of Dirac's delta functions.

**Smooth** effective potentials require coarse-graining.



## The construction of the entropy density $s(e)$

- Artificially add conjugate momenta  $\{\sigma_{\mathbf{x}}\} \longrightarrow \{\sigma_{\mathbf{x}}, \pi_{\mathbf{x}}\}$   
 $\pi_{\mathbf{x}}$  : decoupled Gaussian bath that multiply  $Z(\beta)$  by trivial factor.

## The construction of the entropy density $s(e)$

- Artificially add conjugate momenta  $\{\sigma_{\mathbf{x}}\} \longrightarrow \{\sigma_{\mathbf{x}}, \pi_{\mathbf{x}}\}$   
 $\pi_{\mathbf{x}}$  : decoupled Gaussian bath that multiply  $Z(\beta)$  by trivial factor.
- Original Hamiltonian  $\longrightarrow$  **potential** energy  $U = uN$ .  
 $K = \frac{1}{2} \sum_{\mathbf{x}} \pi_{\mathbf{x}}^2 = N\kappa$ . Total Energy:  $\mathcal{H} = U + K = Ne$ .

## The construction of the entropy density $s(e)$

- Artificially add conjugate momenta  $\{\sigma_{\mathbf{x}}\} \longrightarrow \{\sigma_{\mathbf{x}}, \pi_{\mathbf{x}}\}$   
 $\pi_{\mathbf{x}}$  : decoupled Gaussian bath that multiply  $Z(\beta)$  by trivial factor.
- Original Hamiltonian  $\longrightarrow$  **potential** energy  $U = \underline{u}N$ .  
 $K = \frac{1}{2} \sum_{\mathbf{x}} \pi_{\mathbf{x}}^2 = N\kappa$ . Total Energy:  $\mathcal{H} = U + K = N\mathbf{e}$ .

$$\tilde{Z}(\beta) = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} d\pi_{\mathbf{x}} e^{-\beta \mathcal{H}} = \frac{(2\pi)^{N/2}}{\beta^{N/2}} Z(\beta),$$

.

## The construction of the entropy density $s(\mathbf{e})$

- Artificially add conjugate momenta  $\{\sigma_{\mathbf{x}}\} \longrightarrow \{\sigma_{\mathbf{x}}, \pi_{\mathbf{x}}\}$   
 $\pi_{\mathbf{x}}$  : decoupled Gaussian bath that multiply  $Z(\beta)$  by trivial factor.
- Original Hamiltonian  $\longrightarrow$  **potential** energy  $U = \mathbf{u}N$ .  
 $K = \frac{1}{2} \sum_{\mathbf{x}} \pi_{\mathbf{x}}^2 = N\kappa$ . Total Energy:  $\mathcal{H} = U + K = N\mathbf{e}$ .

$$\tilde{Z}(\beta) = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} d\pi_{\mathbf{x}} e^{-\beta \mathcal{H}} = \frac{(2\pi)^{N/2}}{\beta^{N/2}} Z(\beta),$$

$$\exp[Ns(\mathbf{e}, N)] = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} d\pi_{\mathbf{x}} \delta(\mathcal{H} - \mathbf{e}N),$$

.

## The construction of the entropy density $s(e)$

- Artificially add conjugate momenta  $\{\sigma_{\mathbf{x}}\} \longrightarrow \{\sigma_{\mathbf{x}}, \pi_{\mathbf{x}}\}$   
 $\pi_{\mathbf{x}}$  : decoupled Gaussian bath that multiply  $Z(\beta)$  by trivial factor.
- Original Hamiltonian  $\longrightarrow$  **potential** energy  $U = \mathbf{u}N$ .  
 $K = \frac{1}{2} \sum_{\mathbf{x}} \pi_{\mathbf{x}}^2 = N\kappa$ . Total Energy:  $\mathcal{H} = U + K = N\mathbf{e}$ .

$$\tilde{Z}(\beta) = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} d\pi_{\mathbf{x}} e^{-\beta \mathcal{H}} = \frac{(2\pi)^{N/2}}{\beta^{N/2}} Z(\beta),$$

$$\exp[Ns(\mathbf{e}, N)] = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} d\pi_{\mathbf{x}} \delta(\mathcal{H} - \mathbf{e}N),$$

$$\tilde{Z}(\beta) = \int d\mathbf{e} e^{N(s(\mathbf{e}, N) - \beta \mathbf{e})}.$$

## The construction of the entropy density $s(e)$

- Artificially add conjugate momenta  $\{\sigma_{\mathbf{x}}\} \longrightarrow \{\sigma_{\mathbf{x}}, \pi_{\mathbf{x}}\}$   
 $\pi_{\mathbf{x}}$  : decoupled Gaussian bath that multiply  $Z(\beta)$  by trivial factor.
- Original Hamiltonian  $\longrightarrow$  **potential** energy  $U = \mathbf{u}N$ .  
 $K = \frac{1}{2} \sum_{\mathbf{x}} \pi_{\mathbf{x}}^2 = N\kappa$ . Total Energy:  $\mathcal{H} = U + K = N\mathbf{e}$ .
- Canonical pdf:

$$p_{\beta}(\mathbf{e}) = \frac{e^{N(s(\mathbf{e}, N) - \beta \mathbf{e})}}{\tilde{Z}(\beta)}.$$

## The construction of the entropy density $s(e)$

- Artificially add conjugate momenta  $\{\sigma_{\mathbf{x}}\} \rightarrow \{\sigma_{\mathbf{x}}, \pi_{\mathbf{x}}\}$   
 $\pi_{\mathbf{x}}$  : decoupled Gaussian bath that multiply  $Z(\beta)$  by trivial factor.
- Original Hamiltonian  $\rightarrow$  **potential** energy  $U = uN$ .  
 $K = \frac{1}{2} \sum_{\mathbf{x}} \pi_{\mathbf{x}}^2 = N\kappa$ . Total Energy:  $\mathcal{H} = U + K = Ne$ .
- Canonical pdf:

$$p_{\beta}(e) = \frac{e^{N(s(e,N) - \beta e)}}{\tilde{Z}(\beta)}.$$

- $p_{\beta}(e)$  is a **convolution** ( $\kappa$  and  $u$  independent!):

$$p_{\beta}(e) = \int \int d\kappa du p_{\beta,u}(u) p_{\beta,\kappa}(\kappa) \delta(e - u - \kappa).$$

$$\kappa = \frac{1}{2\beta} \left[ 1 + \zeta \sqrt{\frac{2}{N}} \right], \zeta \sim 1 \text{ and (almost) Gaussian distributed.}$$

$p_{\beta,u}(u)$  gets shifted by  $\frac{1}{2\beta}$  and **smoothed**.

Tethered effective potential  $\Omega_N(\hat{m}, \beta)$ , analogue to  $s(e, N)$

- Conjugate momenta now leave unchanged  $Z(\beta)$

$$Z(\beta) = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} \frac{d\pi_{\mathbf{x}}}{\sqrt{2\pi}} e^{-\beta U - \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2}}.$$



## Tethered effective potential $\Omega_N(\hat{m}, \beta)$ , analogue to $s(e, N)$

- Conjugate momenta now leave unchanged  $Z(\beta)$

$$Z(\beta) = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} \frac{d\pi_{\mathbf{x}}}{\sqrt{2\pi}} e^{-\beta U - \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2}}.$$

- We couple  $\pi_{\mathbf{x}}$  with order parameter:

$$p_{\beta}(\hat{m}) = \frac{1}{Z} \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} \frac{d\pi_{\mathbf{x}}}{\sqrt{2\pi}} e^{-\beta U} \delta(N\hat{m} - Nm - \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2}) = e^{N\Omega_N(\hat{m}, \beta)}$$

## Tethered effective potential $\Omega_N(\hat{m}, \beta)$ , analogue to $s(e, N)$

- Conjugate momenta now leave unchanged  $Z(\beta)$

$$Z(\beta) = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} \frac{d\pi_{\mathbf{x}}}{\sqrt{2\pi}} e^{-\beta U - \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2}}.$$

- We couple  $\pi_{\mathbf{x}}$  with order parameter:

$$p_{\beta}(\hat{m}) = \frac{1}{Z} \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} \frac{d\pi_{\mathbf{x}}}{\sqrt{2\pi}} e^{-\beta U} \delta(N\hat{m} - Nm - \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2}) = e^{N\Omega_N(\hat{m}, \beta)}$$

- Again,  $r = \frac{1}{N} \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2} = \frac{1}{2} + \frac{\zeta}{\sqrt{N}}$ , with  $\zeta \sim 1$ .

## Tethered effective potential $\Omega_N(\hat{m}, \beta)$ , analogue to $s(e, N)$

- Conjugate momenta now leave unchanged  $Z(\beta)$

$$Z(\beta) = \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} \frac{d\pi_{\mathbf{x}}}{\sqrt{2\pi}} e^{-\beta U - \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2}}.$$

- We couple  $\pi_{\mathbf{x}}$  with order parameter:

$$p_{\beta}(\hat{m}) = \frac{1}{Z} \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} \frac{d\pi_{\mathbf{x}}}{\sqrt{2\pi}} e^{-\beta U} \delta(N\hat{m} - Nm - \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2}) = e^{N\Omega_N(\hat{m}, \beta)}$$

- Again,  $r = \frac{1}{N} \sum_{\mathbf{x}} \frac{\pi_{\mathbf{x}}^2}{2} = \frac{1}{2} + \frac{\zeta}{\sqrt{N}}$ , with  $\zeta \sim 1$ .
- $\hat{m} \approx m + \frac{1}{2}$ :  $p_{\beta}(\hat{m})$  is a **convolution** of  $p_{\beta, m}$  and  $p_{\beta, r}$ .

# The microcanonical ensemble

( $\beta$  and  $e$  reverse their roles)

- Integrate-out  $\{\pi_{\mathbf{x}}\}$ : **microcanonical** mean value at fixed  $e$ ! (Lustig 98')

$$\begin{aligned}\exp[N s(\mathbf{e}, N)] &= \sum_{\{\sigma_{\mathbf{x}}\}} \int \prod_{\mathbf{x}} d\pi_{\mathbf{x}} \delta\left(\frac{1}{2} \sum_{\{\sigma_{\mathbf{x}}\}} \pi_{\mathbf{x}}^2 + uN - eN\right), \\ &= \text{constant} \times \sum_{\{\sigma_{\mathbf{x}}\}} (e - u)^{\frac{N-2}{2}} \theta(e - u).\end{aligned}$$

$$\langle O \rangle_e = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(\mathbf{e}; \{\sigma\}) \omega^{\text{mic}}(\{\sigma\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{mic}}(\{\sigma\})}, \quad \omega^{\text{mic}} = (e - u)^{\frac{N-2}{2}} \theta(e - u).$$

# The microcanonical ensemble

( $\beta$  and  $e$  reverse their roles)

- Integrate-out  $\{\pi_{\mathbf{x}}\}$ : **microcanonical** mean value at fixed  $e$ ! (Lustig 98')

$$\langle O \rangle_e = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(e; \{\sigma\}) \omega^{\text{mic}}(\{\sigma\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{mic}}(\{\sigma\})}, \quad \omega^{\text{mic}} = (e - u)^{\frac{N-2}{2}} \theta(e - u).$$

- $\hat{\beta} = \frac{N-2}{2N} \frac{1}{e-u}, \quad \frac{ds(e, N)}{de} = \langle \hat{\beta} \rangle_e.$

# The microcanonical ensemble

( $\beta$  and  $e$  reverse their roles)

- Integrate-out  $\{\pi_{\mathbf{x}}\}$ : **microcanonical** mean value at fixed  $e$ ! (Lustig 98')

$$\langle O \rangle_e = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(\mathbf{e}; \{\sigma\}) \omega^{\text{mic}}(\{\sigma\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{mic}}(\{\sigma\})}, \quad \omega^{\text{mic}} = (e - u)^{\frac{N-2}{2}} \theta(e - u).$$

- $\hat{\beta} = \frac{N-2}{2N} \frac{1}{e-u}$  ,  $\frac{ds(\mathbf{e}, N)}{d\mathbf{e}} = \langle \hat{\beta} \rangle_e$ .
- Canonical** pdf:  $\log P_{\beta}(e_2) - \log P_{\beta}(e_1) = N \int_{e_1}^{e_2} d\mathbf{e} \left( \langle \hat{\beta} \rangle_e - \beta \right)$ .

# The microcanonical ensemble

( $\beta$  and  $e$  reverse their roles)

- Integrate-out  $\{\pi_{\mathbf{x}}\}$ : **microcanonical** mean value at fixed  $e$ ! (Lustig 98')

$$\langle O \rangle_e = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(e; \{\sigma\}) \omega^{\text{mic}}(\{\sigma\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{mic}}(\{\sigma\})}, \quad \omega^{\text{mic}} = (e - u)^{\frac{N-2}{2}} \theta(e - u).$$

- $\hat{\beta} = \frac{N-2}{2N} \frac{1}{e-u}$  ,  $\frac{ds(e, N)}{de} = \langle \hat{\beta} \rangle_e$ .
- Canonical** pdf:  $\log P_{\beta}(e_2) - \log P_{\beta}(e_1) = N \int_{e_1}^{e_2} de \left( \langle \hat{\beta} \rangle_e - \beta \right)$ .
- Maximum** (actually extrema) of canonical pdf:  $\langle \hat{\beta} \rangle_e = \beta$ .

# The microcanonical ensemble

( $\beta$  and  $e$  reverse their roles)

- Integrate-out  $\{\pi_{\mathbf{x}}\}$ : **microcanonical** mean value at fixed  $e$ ! (Lustig 98')

$$\langle O \rangle_e = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(e; \{\sigma\}) \omega^{\text{mic}}(\{\sigma\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{mic}}(\{\sigma\})}, \quad \omega^{\text{mic}} = (e - u)^{\frac{N-2}{2}} \theta(e - u).$$

- $\hat{\beta} = \frac{N-2}{2N} \frac{1}{e-u}$  ,  $\frac{ds(e, N)}{de} = \langle \hat{\beta} \rangle_e$ .
- **Canonical** pdf:  $\log P_{\beta}(e_2) - \log P_{\beta}(e_1) = N \int_{e_1}^{e_2} de \left( \langle \hat{\beta} \rangle_e - \beta \right)$ .
- **Maximum** (actually extrema) of canonical pdf:  $\langle \hat{\beta} \rangle_e = \beta$ .
- Similar to **Creutz's microcanonical daemon** but



# The microcanonical ensemble

( $\beta$  and  $e$  reverse their roles)

- Integrate-out  $\{\pi_{\mathbf{x}}\}$ : **microcanonical** mean value at fixed  $e$ ! (Lustig 98')

$$\langle O \rangle_e = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(e; \{\sigma\}) \omega^{\text{mic}}(\{\sigma\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{mic}}(\{\sigma\})}, \quad \omega^{\text{mic}} = (e - u)^{\frac{N-2}{2}} \theta(e - u).$$

- $\hat{\beta} = \frac{N-2}{2N} \frac{1}{e-u}$ ,  $\frac{ds(e, N)}{de} = \langle \hat{\beta} \rangle_e$ .
- Canonical** pdf:  $\log P_{\beta}(e_2) - \log P_{\beta}(e_1) = N \int_{e_1}^{e_2} de \left( \langle \hat{\beta} \rangle_e - \beta \right)$ .
- Maximum** (actually extrema) of canonical pdf:  $\langle \hat{\beta} \rangle_e = \beta$ .
- Similar to **Creutz's microcanonical daemon** but
  - We have an **extensive** number of daemons.
  - Daemons are **continuous** variables.
  - We integrate them out.

# The tethered ensemble

- Integrating demons out in the *constrained* (fixed  $\hat{m}$ ) partition function  $\rightarrow$  **tethered expectation values**:

$$\langle O \rangle_{\hat{m}, \beta} = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(\hat{m}; \{\sigma_{\mathbf{x}}\}) \omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\})},$$

# The tethered ensemble

- Integrating demons out in the *constrained* (fixed  $\hat{m}$ ) partition function  $\rightarrow$  **tethered expectation values**:

$$\langle O \rangle_{\hat{m}, \beta} = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(\hat{m}; \{\sigma_{\mathbf{x}}\}) \omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\})},$$
$$\omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\}) = e^{-\beta U + M - N \hat{m}} (\hat{m} - m)^{(N-2)/2} \theta(\hat{m} - m).$$

# The tethered ensemble

- Integrating demons out in the *constrained* (fixed  $\hat{m}$ ) partition function  $\rightarrow$  **tethered expectation values**:

$$\langle O \rangle_{\hat{m}, \beta} = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(\hat{m}; \{\sigma_{\mathbf{x}}\}) \omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\})},$$

$$\omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\}) = e^{-\beta U + M - N \hat{m}} (\hat{m} - m)^{(N-2)/2} \theta(\hat{m} - m).$$

- The *canonical*  $\Omega_N$  follows from **Fluctuation-Dissipation**

$$\hat{h}(\hat{m}; \{\sigma_{\mathbf{x}}\}) = -1 + \frac{N/2 - 1}{\hat{M} - M} \quad \Rightarrow \quad \langle \hat{h} \rangle_{\hat{m}, \beta} = \frac{\partial \Omega_N(\hat{m}, \beta)}{\partial \hat{m}}.$$

# The tethered ensemble

- Integrating demons out in the *constrained* (fixed  $\hat{m}$ ) partition function  $\rightarrow$  **tethered expectation values**:

$$\langle O \rangle_{\hat{m},\beta} = \frac{\sum_{\{\sigma_{\mathbf{x}}\}} O(\hat{m}; \{\sigma_{\mathbf{x}}\}) \omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\})}{\sum_{\{\sigma_{\mathbf{x}}\}} \omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\})},$$

$$\omega^{\text{teth}}(\beta, \hat{m}, N; \{\sigma_{\mathbf{x}}\}) = e^{-\beta U + M - N\hat{m}} (\hat{m} - m)^{(N-2)/2} \theta(\hat{m} - m).$$


- The *canonical*  $\Omega_N$  follows from **Fluctuation-Dissipation**

$$\hat{h}(\hat{m}; \{\sigma_{\mathbf{x}}\}) = -1 + \frac{N/2 - 1}{\hat{M} - M} \implies \langle \hat{h} \rangle_{\hat{m},\beta} = \frac{\partial \Omega_N(\hat{m}, \beta)}{\partial \hat{m}}.$$

- Tethered mean values  $\langle O \rangle_{\hat{m},\beta} \leftrightarrow$  canonical mean values  $\langle O \rangle_{\beta}(h)$ , for any external field  $h$ :

$$\langle O \rangle_{\beta}(h) = \int d\hat{m} \langle O \rangle_{\hat{m},\beta} \exp[N(\Omega_N(\hat{m}, \beta) + h\hat{m})].$$

# A prototypical tethered computation




Result:  $\langle u \rangle_{\beta_c} = -0.3309831(15)$

(Ising,  $L = 128$ ,  $D = 3$ )

## Steps

- 1 Select a mesh of  $\hat{m}$  values.

# A prototypical tethered computation




Result:  $\langle u \rangle_{\beta_c} = -0.3309831(15)$

(Ising,  $L = 128$ ,  $D = 3$ )

## Steps

- 1 Select a mesh of  $\hat{m}$  values.
- 2 Independent simulation for each  $\hat{m}$ . Get  $\langle O \rangle_{\hat{m}, \beta}$ .

# A prototypical tethered computation



Result:  $\langle u \rangle_{\beta_c} = -0.3309831(15)$


(Ising,  $L = 128$ ,  $D = 3$ )

## Steps

- 1 Select a mesh of  $\hat{m}$  values.
- 2 Independent simulation for each  $\hat{m}$ . Get  $\langle O \rangle_{\hat{m}, \beta}$ .
- 3  $\langle O \rangle_{\hat{m}, \beta}$  smooth functions of  $\hat{m}$   
 $\rightarrow$  **interpolate** (e.g. spline).



# A prototypical tethered computation




Result:  $\langle u \rangle_{\beta_c} = -0.3309831(15)$

(Ising,  $L = 128$ ,  $D = 3$ )

## Steps

- 1 Select a mesh of  $\hat{m}$  values.
- 2 Independent simulation for each  $\hat{m}$ . Get  $\langle O \rangle_{\hat{m},\beta}$ .
- 3  $\langle O \rangle_{\hat{m},\beta}$  smooth functions of  $\hat{m}$   
→ **interpolate** (e.g. spline).
- 4 Numerical integration of  $\langle \hat{h} \rangle_{\hat{m},\beta}$  yields  $\Omega_N(\hat{m}, \beta)$ .

# A prototypical tethered computation




Result:  $\langle u \rangle_{\beta_c} = -0.3309831(15)$

(Ising,  $L = 128$ ,  $D = 3$ )

## Steps

- 1 Select a mesh of  $\hat{m}$  values.
- 2 Independent simulation for each  $\hat{m}$ . Get  $\langle O \rangle_{\hat{m}, \beta}$ .
- 3  $\langle O \rangle_{\hat{m}, \beta}$  smooth functions of  $\hat{m}$   
→ **interpolate** (e.g. spline).
- 4 Numerical integration of  $\langle \hat{h} \rangle_{\hat{m}, \beta}$  yields  $\Omega_N(\hat{m}, \beta)$ .
- 5 Reconstruct canonical  $\langle O \rangle_{\beta}$  from  $p(\hat{m})$  and  $\langle O \rangle_{\hat{m}, \beta}$ .

# A prototypical tethered computation




Result:  $\langle u \rangle_{\beta_c} = -0.3309831(15)$

(Ising,  $L = 128$ ,  $D = 3$ )

## Steps

- 1 Select a mesh of  $\hat{m}$  values.
- 2 Independent simulation for each  $\hat{m}$ . Get  $\langle O \rangle_{\hat{m},\beta}$ .
- 3  $\langle O \rangle_{\hat{m},\beta}$  smooth functions of  $\hat{m}$   
→ **interpolate** (e.g. spline).
- 4 Numerical integration of  $\langle \hat{h} \rangle_{\hat{m},\beta}$  yields  $\Omega_N(\hat{m}, \beta)$ .
- 5 Reconstruct canonical  $\langle O \rangle_{\beta}$  from  $p(\hat{m})$  and  $\langle O \rangle_{\hat{m},\beta}$ .
- 6 Statistical errors: jackknife.

# A prototypical tethered computation



Result:  $\langle u \rangle_{\beta_c} = -0.3309831(15)$

(Ising,  $L = 128$ ,  $D = 3$ )

## Steps

- 1 Select a mesh of  $\hat{m}$  values.
- 2 Independent simulation for each  $\hat{m}$ . Get  $\langle O \rangle_{\hat{m}, \beta}$ .
- 3  $\langle O \rangle_{\hat{m}, \beta}$  smooth functions of  $\hat{m}$   
→ **interpolate** (e.g. spline).
- 4 Numerical integration of  $\langle \hat{h} \rangle_{\hat{m}, \beta}$  yields  $\Omega_N(\hat{m}, \beta)$ .
- 5 Reconstruct canonical  $\langle O \rangle_{\beta}$  from  $p(\hat{m})$  and  $\langle O \rangle_{\hat{m}, \beta}$ .
- 6 Statistical errors: jackknife.
- 7 Systematic errors: refine  $\hat{m}$  mesh.

# Outline

- 1 Introduction
- 2 Quasi constrained statistical ensembles
- 3 Local simulation algorithm**
- 4 Cluster methods
- 5 First order phase transitions
- 6 Metastability in disordered systems
- 7 Conclusions

# Metropolis algorithm

Start from your working Monte Carlo (canonical) program:

- Try a spin-flip, (or to displace a particle, or whatever local move).

# Metropolis algorithm

Start from your working Monte Carlo (canonical) program:

- Try a spin-flip, (or to displace a particle, or whatever local move).
- Accept the move with Metropolis probability:

$$p_{\text{accept}} = \min \left[ 1, \frac{\omega_{\text{new}}}{\omega_{\text{old}}} \right], \quad \omega = \omega^{\text{mic}}, \omega^{\text{teth}}.$$

# Metropolis algorithm

Start from your working Monte Carlo (canonical) program:


- Try a spin-flip, (or to displace a particle, or whatever local move).
- Accept the move with Metropolis probability:

$$p_{\text{accept}} = \min \left[ 1, \frac{\omega_{\text{new}}}{\omega_{\text{old}}} \right], \quad \omega = \omega^{\text{mic}}, \omega^{\text{teth}}.$$

- Discrete spin systems:  $p_{\text{accept}}$  takes a finite number of values  $\Rightarrow$  Look-up-table provide significant **speed-up**.



# Autocorrelation times for Metropolis



- $\tau_{\text{int}}$ : dramatic dependence on **observable**, and on  $\hat{m}$ .
- Functions of  $m$  (e.g.  $\hat{h}$ ): **no measurable critical slowing down**.
- Energy or propagator's Fourier transform ( $\vec{k} \neq 0$ )  
 $\tau_{\text{int}}(\hat{m} = 0.5) \approx L^2$   
Worst case:  $m \sim 0$  or  $\hat{m} = \frac{1}{2}$ .

# Outline

- 1 Introduction
- 2 Quasi constrained statistical ensembles
- 3 Local simulation algorithm
- 4 Cluster methods**
- 5 First order phase transitions
- 6 Metastability in disordered systems
- 7 Conclusions

# Cluster method with constraint

- Nothing really new: Fortuyn-Kasteleyn.

# Cluster method with constraint

- Nothing really new: Fortuyn-Kasteleyn.
- Microcanonic case straightforward:

# Cluster method with constraint

- Nothing really new: Fortuyn-Kasteleyn.
- Microcanonic case straightforward:
  - $\omega^{\text{mic}} = [\omega^{\text{mic}} e^{kU}] e^{-kU}$ ,  $k$ : tunable parameter.

# Cluster method with constraint

- Nothing really new: Fortuyn-Kasteleyn.
- Microcanonic case straightforward:
  - $\omega^{\text{mic}} = [\omega^{\text{mic}} e^{kU}] e^{-kU}$ ,  $k$ : tunable parameter.
  - Apply Fortuyn-Kasteleyn to  $e^{-kU} \implies$  trace clusters as canonical.

# Cluster method with constraint

- Nothing really new: Fortuyn-Kasteleyn.
- Microcanonic case straightforward:
  - $\omega^{\text{mic}} = [\omega^{\text{mic}} e^{kU}] e^{-kU}$ ,  $k$ : tunable parameter.
  - Apply Fortuyn-Kasteleyn to  $e^{-kU} \implies$  trace clusters as canonical.
  - Choose spin-values for clusters at random.

# Cluster method with constraint

- Nothing really new: Fortuyn-Kasteleyn.
- Microcanonic case straightforward:
  - $\omega^{\text{mic}} = [\omega^{\text{mic}} e^{kU}] e^{-kU}$ ,  $k$ : tunable parameter.
  - Apply Fortuyn-Kasteleyn to  $e^{-kU} \Rightarrow$  trace clusters as canonical.
  - Choose spin-values for clusters at random.
  - Accept spin assignment with Metropolis probability.

$$p_{\text{accept}}^{\text{cluster}} = \min \left[ 1, \frac{\omega_{\text{new}}^{\text{mic}} e^{kU_{\text{new}}}}{\omega_{\text{old}}^{\text{mic}} e^{kU_{\text{old}}}} \right].$$



# Cluster method with constraint

- Nothing really new: Fortuyn-Kasteleyn.
- Microcanonic case straightforward:
  - $\omega^{\text{mic}} = [\omega^{\text{mic}} e^{kU}] e^{-kU}$ ,  $k$ : tunable parameter.
  - Apply Fortuyn-Kasteleyn to  $e^{-kU} \Rightarrow$  trace clusters as canonical.
  - Choose spin-values for clusters at random.
  - Accept spin assignment with Metropolis probability.

$$p_{\text{accept}}^{\text{cluster}} = \min \left[ 1, \frac{\omega_{\text{new}}^{\text{mic}} e^{kU_{\text{new}}}}{\omega_{\text{old}}^{\text{mic}} e^{kU_{\text{old}}}} \right].$$

- $k = \langle \hat{\beta} \rangle_e \rightarrow$  Potts model:  $p_{\text{accept}}^{\text{cluster}} \approx 60\%$ .

# The tethered cluster update algorithm (I)

- We can follow the Fortuin-Kasteleyn construction.
- Edwards-Sokal: introduce bond-occupation variables  $n_{xy}$  ( $= 0, 1$ )

$$\exp[\beta(\sigma_x\sigma_y-1)] = \sum_{n_{xy}=0,1} [(1-p)\delta_{n_{xy},0} + p\delta_{\sigma_x,\sigma_y}\delta_{n_{xy},1}], \quad p = 1 - e^{-2\beta}.$$

these bonds form  $N_C$  clusters.

# The tethered cluster update algorithm (I)

- We can follow the Fortuin-Kasteleyn construction.
- Edwards-Sokal: introduce bond-occupation variables  $n_{xy}$  ( $= 0, 1$ )

$$\exp[\beta(\sigma_x \sigma_y - 1)] = \sum_{n_{xy}=0,1} [(1-p)\delta_{n_{xy},0} + p\delta_{\sigma_x, \sigma_y} \delta_{n_{xy},1}], \quad p = 1 - e^{-2\beta}.$$

these bonds form  $N_C$  clusters.

- Combine this identity with  $\omega(\beta, \hat{m}; \{\sigma_x\})$ :

$$\omega(\beta, \hat{m}; \{\sigma_x\}) = e^{-\beta U + M - N\hat{m}} (\hat{m} - m)^{(N-2)/2} \theta(\hat{m} - m).$$

# The tethered cluster update algorithm (I)

- We can follow the Fortuin-Kasteleyn construction.
- Edwards-Sokal: introduce bond-occupation variables  $n_{xy}$  ( $= 0, 1$ )

$$\exp[\beta(\sigma_x\sigma_y-1)] = \sum_{n_{xy}=0,1} [(1-p)\delta_{n_{xy},0} + p\delta_{\sigma_x,\sigma_y}\delta_{n_{xy},1}], \quad p = 1 - e^{-2\beta}.$$

these bonds form  $N_C$  clusters.

- Combine this identity with  $\omega(\beta, \hat{m}; \{\sigma_x\})$ :

$$\omega(\beta, \hat{m}; \{\sigma_x\}) = e^{-\beta U + M - N\hat{m}} (\hat{m} - m)^{(N-2)/2} \theta(\hat{m} - m).$$

- We have the following conditional probabilities:
  - Given  $\{\sigma_x\}$ , just as in the canonical case, bonds are independent and  $n_{xy}$  is 1 with probability  $p\delta_{\sigma_x,\sigma_y}$ .

# The tethered cluster update algorithm (I)

- We can follow the Fortuin-Kasteleyn construction.
- Edwards-Sokal: introduce bond-occupation variables  $n_{xy}$  ( $= 0, 1$ )

$$\exp[\beta(\sigma_x \sigma_y - 1)] = \sum_{n_{xy}=0,1} [(1-p)\delta_{n_{xy},0} + p\delta_{\sigma_x, \sigma_y} \delta_{n_{xy},1}], \quad p = 1 - e^{-2\beta}.$$

these bonds form  $N_C$  clusters.

- Combine this identity with  $\omega(\beta, \hat{m}; \{\sigma_x\})$ :

$$\omega(\beta, \hat{m}; \{\sigma_x\}) = e^{-\beta U + M - N\hat{m}} (\hat{m} - m)^{(N-2)/2} \theta(\hat{m} - m).$$

- We have the following conditional probabilities:
  - Given  $\{\sigma_x\}$ , just as in the canonical case, bonds are independent and  $n_{xy}$  is 1 with probability  $p\delta_{\sigma_x, \sigma_y}$ .
  - Given  $\{n_{xy}\}$ , the spins within cluster  $i$  are equal to  $S_i = \pm 1$ .  
Not all  $\{S_i\}$  configurations have the same probability:

$$p(\{S_i\}) \propto e^{M - \hat{M}} (\hat{M} - M)^{(N-2)/2} \theta(\hat{M} - M).$$

# The tethered cluster update algorithm (II)

- a) Cluster tracing
- b) Cluster flipping:

.

# The tethered cluster update algorithm (II)

- a) Cluster tracing (as in canonical SW)
- b) Cluster flipping:

.

# The tethered cluster update algorithm (II)

- a) Cluster tracing (as in canonical SW)
- b) Cluster flipping: It is impossible to consider all  $2^{N_c}$  cluster orientations



# The tethered cluster update algorithm (II)

- a) Cluster tracing (as in canonical SW)
- b) Cluster flipping: We perform a dynamic MC
  - 1 Select  $N'_C \ll N_C$  clusters (for instance  $N'_C \approx 5$ ).

# The tethered cluster update algorithm (II)

- a) Cluster tracing (as in canonical SW)
- b) Cluster flipping: We perform a dynamic MC
  - 1 Select  $N'_C \ll N_C$  clusters (for instance  $N'_C \approx 5$ ).
  - 2 Heat bath among the  $2^{N'_C}$  configurations with  $N_C - N'_C$  clusters fixed.

# The tethered cluster update algorithm (II)

- a) Cluster tracing (as in canonical SW)
- b) Cluster flipping: We perform a dynamic MC
  - ① Select  $N'_C \ll N_C$  clusters (for instance  $N'_C \approx 5$ ).
  - ② Heat bath among the  $2^{N'_C}$  configurations with  $N_C - N'_C$  clusters fixed.
  - ③ Take measurements.

Steps 1–3 are repeated  $N_{\text{REP}}$  times, without retracing the clusters.

# The tethered cluster update algorithm (II)


- a) Cluster tracing (as in canonical SW)
- b) Cluster flipping: We perform a dynamic MC
  - ① Select  $N'_C \ll N_C$  clusters (for instance  $N'_C \approx 5$ ).
  - ② Heat bath among the  $2^{N'_C}$  configurations with  $N_C - N'_C$  clusters fixed.
  - ③ Take measurements.

Steps 1–3 are repeated  $N_{\text{REP}}$  times, without retracing the clusters.

- We consider the overlap

$$o = \frac{\langle \sum_{\mathbf{x}} [\sigma_{\mathbf{x}}^{t=0} \sigma_{\mathbf{x}}^{t=N_{\text{rep}}} - \langle m \rangle_{\hat{m},\beta}^2] \rangle_{\hat{m},\beta}}{N(1 - \langle m \rangle_{\hat{m},\beta}^2)},$$

- Our  $N_{\text{REP}}$ : Tracing time  $\approx$  Flipping time



# The tethered cluster update algorithm (II)


- a) Cluster tracing (as in canonical SW)
- b) Cluster flipping: We perform a dynamic MC
  - ① Select  $N'_C \ll N_C$  clusters (for instance  $N'_C \approx 5$ ).
  - ② Heat bath among the  $2^{N'_C}$  configurations with  $N_C - N'_C$  clusters fixed.
  - ③ Take measurements.

Steps 1–3 are repeated  $N_{\text{REP}}$  times, without retracing the clusters.

- We consider the overlap

$$o = \frac{\langle \sum_{\mathbf{x}} [\sigma_{\mathbf{x}}^{t=0} \sigma_{\mathbf{x}}^{t=N_{\text{rep}}} - \langle m \rangle_{\hat{m}, \beta}^2] \rangle_{\hat{m}, \beta}}{N(1 - \langle m \rangle_{\hat{m}, \beta}^2)},$$

- Our  $N_{\text{REP}}$ : Tracing time  $\approx$  Flipping time
- Measuring  $\hat{h}$  at each of the  $N_{\text{REP}}$  steps reduces errors by up to a factor 25



# Results at the critical point

## Integrated autocorrelation times

- The slowest observable is  $E$ .
- $\tau_{\text{int}}$  largest at  $\hat{m} = 0.5$  ( $m \simeq 0$ ).

# Results at the critical point


## Integrated autocorrelation times

- The slowest observable is  $E$ .
- $\tau_{\text{int}}$  largest at  $\hat{m} = 0.5$  ( $m \simeq 0$ ).
- We fit to  $\tau_{\text{int}} = AL^z$ :
  - $D = 2 \rightarrow z = 0.241(7)$

# Results at the critical point

## Integrated autocorrelation times

- The slowest observable is  $E$ .
- $\tau_{\text{int}}$  largest at  $\hat{m} = 0.5$  ( $m \simeq 0$ ).
- We fit to  $\tau_{\text{int}} = AL^z$ :
  - $D = 2 \rightarrow z = 0.241(7)$
  - $D = 3 \rightarrow z = 0.472(8)$






# Results at the critical point

## Integrated autocorrelation times


- The slowest observable is  $E$ .
- $\tau_{\text{int}}$  largest at  $\hat{m} = 0.5$  ( $m \simeq 0$ ).
- We fit to  $\tau_{\text{int}} = AL^z$ :
  - $D = 2 \rightarrow z = 0.241(7)$
  - $D = 3 \rightarrow z = 0.472(8)$
- $D = 3$ ,  $z$  compatible with that of canonical Swendsen-Wang (data from Ossola and Sokal, 2001)



# Results at the critical point

## Integrated autocorrelation times

- The slowest observable is  $E$ .
- $\tau_{\text{int}}$  largest at  $\hat{m} = 0.5$  ( $m \simeq 0$ ).
- We fit to  $\tau_{\text{int}} = AL^z$ :
  - $D = 2 \rightarrow z = 0.241(7)$
  - $D = 3 \rightarrow z = 0.472(8)$
- $D = 3$ ,  $z$  compatible with that of canonical Swendsen-Wang (data from Ossola and Sokal, 2001)



## Canonical averages for $L = 128$ , $D = 3$


	MCS	$-\langle e \rangle_\beta$	$\mathcal{C}$	$\chi$	$\xi$
SW	$48 \times 10^6$	0.3309822(16)	22.155(18)	21193(13)	82.20(3)
TMC	$50 \times 10^6$	0.3309831(15)	22.174(13)	21202(13)	82.20(5)

# Outline

- 1 Introduction
- 2 Quasi constrained statistical ensembles
- 3 Local simulation algorithm
- 4 Cluster methods
- 5 First order phase transitions**
- 6 Metastability in disordered systems
- 7 Conclusions

# A standard example: $D = 2$ , $Q = 10$ , Potts model

- Use **cluster** method




# A standard example: $D = 2$ , $Q = 10$ , Potts model

- Use **cluster** method
- At phase coexistence  $\langle \hat{\beta} \rangle_e$  **non decreasing**,

$$\langle \hat{\beta} \rangle_e = \beta_c \quad \text{at}$$

$e_d$ : disordered phase,  
 $e^*$ : saddle-point,  
 $e_o$ : ordered phase.




# A standard example: $D = 2$ , $Q = 10$ , Potts model

- Use **cluster** method
- At phase coexistence  $\langle \hat{\beta} \rangle_e$  **non decreasing**,

$$\langle \hat{\beta} \rangle_e = \beta_c \quad \text{at}$$

$e_d$ : disordered phase,  
 $e^*$ : saddle-point,  
 $e_o$ : ordered phase.

- $\beta_c$  from **Maxwell** rule  
6 significant digits, on  $L = 1024$ .



# A standard example: $D = 2$ , $Q = 10$ , Potts model


- Use **cluster** method
- At phase coexistence  $\langle \hat{\beta} \rangle_e$  **non decreasing**,

$$\langle \hat{\beta} \rangle_e = \beta_c \quad \text{at}$$



$e_d$ : disordered phase,  
 $e^*$ : saddle-point,  
 $e_o$ : ordered phase.

- $\beta_c$  from **Maxwell** rule  
6 significant digits, on  $L = 1024$ .
- **Surface tension**:

$$\Sigma_L = \frac{L}{2} \int_{e^*}^{e_d} de (\langle \hat{\beta} \rangle_e - \beta_c)$$





# Geometric transitions





# Geometric transitions



# Outline

- 1 Introduction
- 2 Quasi constrained statistical ensembles
- 3 Local simulation algorithm
- 4 Cluster methods
- 5 First order phase transitions
- 6 Metastability in disordered systems**
- 7 Conclusions

# Short reminder: disordered systems

- Two types of variables, **dynamical** (i.e. spins) and **quenched** (geologically slow: atomic impurities, vacancies, etc.).

# Short reminder: disordered systems

- Two types of variables, **dynamical** (i.e. spins) and **quenched** (geologically slow: atomic impurities, vacancies, etc.).
- Impurities  $\longrightarrow$  **random** Hamiltonian  $H_J$ . Site-dependent coupling constants (**sample**), extracted from some pdf  $P(J)$ .

# Short reminder: disordered systems

- Two types of variables, **dynamical** (i.e. spins) and **quenched** (geologically slow: atomic impurities, vacancies, etc.).
- Impurities  $\longrightarrow$  **random** Hamiltonian  $H_J$ . Site-dependent coupling constants (**sample**), extracted from some pdf  $P(J)$ .
- **Quenched** approximation: fast variables (spins) allowed **no backreaction** on slow ones. In practice:

# Short reminder: disordered systems

- Two types of variables, **dynamical** (i.e. spins) and **quenched** (geologically slow: atomic impurities, vacancies, etc.).
- Impurities  $\longrightarrow$  **random** Hamiltonian  $H_J$ . Site-dependent coupling constants (**sample**), extracted from some pdf  $P(J)$ .
- **Quenched** approximation: fast variables (spins) allowed **no backreaction** on slow ones. In practice:
  - 1 Take  $J$ -dependent thermal average  $\langle \dots \rangle_J$ .

# Short reminder: disordered systems

- Two types of variables, **dynamical** (i.e. spins) and **quenched** (geologically slow: atomic impurities, vacancies, etc.).
- Impurities  $\longrightarrow$  **random** Hamiltonian  $H_J$ . Site-dependent coupling constants (**sample**), extracted from some pdf  $P(J)$ .
- **Quenched** approximation: fast variables (spins) allowed **no backreaction** on slow ones. In practice:
  - 1 Take  $J$ -dependent thermal average  $\langle \dots \rangle_J$ .
  - 2 Average  $\langle \dots \rangle_J$  with  $P(J)$ .

# Short reminder: disordered systems

- Two types of variables, **dynamical** (i.e. spins) and **quenched** (geologically slow: atomic impurities, vacancies, etc.).
- Impurities  $\longrightarrow$  **random** Hamiltonian  $H_J$ . Site-dependent coupling constants (**sample**), extracted from some pdf  $P(J)$ .
- **Quenched** approximation: fast variables (spins) allowed **no backreaction** on slow ones. In practice:
  - 1 Take  $J$ -dependent thermal average  $\langle \dots \rangle_J$ .
  - 2 Average  $\langle \dots \rangle_J$  with  $P(J)$ .
- Metastability in single sample already bad enough, yet even worse...



# Short reminder: disordered systems

- Two types of variables, **dynamical** (i.e. spins) and **quenched** (geologically slow: atomic impurities, vacancies, etc.).
- Impurities  $\longrightarrow$  **random** Hamiltonian  $H_J$ . Site-dependent coupling constants (**sample**), extracted from some pdf  $P(J)$ .
- **Quenched** approximation: fast variables (spins) allowed **no backreaction** on slow ones. In practice:
  - 1 Take  $J$ -dependent thermal average  $\langle \dots \rangle_J$ .
  - 2 Average  $\langle \dots \rangle_J$  with  $P(J)$ .
- Metastability in single sample already bad enough, yet even worse...

**Sample-to-sample variance** of specific-heat, magnetic susceptibility, etc., expressed in natural units, is **diverging** in large  $L$  limit (long-tailed  $P(\chi_J/\overline{\chi_J})$ , Berche et al. 05)

# Short reminder: disordered systems

- Two types of variables, **dynamical** (i.e. spins) and **quenched** (geologically slow: atomic impurities, vacancies, etc.).
- Impurities  $\rightarrow$  **random** Hamiltonian  $H_J$ . Site-dependent coupling constants (**sample**), extracted from some pdf  $P(J)$ .
- **Quenched** approximation: fast variables (spins) allowed **no backreaction** on slow ones. In practice:
  - 1 Take  $J$ -dependent thermal average  $\langle \dots \rangle_J$ .
  - 2 Average  $\langle \dots \rangle_J$  with  $P(J)$ .
- Metastability in single sample already bad enough, yet even worse...


**Sample-to-sample variance** of specific-heat, magnetic susceptibility, etc., expressed in natural units, is **diverging** in large  $L$  limit

(long-tailed  $P(\chi_J/\overline{\chi_J})$ , Berche et al. 05)


$\Rightarrow$  need **huge** number of samples.

# Why rare events (Berche et al' 05)?

- Phase-coexistence:  
specific heat is  
 $C \sim L^D \propto N$   
 $(T - T_c \sim \frac{1}{N})$




# Why rare events (Berche et al' 05)?




- Phase-coexistence:  
specific heat is  
 $C \sim L^D \propto N$   
( $T - T_c \sim \frac{1}{N}$ )
- Critical region  
 $T - T_c \sim \frac{1}{\sqrt{N}}$   
Probability of phase  
coexistence  $\sim \frac{1}{\sqrt{N}}$

# Why rare events (Berche et al' 05)?




- Phase-coexistence:  
specific heat is  
 $C \sim L^D \propto N$   
( $T - T_c \sim \frac{1}{N}$ )
- Critical region  
 $T - T_c \sim \frac{1}{\sqrt{N}}$   
Probability of phase  
coexistence  $\sim \frac{1}{\sqrt{N}}$
- Rare events dominated:  
 $\bar{C} \propto N \times \frac{1}{\sqrt{N}} = \sqrt{N}$ ,  
(saturates Chayes bound)

# The microcanonical cure




- Disorder-averaged **entropy** (rather than free-energy)

# The microcanonical cure



- Disorder-averaged **entropy** (rather than free-energy)
- $\langle \hat{\beta} \rangle_e$  **self-averaging**

# The microcanonical cure



- Disorder-averaged **entropy** (rather than free-energy)
- $\langle \hat{\beta} \rangle_e$  **self-averaging**
- **Scaling analysis**: survival of first-order transition (Gordillo et al. 08).



# Outline

- 1 Introduction
- 2 Quasi constrained statistical ensembles
- 3 Local simulation algorithm
- 4 Cluster methods
- 5 First order phase transitions
- 6 Metastability in disordered systems
- 7 Conclusions**

# Conclusions

- (Quasi) **constrained** ensembles (for **energy** or other **reaction** coordinate) can be simulated **easily**.

# Conclusions

- (Quasi) **constrained** ensembles (for **energy** or other **reaction** coordinate) can be simulated **easily**.
- We visit **crucial** parts of phase space, with **exponentially small canonical probabilities**.

# Conclusions

- (Quasi) **constrained** ensembles (for **energy** or other **reaction** coordinate) can be simulated **easily**.
- We visit **crucial** parts of phase space, with **exponentially small canonical probabilities**.
- Fluctuation-Dissipation formalism allows to perform a **fully canonical** analysis.

# Conclusions

- (Quasi) **constrained** ensembles (for **energy** or other **reaction coordinate**) can be simulated **easily**.
- We visit **crucial** parts of phase space, with **exponentially small canonical probabilities**.
- Fluctuation-Dissipation formalism allows to perform a **fully canonical** analysis.
- **Cluster methods effective**, even for first-order phase transitions.

# Conclusions

- (Quasi) **constrained** ensembles (for **energy** or other **reaction coordinate**) can be simulated **easily**.
- We visit **crucial** parts of phase space, with **exponentially small canonical probabilities**.
- Fluctuation-Dissipation formalism allows to perform a **fully canonical** analysis.
- **Cluster methods effective**, even for first-order phase transitions.
- It is crucial to find **useful reaction coordinate**, see talk by B. Seoane on hard-spheres crystallization.

# Conclusions

- (Quasi) **constrained** ensembles (for **energy** or other **reaction coordinate**) can be simulated **easily**.
- We visit **crucial** parts of phase space, with **exponentially small canonical probabilities**.
- Fluctuation-Dissipation formalism allows to perform a **fully canonical** analysis.
- **Cluster methods effective**, even for first-order phase transitions.
- It is crucial to find **useful reaction coordinate**, see talk by B. Seoane on hard-spheres crystallization.
- **Geometric** transitions: metastability remains.

# Conclusions

- (Quasi) **constrained** ensembles (for **energy** or other **reaction coordinate**) can be simulated **easily**.
- We visit **crucial** parts of phase space, with **exponentially small canonical probabilities**.
- Fluctuation-Dissipation formalism allows to perform a **fully canonical** analysis.
- **Cluster methods effective**, even for first-order phase transitions.
- It is crucial to find **useful reaction coordinate**, see talk by B. Seoane on hard-spheres crystallization.
- **Geometric** transitions: metastability remains.

Angioletti-Uberti et al. 2010: promising cure, order parameters defined in half the simulation box.



# Conclusions

- (Quasi) **constrained** ensembles (for **energy** or other **reaction coordinate**) can be simulated **easily**.
- We visit **crucial** parts of phase space, with **exponentially small canonical probabilities**.
- Fluctuation-Dissipation formalism allows to perform a **fully canonical** analysis.
- **Cluster methods effective**, even for first-order phase transitions.
- It is crucial to find **useful reaction coordinate**, see talk by B. Seoane on hard-spheres crystallization.
- **Geometric** transitions: metastability remains.  
Angioletti-Uberti et al. 2010: promising cure, order parameters defined in half the simulation box.
- For **disordered systems**, these methods allow for a **redefinition** of the quenched averaged: we **cured the rare events syndrome**.