

MEDICAL CONDITIONS AND CRITICAL ILLNESS

GUIDE FOR INDIVIDUAL LIFE

Sun Life of Canada (Philippines), Inc.
October 1, 2019

The following chart outlines potential underwriting decisions for many commonly encountered medical conditions that might result in a rated, decline or postpone decision for life insurance coverage.

Decisions on medical conditions depend on a wide range of factors, and it is possible that after review of all underwriting evidences, the final underwriting decision may be different from the potential underwriting decision quoted in this guide.

This list is intended to be used as a quick reference guide only and is subject to change.

Glossary:

- **Standard** means clients are charged the standard premium rate quoted in the proposal.
- **Standard to Rated** means the best cases will be accepted at standard rates but some will be rated.
- **Rated** means the client has an above average risk and needs to pay additional premium to cover the extra mortality risk.
- **Postpone** means the client's application is not approved temporarily until more information can be obtained or a predetermined time has passed or until factors that had a bearing in the underwriting decision has changed. If a time requirement applies (example: postponed until age 10 or postponed for 1 year), the time frame given is the earliest we can consider the client.
- **Decline** means the client poses an uninsurable risk and no offer will be made.
- **Individual consideration (IC)** means that depending on the nature of the case, we may be able to offer insurance in some instances; however, in others, the case may be declined.

Medical condition	Potential underwriting decision for individual life coverage	Potential underwriting decision for critical illness coverage	Potential underwriting decision for staged products (Sun Fit and Well)
A			
Angina (blocked arteries of the heart)	See Coronary Artery Disease	See Coronary Artery Disease	See Coronary Artery Disease
Alzheimer's disease or other dementia	Decline	Decline	Decline
Angioplasty	See Bypass	See Bypass	See Bypass
Aneurysm (of the abdominal aorta)			
• Currently under age 50	Decline	Decline	Decline
• Poor control of cardiac risk factors or currently smoking	Decline	Decline	Decline
• Others	Individual consideration	Individual consideration	Decline
Anorexia Nervosa / Bulimia (eating disorder)			
• Under age 21 or within 3 years	Postpone	Postpone	Current age < or = 18 years old: Decline
• Over age 21, fully recovered 3 to 10 years ago, not underweight, no psychiatric disorder	Rated	Individual consideration	Over 18 years old: Individual consideration
• Recovered over 10 years ago	Standard	Individual consideration	Individual consideration

Anxiety			
• Currently disabled	Decline	Decline	Individual consideration
• Others	Standard to Rated	Individual consideration	Individual consideration
Arrhythmia			
• With history of stroke, transient ischemic attack, heart valve disease, coronary heart disease or cardiomyopathy	Decline	Decline	Decline
• Others, investigated, no apparent heart disease, stable	Standard to Rated	Individual consideration	Individual consideration
Artery disease, peripheral (lower legs)			
• Under age 40	Decline	Decline	Decline
• Currently smoking	Decline	Decline	Decline
• With other arterial disease (cerebral or heart)	Decline	Decline	Decline
• With diabetes	Individual consideration	Decline	Decline
• Age 40 and up, non-smoker, mild to moderate disease, stable, no complications, no other arterial disease (heart or cerebral), no diabetes	Rated	Decline	Decline
Asperger's syndrome (high functioning autism)			
• Up to age 10	Postpone	Postpone	< 18 years old: Postpone
• Age 11 and up, no behavior problems or mental or developmental disorders	Standard	Rated	> 18 years old: Individual consideration
• Others	See Autism	See Autism	See Autism
Asthma			
• Poorly controlled	Decline	Decline	Decline
• Mild, no history of hospitalization, well controlled with inhalers and/or bronchodilators, no oral steroids, smoking or not smoking	Standard	Standard to Rated	Individual consideration
• Others	Rated	Individual consideration	Individual consideration
Atrial fibrillation			
• With history of stroke, transient ischemic attack (TIA), heart valve disease, coronary heart disease or cardiomyopathy	Decline	Decline	Decline
• Others, stable	Individual consideration	Individual consideration	Individual consideration
Attention Deficit Disorder			
• Mild, no treatment or single medication with good response	Standard	Standard	Individual consideration
• Moderate, taking two medications, under age 10	Postpone	Postpone	Individual consideration
• Moderate, age 11 and up, taking two medications with good	Rated	Individual consideration	Individual consideration

response			
• Others	Decline	Decline	Decline
Autism			
• Up to age 15	Postpone	Decline	< 18 years old: Postpone
• Age 16 and up, best cases, no more than mild developmental delay, satisfactory social adaptation, no other medical problem	Rated	Decline	> 18 years old: Individual consideration
• Others	Decline	Decline	Decline
• Asperger's syndrome	See Asperger's syndrome	See Asperger's syndrome	See Asperger's syndrome
B			
Bipolar disorder	See Depression	See Depression	See Depression
Birth weight less than 5lbs	See Premature birth	See Premature birth	See Premature birth
Blood clot in the legs (deep vein thrombosis)			
• Single attack, within 3 months	Postpone	Postpone	Individual consideration
• Recurrent attacks, last one within 6 months	Postpone	Postpone	Individual consideration
• Others, no complications, no blood coagulation disorder	Standard to Rated	Individual consideration	Individual consideration
Blood pressure			
• Well controlled	Standard	Standard	Standard
• Fair control, mild to moderate elevation	Rated	Individual consideration	Individual consideration
• Poor control and/or non compliant to treatment	Decline	Decline	Decline
Brain tumour (benign)			
• Completely removed and confirmed benign	Individual consideration	Individual consideration	Individual consideration
• Small, no symptoms, no surgery required	Individual consideration	Individual consideration	Individual consideration
Bulimia			
• Recovered within 1 year	Postpone	Postpone	< 2 years: Postpone
• Recovered within 1 to 4 years	Rated	Postpone	Individual consideration
• Recovered within > 4 years	Standard	Individual consideration	Individual consideration
Bypass or angioplasty (coronary)			
• 0 – 6 months since surgery	Postpone	Decline	Decline
• Age at onset less than 35	Decline	Decline	Decline
• Recurrence of angina (chest pains)	Decline	Decline	Decline
• With other disease of the arteries (cerebral or limbs)	Decline	Decline	Decline
• With type 1 diabetes (insulin-dependent diabetes)	Decline	Decline	Decline
• With type 2 diabetes (non-insulin-dependent diabetes) and smoking	Decline	Decline	Decline

With type 2 diabetes (non-insulin-dependent diabetes) and non-smoker	Rated to Decline	Decline	Decline
Others	Rated to Decline	Decline	Decline
C			
Cancer	See table for cancers that are most often decline	See table for cancers that are most often decline	See table for cancers that are most often decline
Cardiac Failure	See Heart Failure	See Heart Failure	See Heart Failure
Cerebral palsy			
Less than 9 years old	Postpone	Postpone	< 16 years old: Postpone
Ages 9 and up, mild physical disability, normal mental development	Standard	Individual consideration	>16 years old: Individual consideration
Ages 9 and up, mild to moderate physical disability, mild developmental delay	Individual consideration	Decline	Individual consideration
Others	Decline	Decline	Decline
Cerebrovascular accident (CVA)	See Stroke	See Stroke	See Stroke
Cholesterol, high			
Well controlled with or without medication	Standard	Standard	Standard
Fair control, moderate elevation of cholesterol	Rated	Rated	Rated
Poor control, non-compliant, cholesterol very elevated	Decline	Decline	Decline
Chronic obstructive pulmonary disease (COPD)			
Severe, symptoms with slight exertion or at rest	Decline	Decline	Decline
Moderate, symptoms with moderate exertion	Postpone	Decline	Smoker: Individual consideration Non – smoker: Rated with exclusion
Mild, symptoms with strenuous exertion	Rated	Individual consideration	Smoker and Non – smoker: Rated with exclusion
Minimal, little or no symptoms	Standard to Rated	Standard to Rated	Smoker and Non – smoker: Rated with exclusion
Cirrhosis			
Currently drinking alcohol	Decline	Decline	Decline
Others, including those with liver biopsy results	Individual consideration	Individual consideration	Decline
Cleft Palate (cleft lip)			
Condition present, under age 2	Postpone	Postpone	Individual consideration
Condition present, age 2 and up	Individual consideration	Individual consideration	Individual consideration
Successful surgical repair, no complications, no residuals (all ages)	Standard	Standard	Individual consideration

Coronary artery disease (blocked arteries of the heart)			
• 0 – 6 months from onset of symptoms or surgery	Postpone	Decline	Decline
• Age at onset less than 35	Decline	Decline	Decline
• Unstable angina	Postpone	Decline	Decline
• With other disease of the arteries (cerebral or limbs)	Decline	Decline	Decline
• With type 1 diabetes (insulin-dependent diabetes)	Decline	Decline	Decline
• With type 2 diabetes (non-insulin-dependent diabetes) and smoking	Decline	Decline	Decline
• With type 2 diabetes (non-insulin-dependent diabetes), non-smoker for more than 12 months	Rated to decline	Decline	Decline
• Others	Rated to decline	Decline	Decline
Crohn's disease			
• Severe, within 5 years	Decline	Decline	Postpone to Rated with exclusion
• With surgery, within 6 months	Postpone	Postpone	Within 1 year: postpone
• Mild, no symptoms for at least 5 years	Standard	Individual consideration	Postpone to Rated with exclusion
• With surgery, no recurrence and no symptoms for at least 2 years	Standard	Individual consideration	Individual consideration
• Others	Rated	Decline	Decline
Cystic fibrosis			
• Under age 20	Decline	Decline	Decline
• Age 20 and up	Rated to Decline	Decline	Decline
D			
Depression (Major) or bipolar disorder			
• Within 1 year or currently disabled	Decline	Decline	Postpone
• Functioning well socially and in occupation 1 to 5 years	Individual consideration	Individual consideration	Individual consideration
• No symptoms 5 years up	Standard to Rated	Individual consideration	Individual consideration
Diabetes			
• Age 10 or younger	Decline	Decline	Decline
• Poorly controlled, non-compliant with treatment, or with complications such as renal disease, disease of the arteries, heart attack or stroke	Decline	Decline	Decline
• Ages 40 or older, non-smoker well controlled on diet alone, no complications	Standard	Individual consideration	Individual consideration
• Others	Rated	Decline	Decline
Dialysis, renal	Individual consideration	Decline	Decline

Down Syndrome			
• Up to age 10	Decline	Decline	Decline
• Ages 11 to 40, mild cases, stable environment, no other apparent disease, at school or gainfully employed	Rated	Decline	Decline
• Others, including those over age 40	Decline	Decline	Decline
E			
Emphysema	See Chronic obstructive pulmonary disease	Decline	Decline
Epilepsy			
• Within one year	Postpone	Postpone	Postpone
• Poorly controlled	Decline	Decline	Decline
• Well controlled with medication	Rated	Individual consideration	Individual consideration
• Seizure free for at least 5 years	Standard	Individual consideration	Individual consideration
• With surgery, no further seizures, 1 year up	Standard	Individual consideration	Individual consideration
Esophageal Varices	Decline	Decline	Decline
H			
Heart attack (myocardial infarct)	See Coronary artery disease	Decline	Decline
Heart failure, congestive			
• Present or within 2 years	Postpone	Postpone	Decline
• Over 2 years ago, underlying cause successfully treated, no recurrence of heart failure	Individual consideration	Individual consideration	Decline
Heart Transplant	Decline	Decline	Decline
Heart valve disease (i.e. aortic stenosis, mitral regurgitation, etc.)			
• Severe and unoperated	Decline	Decline	Decline
• Valvuloplasty, balloon or surgical, within 1 year	Decline	Decline	Decline
• With atrial fibrillation or coronary disease	Decline	Decline	Decline
• Pulmonary stenosis, under age 1	Decline	Decline	Decline
• Others	Individual consideration	Individual consideration	Individual consideration
Heart Valve Replacement			
• Within 1 year or awaiting surgery	Postpone	Decline	Decline
• Replacement prior to age 26	Decline	Decline	Decline
• Tricuspid valve replacement	Decline	Decline	Decline
• Others	Rated to Decline	Decline	Decline
Hemophilia			
• Mild, under age 20 and within 2 years of diagnosis	Postpone	Decline	Decline
• Moderate or severe, under age 20	Postpone	Decline	Decline
• Others(including chronic hepatitis)	Rated	Decline	Decline
Hepatitis B			
• Acute hepatitis B within the last	Postpone	Postpone	Postpone

6 months			
Acute hepatitis B, full recovery for at least 6 months (virus no longer present)	Standard	Individual consideration	Individual consideration
Others	Individual consideration	Individual consideration	Individual consideration
Hepatitis C			
Newly diagnosed or treated within 6 months	Postpone	Postpone	Decline
Treated after 6 months but with abnormal liver enzymes	Declined	Decline	Decline
Full recovery, normal liver enzymes, virus no longer detectable	Standard	Individual consideration	Individual consideration
Others	Rated to Decline	Individual consideration	Individual consideration
High blood pressure	See Blood pressure, elevated	See Blood pressure, elevated	See blood pressure, elevated
Huntington's chorea			
Present	Decline	Decline	Decline
Family history only, currently less than 60 years of age	Rated	Decline	Individual consideration
Family history only, currently age 60 and over	Standard	Decline	Individual consideration
Family history only with negative genetic test (all ages)	Standard	Individual Consideration	Individual consideration
Hydrocephalus			
Under age 10	Postpone	Postpone	Postpone
Within 3 years of shunt installation or revision	Postpone	Postpone	Decline
With abnormal development or mental function	Decline	Decline	Decline
Others	Rated	Decline	Individual consideration
Hypertension	See Blood Pressure, elevated	See Blood Pressure, elevated	See Blood Pressure, elevated
K			
Kidney transplant donor			
Not fully recovered or abnormal kidney function	Decline	Decline	Individual consideration
Fully recovered, normal kidney function	Standard	Standard	Standard
Kidney transplant recipient			
Age 25 or under at time of transplant	Decline	Decline	Decline
Within 2 years of transplant	Decline	Decline	Decline
Others	Individual consideration	Decline	Decline
L			
Liver function tests, elevated			
Severe elevation	Decline	Decline	Postpone
Moderate elevation, cause unknown	Decline	Decline	Rated
Mild elevation, cause unknown	Standard	Individual consideration	Rated

Mild or moderate elevation, cause known	Individual consideration	Individual consideration	Individual consideration
Liver transplant recipient	Decline	Decline	Individual consideration
Lou Gehrig's disease (amyotrophic lateral sclerosis)	Decline	Decline	Decline
Lung transplant	Decline	Decline	Decline
Lupus (SLE)			
Severe active disease or with kidney, heart, central nervous system, lung or blood involvement	Decline	Decline	Decline
Active disease, within 1 year	Decline	Postpone	Decline
Others	Individual consideration	Individual consideration	Decline
M			
Mammogram and/or breast ultrasound, abnormal			
Suspicious findings requiring repeat mammogram or additional testing	Postpone	Postpone	Postpone
Others	Individual consideration	Individual consideration	Individual consideration
Mental Retardation	See Developmental delay	See Developmental delay	See developmental delay
Multiple Sclerosis			
Within 1 year of diagnosis	Decline	Decline	Decline
Significant disability, unable to walk without aid or wheelchair bound, with cerebellar symptoms or bladder or bowel involvement	Decline	Decline	Decline
Progressive disease	Decline	Decline	Decline
Others, stable, fully ambulatory, may have mild residuals	Rated	Decline	Decline
Muscular dystrophy			
Limb girdle, Becker's/Benign X-linked	Rated to decline	Decline	Individual consideration
Fascioscapulohumeral, oculomuscular types (eye muscles only)	Individual consideration	Individual consideration	Individual consideration
Others or type unknown	Decline	Decline	Decline
- Family history only: - Negative genetic testing - No genetic testing, under age 20 - No genetic testing, age 20 to 45 - No genetic testing, age over 45, with favourable doctor's report - See note on genetic testing* at the end of this table	Standard	Standard	Individual consideration
	Decline	Decline	Individual consideration
	Individual consideration	Individual consideration	Individual consideration
	Standard	Standard	Individual consideration

Myocardial infarct (MI)	See Coronary artery disease	See Coronary artery disease	See Coronary artery disease
O			
Osteogenesis imperfecta (osteitis fragilitas)			
• Up to age 17	Postpone	Decline	Decline
• Age 18 and up	Rated to Decline	Decline	Decline
P			
Pacemaker			
• Within 1 year from implantation	Postpone	Postpone	Decline
• Under age 18	Postpone	Postpone	Decline
• Associated with cardiomyopathy, heart attack or heart valve disease	Decline	Decline	Decline
• Others	Rated to Decline	Individual consideration	Individual consideration
Pancreatitis			
• Acute, not related to alcohol, within 6 months	Postpone	Postpone	Postpone
• Acute, related to alcohol, within 2 years or still drinking	Decline	Decline	Decline
• Chronic, within 2 years or currently using alcohol	Decline	Decline	Decline
• Others, recovered, no complications	Standard to Rated	Standard to Rated	Rated to Individual consideration
Parkinson's disease	Rated to Decline	Decline	Individual consideration to Decline
Pap test, abnormal			
• Dysplasia (pre-cancerous), not compliant with recommended treatment or follow-up	Postpone	Postpone	Postpone
• Dysplasia (pre-cancerous), compliant with recommended treatment and follow-ups	Standard	Individual consideration	Standard to Rated
• Benign abnormalities, with or without normal follow-up pap test	Standard	Standard	Rated to Postpone
Paraplegia (wheelchair bound)			
• Within 2 years of onset; or not well-adjusted or not yet self-supporting	Postpone	Decline	Decline
• Without bowel and bladder control, under age 20	Postpone	Decline	Decline
• Others	Rated to Decline	Individual consideration	Rated to Decline
Pending Medical Investigations	Postpone	Postpone	Individual consideration
Pending Surgery	Individual consideration	Postpone	Individual consideration
Polycystic kidney disease			
• Present, up to age 44	Decline	Decline	Decline
• Present, age 45 and over	Rated to Decline	Decline	Decline

• Family history only:			
○ With negative DNA probe	Standard	Individual consideration	Standard
○ No DNA test, client up to age 19	Postpone	Postpone	Postpone
○ No DNA test, age 20 and up with investigation	Individual consideration	Postpone	Postpone
○ No DNA test, age up to 35, without investigation	Postpone	Postpone	Postpone
○ No DNA test, age 36 and up	Individual consideration	Postpone	Postpone
○ See note on genetic testing* at the end of this table			
Premature birth			
• Weight birth under 3.3 lbs (1.5 kg)			
○ Age under 12 months	Postpone	Postpone	Postpone
○ Age 12 months up and meeting their development milestones	Standard	Postpone to 24 months	Postpone to age 3 months
• Weight birth 3.3 lbs to 5 lbs (1.5 kg to 2.3 kg)			
○ Age under 6 months	Postpone	Postpone	Postpone
○ Age 6 months up and meeting their development milestones	Standard	Postpone to 18 months	Postpone to age 1
• Weight at birth greater than 5 lbs (2.3 kg), no complications	Standard	Standard	Postpone to age 1
Prostate-specific antigen (PSA), abnormal			
• Moderate to severe elevations	Decline	Decline	Decline
• Rising PSA, not investigated, under observation or pending biopsy	Postpone	Postpone	Postpone to Decline
• With negative ultrasound and/or biopsies	Individual consideration	Individual consideration	Individual consideration
• Mild elevation	Individual consideration	Individual consideration	Individual consideration
Q			
Quadriplegic	Decline	Decline	Decline
R			
Renal Failure (Chronic)			
• Mild stable impairment, normal blood pressure, near normal kidney function, no dialysis	Rated to Decline	Decline	Rated to Decline
• Others, including those with regular dialysis	Decline	Decline	Decline
Renal Failure (Acute), recovered			
• Within 6 months	Postpone	Postpone	Postpone
• Fully recovered at least 6 months, normal urinalysis and renal function	Standard	Postpone up to 2 years	Postpone to 1 year; Rated thereafter

Renal transplant recipient	See Kidney transplant recipient	See Kidney transplant recipient	See Kidney transplant, recipient
Rheumatoid arthritis			
• With heart, lung, vascular or nerve involvement	Decline	Decline	Decline
• Moderate to severe, stable with medication (such as methotrexate)	Rated	Rated	Individual consideration
• Mild, treatment with analgesics or mild non-steroidal anti-inflammatory drugs (NSAIDs) only	Standard	Standard	Individual consideration
S			
Schizophrenia or chronic psychotic disorders			
Within 2 years of last episode	Postpone	Decline	Postpone
Disabled or not functioning socially	Decline	Decline	Decline
Others, recovered, functioning socially and in occupation	Individual consideration	Decline	Postpone 5 years then Individual consideration
Scleroderma (systemic)			
• With 3 years of diagnosis	Postpone	Decline	Decline
• Heart, lung or kidney involvement	Decline	Decline	Decline
• Disease confined to the fingers and toes only	Standard	Rated	Postpone 5 years then Individual consideration
• Others	Rated to Decline	Decline	Decline
Sickle cell anaemia			
• Age less than 25	Postpone	Decline	Individual consideration
• Age 25 and up	Rated to Decline	Decline	Individual consideration
• All ages, sickle cell trait only, no anemia	Standard	Decline	Standard
Sleep apnea			
• Severe and untreated or severe and treated less than 6 months or severe and poor CPAP compliance	Decline	Decline	Rated
• Mild to moderate, untreated	Individual consideration	Individual consideration	Rated
• Treated with CPAP, good compliance, well controlled	Individual consideration	Individual consideration	Individual consideration
• Surgically treated, within 6 months	Postpone	Postpone	Individual consideration
• Successful surgery, 6 months up	Standard to Rated	Individual consideration	Individual consideration
Stroke (cerebrovascular accident)			
• Within 1 year	Postpone	Decline	Decline
• More than one stroke	Decline	Decline	Decline
• With diabetes, type I or II	Decline	Decline	Decline
• Smoker	Decline	Decline	Decline

• With history of heart attack, diabetes or poorly controlled hypertension	Decline	Decline	Decline
• Others	Individual consideration	Decline	Decline
Suicide attempt			
• Up to age 20, within 3 years of attempt	Postpone	Postpone	Postpone
• Age 21 and up, within 1 year of attempt	Postpone	Postpone	Postpone
• More than one attempt, last attempt within 5 years	Decline	Decline	Individual consideration
• Others, recovered with favourable prognosis (no substance abuse, no mental illness)	Individual consideration	Individual consideration	Rated
T			
Tetralogy of fallot			
• Without surgery or palliative surgery only	Decline	Decline	Decline
• With surgery, within 2 years	Postpone	Decline	Decline
• With surgery over 2 years ago	Rated to Decline	Decline	Decline
Thalassemia			
• Thalassemia carrier	Individual consideration	Individual consideration	Individual consideration
• Thalassemia trait or thalassemia minor of any type, minimal or no symptoms, mild anemia only	Individual consideration	Individual consideration	Standard
• Others	Individual consideration	Decline	Decline
Transient ischemic attack (TIA)			
• Single attack within 6 months	Postpone	Decline	Decline
• Single attack, 6 months to 5 years ago	Individual consideration	Decline	Decline
• Two attacks, last one within 1 year, or last attack up to age 40	Postpone	Decline	Decline
• Two attacks, last one within 1 year, or last attack at age 41 and above	Postpone	Decline	Decline
• More than 2 attacks	Decline	Decline	Decline
• With diabetes, type I or type II or with heart disease	Decline	Decline	Decline
• Others	Decline	Postpone	Decline
Transposition of the great vessels			
• Complete TGA (D Transposition)	Decline	Decline	Decline
• Congenitally corrected TGA, up to age 21	Decline	Decline	Decline
• Congenitally corrected TGA , age 22 and up	Individual consideration	Decline	Decline
U			
Ulcerative colitis			
• Severe attack within 2 years requiring hospitalization	Postpone	Decline	Decline
• Ongoing symptoms, within 1 year of diagnosis	Decline	Decline	Postpone

• Intermittent symptoms, within 6 months of diagnosis	Decline	Decline	Postpone
• Surgery within 6 months	Decline	Decline	Postpone for 1 year
• Surgically treated, 6 months and up	Rated	Individual consideration	Postpone for 1 year
• Others	Individual consideration	Individual consideration	Postpone (active disease)

***Genetic testing:** Sun Life Financial does not require a proposed insured to undergo genetic testing for an insurance application but if genetic testing has been done, it must be disclosed on the application