

# 1D ADCIRC Derivation

Tristan Dyer

March 7, 2018

## 1 Continuity Equation

Start with the vertically integrated continuity equation:

$$\frac{\partial H}{\partial t} + \frac{\partial}{\partial x}(UH) = 0 \quad (1)$$

where

$$H \equiv \zeta + h$$

$\zeta$  = free surface departure from the geoid

$h$  = bathymetric depth (distance from the geoid to the bottom)

$u$  = vertically varying velocity in the x-direction

$$U = \frac{1}{H} \int_{-h}^{\zeta} u dz = \text{depth-averaged velocity in the x-direction}$$

Take  $\partial/\partial t$  of (2):

$$\frac{\partial^2 H}{\partial t^2} + \frac{\partial}{\partial x} \frac{\partial}{\partial t}(UH) = 0 \quad (2)$$

Add (2) to (1) multiplied by the parameter  $\tau_0$ , which may be variable in space:

$$\begin{aligned} \frac{\partial^2 H}{\partial t^2} + \frac{\partial}{\partial x} \frac{\partial}{\partial t}(UH) + \tau_0 \left( \frac{\partial H}{\partial t} + \frac{\partial}{\partial x}(UH) \right) &= 0 \\ \frac{\partial^2 H}{\partial t^2} + \frac{\partial}{\partial x} \frac{\partial}{\partial t}(UH) + \tau_0 \frac{\partial H}{\partial t} + \tau_0 \frac{\partial}{\partial x}(UH) &= 0 \\ \frac{\partial^2 H}{\partial t^2} + \tau_0 \frac{\partial H}{\partial t} + \tau_0 \frac{\partial}{\partial x}(UH) + \frac{\partial}{\partial x} \frac{\partial}{\partial t}(UH) &= 0 \end{aligned} \quad (3)$$

Now, define  $\tilde{J}_x$ :

$$\tilde{J}_x \equiv \tau_0(UH) + \frac{\partial}{\partial t}(UH) \quad (4)$$

$$\tilde{J}_x = \tau_0 Q + \frac{\partial Q}{\partial t} \quad (5)$$

where

$$Q = UH$$

Recall that  $\tau_0$ ,  $U$ , and  $H$  are all variable in  $x$  and take  $\partial/\partial x$  of (5), noting the use of the product rule:

$$\begin{aligned} \frac{\partial \tilde{J}_x}{\partial x} &= \frac{\partial}{\partial x} \left[ \tau_0 Q + \frac{\partial Q}{\partial t} \right] \\ &= \frac{\partial}{\partial x}(\tau_0 Q) + \frac{\partial}{\partial x} \frac{\partial}{\partial t} Q \\ &= Q \frac{\partial \tau_0}{\partial x} + \tau_0 \frac{\partial Q}{\partial x} + \frac{\partial}{\partial x} \frac{\partial}{\partial t} Q \\ &= \tau_0 \frac{\partial Q}{\partial x} + \frac{\partial}{\partial x} \frac{\partial}{\partial t} Q + Q \frac{\partial \tau_0}{\partial x} \\ &= \tau_0 \frac{\partial(UH)}{\partial x} + \frac{\partial}{\partial x} \frac{\partial}{\partial t}(UH) + UH \frac{\partial \tau_0}{\partial x} \end{aligned} \quad (6)$$

Now, returning to equation (3), let's add zero to it in the form of:

$$\begin{aligned} UH \frac{\partial \tau_0}{\partial x} - UH \frac{\partial \tau_0}{\partial x} &= 0 \\ \frac{\partial^2 H}{\partial t^2} + \tau_0 \frac{\partial H}{\partial t} + \underbrace{\tau_0 \frac{\partial}{\partial x}(UH) + \frac{\partial}{\partial x} \frac{\partial}{\partial t}(UH) + UH \frac{\partial \tau_0}{\partial x}}_{\text{Note that this is equivalent to (6)}} - UH \frac{\partial \tau_0}{\partial x} &= 0 \end{aligned}$$

and substituting (6) in gives us:

$$\frac{\partial^2 H}{\partial t^2} + \tau_0 \frac{\partial H}{\partial t} + \frac{\partial \tilde{J}_x}{\partial x} - UH \frac{\partial \tau_0}{\partial x} = 0 \quad (7)$$

If we assume that bathymetric depth is constant, then

$$\frac{\partial H}{\partial t} = \frac{\partial \zeta}{\partial t}$$

$$\frac{\partial^2 H}{\partial t^2} = \frac{\partial^2 \zeta}{\partial t^2}$$

and (7) can be rewritten as

$$\frac{\partial^2 \zeta}{\partial t^2} + \tau_0 \frac{\partial \zeta}{\partial t} + \frac{\partial \tilde{J}_x}{\partial x} - UH \frac{\partial \tau_0}{\partial x} = 0 \quad (8)$$