

# NUMERICAL SIMULATIONS

František Dinnbier

NCTS (NTU)  
3rd July 2025

## Motivation for N-body dynamics

The problem is called N-body problem.

# Motivation for N-body dynamics

The problem is called N-body problem.

Lets' start with  $N = 2$ .

# Motivation for the N-body problem

- Let's start with two bodies
- The law of universal gravitation

$$\vec{F} = -\frac{GM_1 M_2 \vec{r}}{r^3}$$



# Motivation for the N-body problem

- Let's start with two bodies
- The law of universal gravitation


$$\vec{F} = -\frac{GM_1 M_2 \vec{r}}{r^3}$$



# Motivation for the N-body problem

- Let's start with two bodies
- The law of universal gravitation

$$\vec{F} = -\frac{GM_1 M_2 \vec{r}}{r^3}$$




## Two body problem - analytical solution

- The solution is well known

# Two body problem - analytical solution


- The solution is well known




# Two body problem - analytical solution

- The solution is well known
- 


$$r = \frac{a(1 - e^2)}{1 + e \cos \phi}$$




## Two body problem - numerical solution



## Two body problem - numerical solution




## Two body problem - numerical solution



- Two stars of masses  $1 M_{\odot}$  and  $0.5 M_{\odot}$  on a mildly eccentric orbit with semi-major axis  $a = 0.64 \text{ AU}$ .

# Two body problem - numerical solution



## The transition to complexity

Two body problem is well understood.

# The transition to complexity

Two body problem is well understood.

What happens if  $N = 3$ .

## The transition to complexity


Two body problem is well understood.

What happens if  $N = 3$ .

Is there also an elegant analytic solution as for  
 $N = 2$ ?


# Introduction to three body problem

- Lets have three stars of masses  $3 M_{\odot}$ ,  $4 M_{\odot}$  and  $5 M_{\odot}$  at rest at the beginning.
- The stars move only as the influence of its mutual gravity (i.e. star 2 is attracted only by star 1 and 3).




# Introduction to three body problem

- Lets have three stars of masses  $3 M_{\odot}$ ,  $4 M_{\odot}$  and  $5 M_{\odot}$  at rest at the beginning.
- The stars move only as the influence of its mutual gravity (i.e. star 2 is attracted only by star 1 and 3).



# Introduction to three body problem


- Lets have three stars of masses  $3 M_{\odot}$ ,  $4 M_{\odot}$  and  $5 M_{\odot}$  at rest at the beginning.
- The stars move only as the influence of its mutual gravity (i.e. star 2 is attracted only by star 1 and 3).
- It is difficult to imagine that the complex trajectories have analytic solution.



## The transition to complexity

What are the typical values of  $N$  in the nature?

# The ecosystem of a spiral galaxy




## Systems with larger N: open star clusters

Open star clusters contain up to several thousand of stars ( $N = 30$  to  $\approx 10^4$  ( $10^6$ );  $\log_{10}(N) = 1.5$  to 4).



## Systems with larger N: open star clusters

There is a large diversity in open star clusters (they differ in their sizes, ages, mass, . . . )




## Systems with large N: globular star clusters

Globular star clusters contain up to several million of stars ( $N \approx 10^3$  to  $\approx 10^7$ ;  $\log_{10}(N) = 3$  to 7).



# Stellar binaries and multiples



- Binary stars are ubiquitous;  
 $\approx 55\%$  of solar-type stars are binaries.
- Among  $15 M_{\odot}$  stars (spectral type O), almost all of them have a companion.

# Stellar binaries and multiples

A: K1V 0.84 M<sub>⊙</sub>  
B: DA4 White Dwarf 0.5 M<sub>⊙</sub>  
C: M4.5 Red Dwarf 0.2 M<sub>⊙</sub>



- Binary stars are ubiquitous;  
 $\approx 55\%$  of solar-type stars are binaries.
- Among  $15 M_{\odot}$  stars (spectral type O), almost all of them have a companion.
- Most of  $15 M_{\odot}$  stars have two companions, some have four, and a few even more.

# Stellar binaries and multiples




- Binary stars are ubiquitous;  
≈ 55 % of solar-type stars are binaries.
- Among  $15 M_{\odot}$  stars (spectral type O), almost all of them have a companion.
- Most of  $15 M_{\odot}$  stars have two companions, some have four, and a few even more.

# Stellar binaries and multiples




- Binary stars are ubiquitous;  
 $\approx 55\%$  of solar-type stars are binaries.
- Among  $15 M_{\odot}$  stars (spectral type O), almost all of them have a companion.
- Most of  $15 M_{\odot}$  stars have two companions, some have four, and a few even more.

## Stellar binaries and multiples




- Binary stars are ubiquitous;  
≈ 55 % of solar-type stars are binaries.
  - Among  $15 M_{\odot}$  stars (spectral type O), almost all of them have a companion.
  - Most of  $15 M_{\odot}$  stars have two companions, some have four, and a few even more.

# Stellar binaries and multiples



- Binary stars are ubiquitous;  $\approx 55\%$  of solar-type stars are binaries.
- Among  $15 M_{\odot}$  stars (spectral type O), almost all of them have a companion.
- Most of  $15 M_{\odot}$  stars have two companions, some have four, and a few even more.
- Also some of Solar-type stars have a third companion ( $\approx 7\%$ ).

# Stellar binaries and multiples



- Binary stars are ubiquitous;  $\approx 55\%$  of solar-type stars are binaries.
- Among  $15 M_{\odot}$  stars (spectral type O), almost all of them have a companion.
- Most of  $15 M_{\odot}$  stars have two companions, some have four, and a few even more.
- Also some of Solar-type stars have a third companion ( $\approx 7\%$ ).
- The Sun has no companion.

## Systems with huge N: galaxies

Milky Way galaxy contains  $\approx 5 \cdot 10^{10}$  stars  
 $(\log_{10}(N) = 10.7)$ .



# Zooming-in on the Andromeda galaxy (M 31)



Panchromatic Hubble Andromeda Treasury (PHAT) program

# Zooming-in on the Andromeda galaxy (M 31)



Panchromatic Hubble Andromeda Treasury (PHAT) program

# Zooming-in on the Andromeda galaxy (M 31)



Panchromatic Hubble Andromeda Treasury (PHAT) program

# Zooming-in on the Andromeda galaxy (M 31)



Panchromatic Hubble Andromeda Treasury (PHAT) program

# Zooming-in on the Andromeda galaxy (M 31)



Panchromatic Hubble Andromeda Treasury (PHAT) program

# Zooming-in on the Andromeda galaxy (M 31)




Panchromatic Hubble Andromeda Treasury (PHAT) program

# Zooming-in on the Andromeda galaxy (M 31)



Panchromatic Hubble Andromeda Treasury (PHAT) program

# Zooming-in on the Andromeda galaxy (M 31)



Panchromatic Hubble Andromeda Treasury (PHAT) program

# N-body integrators

- Each star is represented by one particle.

# N-body integrators

- Each star is represented by one particle.
- More detailed models include binary stars.

# N-body integrators

- Each star is represented by one particle.
- More detailed models include binary stars.
- The models can include stellar evolution from pre-calculated fitting formulae (stellar evolution is not calculated on-the-fly).

# N-body integrators


- Each star is represented by one particle.
- More detailed models include binary stars.
- The models can include stellar evolution from pre-calculated fitting formulae (stellar evolution is not calculated on-the-fly).
- Tidal field of the galaxy, molecular clouds, ...

# N-body integrators

- Each star is represented by one particle.
- More detailed models include binary stars.
- The models can include stellar evolution from pre-calculated fitting formulae (stellar evolution is not calculated on-the-fly).
- Tidal field of the galaxy, molecular clouds, ...
- → here we assume that all stars are point-mass (the integrating scheme is the most difficult and complex part of the numerical scheme).


# The initial conditions

- For each star, we know its position  $\vec{r}$ , velocity  $\vec{v}$ , and mass  $m$ .




# The initial conditions

- For each star, we know its position  $\vec{r}$ , velocity  $\vec{v}$ , and mass  $m$ .
- We need to determine the integration time-step.



# The initial conditions

- For each star, we know its position  $\vec{r}$ , velocity  $\vec{v}$ , and mass  $m$ .
- We need to determine the integration time-step.
- Each integration time-step, the position and velocity of the stars is updated.



## The types of N-body integration schemes

- Predictor-corrector (adaptive and individual time-steps).

## The types of N-body integration schemes

- Predictor-corrector (adaptive and individual time-steps).
- Leap frog (symplectic, the energy error does not increase with time).

## Overview of the predictor-corrector scheme

- Star 1 experiences the following acceleration

$$\vec{a}_1 = -\frac{GM_2\vec{r}_{1,2}}{r_{1,2}^3}$$

## Overview of the predictor-corrector scheme

- Star 1 experiences the following acceleration

$$\vec{a}_1 = -\frac{GM_2 \vec{r}_{1,2}}{r_{1,2}^3}$$

- Thus, the acceleration changes with time as

$$\dot{\vec{a}}_1 = -\frac{GM_2 \vec{v}_{1,2}}{r_{1,2}^3} + 3GM_2 \frac{\vec{v}_{1,2} \cdot \vec{r}_{1,2}}{r_{1,2}^5}$$

# Overview of the predictor-corrector scheme

- Star 1 experiences the following acceleration

$$\vec{a}_1 = -\frac{GM_2 \vec{r}_{1,2}}{r_{1,2}^3}$$

- Thus, the acceleration changes with time as

$$\dot{\vec{a}}_1 = -\frac{GM_2 \vec{v}_{1,2}}{r_{1,2}^3} + 3GM_2 \frac{\vec{v}_{1,2} \cdot \vec{r}_{1,2}}{r_{1,2}^5}$$

- From these formulae, we can predict the positions at time  $dt$

$$\vec{r}_p = \vec{r}_0 + \vec{v}_0 dt + \frac{1}{2} \vec{a}_0 (dt)^2 + \frac{1}{6} \vec{a}_0 \cdot$$

# Overview of the predictor-corrector scheme

- Star 1 experiences the following acceleration

$$\vec{a}_1 = -\frac{GM_2 \vec{r}_{1,2}}{r_{1,2}^3}$$

- Thus, the acceleration changes with time as

$$\dot{\vec{a}}_1 = -\frac{GM_2 \vec{v}_{1,2}}{r_{1,2}^3} + 3GM_2 \frac{\vec{v}_{1,2} \cdot \vec{r}_{1,2}}{r_{1,2}^5}$$

- From these formulae, we can predict the positions at time  $dt$

$$\vec{r}_p = \vec{r}_0 + \vec{v}_0 dt + \frac{1}{2} \vec{a}_0 (dt)^2 + \frac{1}{6} \dot{\vec{a}}_0 (dt)^3$$

- And velocities

$$\vec{v}_p = \vec{v}_0 + \vec{a}_0 dt + \frac{1}{2} \dot{\vec{a}}_0 (dt)^2$$

# Overview of the predictor-corrector scheme

- Star 1 experiences the following acceleration

$$\vec{a}_1 = -\frac{GM_2 \vec{r}_{1,2}}{r_{1,2}^3}$$

- Thus, the acceleration changes with time as

$$\dot{\vec{a}}_1 = -\frac{GM_2 \vec{v}_{1,2}}{r_{1,2}^3} + 3GM_2 \frac{\vec{v}_{1,2} \cdot \vec{r}_{1,2}}{r_{1,2}^5}$$

- From these formulae, we can predict the positions at time  $dt$

$$\vec{r}_p = \vec{r}_0 + \vec{v}_0 dt + \frac{1}{2} \vec{a}_0 (dt)^2 + \frac{1}{6} \vec{a}_0 \dot{(dt)}$$

- And velocities

$$\vec{v}_p = \vec{v}_0 + \vec{a}_0 dt + \frac{1}{2} \vec{a}_0 \dot{(dt)^2}$$

- Note that positions and velocities  $\vec{r}_p$  and  $\vec{v}_p$  are only predicted values; their accuracy can be further increased by correcting.

## Overview of the predictor-corrector scheme

- We evaluate acceleration ( $\vec{a}_1$ ) and its time derivative ( $\dot{\vec{a}}_1$ ) at the predicted positions at new time  $t_1 = t_0 + dt$ .

## Overview of the predictor-corrector scheme

- We evaluate acceleration ( $\vec{a}_1$ ) and its time derivative ( $\dot{\vec{a}}_1$ ) at the predicted positions at new time  $t_1 = t_0 + dt$ .
- It is reasonable to assume that the velocity and its derivatives change smoothly (lets drop the vector symbols for simplicity), i.e.

$$v(t_0) = v_0$$

$$\dot{v}(t_0) = a_0$$

$$\ddot{v}(t_0) = \dot{a}_0$$

$$\dot{v}(t_1) = a_1$$

$$\ddot{v}(t_1) = \dot{a}_1$$

## Overview of the predictor-corrector scheme

- Correcting step is achieved by fitting by Hermite polynomials:  
 $r(t_1) = r_p + 3/20(-a_0 + a_1)(dt)^2 - 1/60(7\dot{a}_0 + 2\dot{a}_1)(dt)^3$   
 $v(t_1) = v_p + 1/2(-a_0 + a_1)dt - 1/12(5\dot{a}_0 + \dot{a}_1)(dt)^2$

## The individual time-steps

- Time-step reflects the speed of change of the environment around the particular star.

## The individual time-steps

- Time-step reflects the speed of change of the environment around the particular star.
- Short time-step is necessary if the star undergoes rapid changes.

## The individual time-steps

- Time-step reflects the speed of change of the environment around the particular star.
- Short time-step is necessary if the star undergoes rapid changes.
- Experimental time-step criterions:

$$dt = \eta \frac{r_{\text{stars}}}{v_{\text{max}}}$$

$$dt = \eta \frac{F}{\ddot{F}}$$

$$dt = \sqrt{\eta \frac{F}{\ddot{F}}}$$

# The individual time-steps

- Time-step reflects the speed of change of the environment around the particular star.
- Short time-step is necessary if the star undergoes rapid changes.
- Experimental time-step criterions:

$$dt = \eta \frac{r_{\text{stars}}}{v_{\text{max}}}$$

$$dt = \eta \frac{F}{\dot{F}}$$


$$dt = \sqrt{\eta \frac{F}{\ddot{F}}}$$

- The best condition turned out to be (Aarseth 1985)

$$dt = \sqrt{\eta \frac{|F||\ddot{F}| + |\dot{F}|^2}{|\dot{F}||\ddot{F}| + |\ddot{F}|^2}}$$

# The individual time-steps

- time steps are generated by a factor of 2 like: ... it is like musical notes



→ particle 3 does not interact so much

→ particles 1 and 2 interact strongly → short time

→ the interval duration is one in N-body units

# Individual time-steps as the main challenge in star cluster modelling

- Star clusters evolve on the time-scale of Myrs to Gyrs.

# Individual time-steps as the main challenge in star cluster modelling

- Star clusters evolve on the time-scale of Myrs to Gyrs.
- Close binary stars in the cluster can have orbital periods as short as several hours → the necessary time-step is several minutes.

# Individual time-steps as the main challenge in star cluster modelling

- Star clusters evolve on the time-scale of Myrs to Gyrs.
- Close binary stars in the cluster can have orbital periods as short as several hours → the necessary time-step is several minutes.
- This would mean  $\approx 10^{14}$  individual integration time-steps for a single star.

# Individual time-steps as the main challenge in star cluster modelling

- Star clusters evolve on the time-scale of Myrs to Gyrs.
- Close binary stars in the cluster can have orbital periods as short as several hours → the necessary time-step is several minutes.
- This would mean  $\approx 10^{14}$  individual integration time-steps for a single star.
- Smart algorithms usually wins over brute force.

## Checking accuracy of N-body models

- Consider evolution of a system of N-point masses under its mutual gravity.

# Checking accuracy of N-body models

- Consider evolution of a system of N-point masses under its mutual gravity.
- Is there a way how to check that the simulation was correctly calculated?

# Checking accuracy of N-body models


- Consider evolution of a system of N-point masses under its mutual gravity.
- Is there a way how to check that the simulation was correctly calculated?
- Conservation of mechanical energy  $E_{\text{tot}} = E_k + E_p$ .

# Checking accuracy of N-body models


- Consider evolution of a system of N-point masses under its mutual gravity.
- Is there a way how to check that the simulation was correctly calculated?
- Conservation of mechanical energy  $E_{\text{tot}} = E_k + E_p$ .
- Relative energy error:

$$\text{d}E = |E_{\text{tot}} - E_{\text{tot},0}| / |E_{\text{tot},0}|$$

# Checking accuracy of N-body models




# Checking accuracy of N-body models




# The sensitivity on initial conditions

$t = 15.5608$  years



$t = 15.5608$  years



Both simulations initially have the same initial conditions with the difference that the  $3 M_{\odot}$  star in the model on the right panel has initial speed of 10cm/s.

## The sensitivity on initial conditions

- A tiny difference in initial conditions leads to a large difference at a later time (the outgoing binary was formed from different stars).

## The sensitivity on initial conditions

- A tiny difference in initial conditions leads to a large difference at a later time (the outgoing binary was formed from different stars).
- Divergence of close trajectories → typical for chaotic systems.


## The sensitivity on initial conditions

- A tiny difference in initial conditions leads to a large difference at a later time (the outgoing binary was formed from different stars).
- Divergence of close trajectories → typical for chaotic systems.
- Initially nearby trajectories diverge as

$$x_p - x_r \propto e^{\lambda t},$$


where  $\lambda$  is a positive number.

# Performance of supercomputers



Speed of computers doubled every 18 months

# The past and future of N-body calculations



- Speed of computers doubles every 18 months (Moore's law).
- In 10 years, it is possible to calculate only 10 times more massive clusters.
- Modelling of globular star clusters is currently possible.

## A general note on performing numerical simulations

- Alongside observations and theory, numerical modelling is a powerful tool in science.
- Numerical algorithms are often very efficient on the task which they were developed.
- Numerical algorithms can become inefficient or provide wrong results when incorrectly understood and setup.
- → avoid using a code as a black box (for doing real science, experiments are OK); it is better to understand at least the basics of the numerical scheme, its strengths and weaknesses.

Thank you for your attention