

# Introduction to Metric Spaces

TA: Nate Clause

# What is a Metric Space

## Definition

A *metric space* is a pair  $(X, d_X)$ , where  $X$  is a set, and  $d_X : X \times X \rightarrow \mathbb{R}$  is a function, called the *metric* or *distance*, satisfying:

# What is a Metric Space

## Definition

A *metric space* is a pair  $(X, d_X)$ , where  $X$  is a set, and  $d_X : X \times X \rightarrow \mathbb{R}$  is a function, called the *metric* or *distance*, satisfying:

- (i) (identity of indiscernibles)  $d_X(x, y) = 0$  if and only if  $x = y$ .

# What is a Metric Space

## Definition

A *metric space* is a pair  $(X, d_X)$ , where  $X$  is a set, and  $d_X : X \times X \rightarrow \mathbb{R}$  is a function, called the *metric* or *distance*, satisfying:

- (i) (identity of indiscernibles)  $d_X(x, y) = 0$  if and only if  $x = y$ .
- (ii) (symmetry)  $d_X(x, y) = d_X(y, x)$  for all  $x, y \in X$ .

# What is a Metric Space

## Definition

A *metric space* is a pair  $(X, d_X)$ , where  $X$  is a set, and  $d_X : X \times X \rightarrow \mathbb{R}$  is a function, called the *metric* or *distance*, satisfying:

- (i) (identity of indiscernibles)  $d_X(x, y) = 0$  if and only if  $x = y$ .
- (ii) (symmetry)  $d_X(x, y) = d_X(y, x)$  for all  $x, y \in X$ .
- (iii) (triangle inequality) For all  $x, y, z \in X$ , we have:

$$d_X(x, z) \leq d_X(x, y) + d_X(y, z)$$

# What is a Metric Space

## Definition


A *metric space* is a pair  $(X, d_X)$ , where  $X$  is a set, and  $d_X : X \times X \rightarrow \mathbb{R}$  is a function, called the *metric* or *distance*, satisfying:

- (i) (identity of indiscernibles)  $d_X(x, y) = 0$  if and only if  $x = y$ .
- (ii) (symmetry)  $d_X(x, y) = d_X(y, x)$  for all  $x, y \in X$ .
- (iii) (triangle inequality) For all  $x, y, z \in X$ , we have:


$$d_X(x, z) \leq d_X(x, y) + d_X(y, z)$$

- If  $X$  is understood, sometimes the subscript is dropped and we just write  $d$ .

# What is a Metric Space



# What is a Metric Space





# Non-negativity

**Claim:** If  $(X, d_X)$  is a metric space, then for all  $x, y \in X$ ,  $d(x, y) \geq 0$ .

# Non-negativity

**Claim:** If  $(X, d_X)$  is a metric space, then for all  $x, y \in X$ ,  $d(x, y) \geq 0$ .

**Proof.**

Let  $x, y \in X$ . Then we have:

$$2d(x, y) \stackrel{(ii)}{=} d(x, y) + d(y, x) \stackrel{(iii)}{\geq} d(x, x) \stackrel{(i)}{=} 0$$



# Metric Space Example

- $X = \mathbb{R}$ ,  $d : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  given by  $d(x, y) = |x - y|$ . We show that this is a metric space:

Proof.

To see (i):  $|x - y| = 0 \iff x - y = 0 \iff x = y$ .

To see (ii):  $x - y = -(y - x) \implies |x - y| = |y - x|$ .

# Metric Space Example

- $X = \mathbb{R}$ ,  $d : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  given by  $d(x, y) = |x - y|$ . We show that this is a metric space:

Proof.

To see (i):  $|x - y| = 0 \iff x - y = 0 \iff x = y$ .

To see (ii):  $x - y = -(y - x) \implies |x - y| = |y - x|$ . As an exercise,

show that for all  $x, y \in \mathbb{R}$ ,  $|x + y| \stackrel{*}{\leq} |x| + |y|$ . Using this fact, for all  $x, y, z \in \mathbb{R}$ , we have:

$$\begin{aligned} d(x, z) &= |x - z| = |(x - y) + (y - z)| \\ &\stackrel{*}{\leq} |x - y| + |y - z| = d(x, y) + d(y, z) \end{aligned}$$



## Definition

Given a vector space  $X$  over  $\mathbb{R}$ , a *norm* on  $X$  is a function  $\| \cdot \| : X \rightarrow \mathbb{R}$  satisfying:

## Definition

Given a vector space  $X$  over  $\mathbb{R}$ , a *norm* on  $X$  is a function  $\| \cdot \| : X \rightarrow \mathbb{R}$  satisfying:

(i)  $\|x + y\| \leq \|x\| + \|y\|$  for all  $x, y \in X$ .

## Definition

Given a vector space  $X$  over  $\mathbb{R}$ , a *norm* on  $X$  is a function  $\| \cdot \| : X \rightarrow \mathbb{R}$  satisfying:

- (i)  $\|x + y\| \leq \|x\| + \|y\|$  for all  $x, y \in X$ .
- (ii)  $\|c \cdot x\| = |c|\|x\|$  for all  $x \in X, c \in \mathbb{R}$ .

## Definition

Given a vector space  $X$  over  $\mathbb{R}$ , a *norm* on  $X$  is a function  $\| \cdot \| : X \rightarrow \mathbb{R}$  satisfying:

- (i)  $\|x + y\| \leq \|x\| + \|y\|$  for all  $x, y \in X$ .
- (ii)  $\|c \cdot x\| = |c|\|x\|$  for all  $x \in X$ ,  $c \in \mathbb{R}$ .
- (iii) For all  $x \in X$ ,  $\|x\| = 0$  if and only if  $x = 0$ .



# Norm Examples

- The function  $\| \cdot \| : \mathbb{R} \rightarrow \mathbb{R}$  given by  $\|x\| = |x|$ .

# Norm Examples

- The function  $\| \cdot \| : \mathbb{R} \rightarrow \mathbb{R}$  given by  $\|x\| = |x|$ .
- The Euclidean norm:  $\| \cdot \| : \mathbb{R}^n \rightarrow \mathbb{R}$  given by  $\|x\| = \sqrt{x \cdot x}$ , where  $x \cdot x$  refers to the *dot product*.

An important class of norms are the  $p$ -norms:

### Definition

For  $n \in \mathbb{N}$ ,  $p \in \mathbb{R}$  with  $p \in [1, \infty)$ , the  $p$ -norm is a norm on  $\mathbb{R}^n$  given by:

$$\|x\|_p := \left( \sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}$$

An important class of norms are the  $p$ -norms:

### Definition

For  $n \in \mathbb{N}$ ,  $p \in \mathbb{R}$  with  $p \in [1, \infty)$ , the  $p$ -norm is a norm on  $\mathbb{R}^n$  given by:

$$\|x\|_p := \left( \sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}$$

Examples:

- If  $n = 1$ ,  $p = 1$ , we get the absolute value norm on  $\mathbb{R}$ .
- If  $n \in \mathbb{N}$ ,  $p = 2$ , we get the Euclidean norm.

## Definition

For  $n \in \mathbb{N}$ ,  $p = \infty$ , the  $p$ -norm is a norm on  $\mathbb{R}^n$  given by:

$$\|x\|_{\infty} := \left( \max_{1 \leq i \leq n} |x_i| \right)$$

## Proposition

Let  $X$  be a vector space over  $\mathbb{R}$  and  $\|\cdot\|$  a norm on  $X$ . Define  $d_X : X \times X \rightarrow \mathbb{R}$  by  $d_X(x, y) := \|x - y\|$ . Then  $(X, d_X)$  is a metric space.

Proof: **Exercise**

# Minkowski distances

## Definition

For  $n \in \mathbb{N}$ ,  $p \in [1, \infty]$ , the *Minkowski distance of order  $p$*  is given by  $d_p : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ , where

$$d_p(x, y) := \|x - y\|_p.$$

# Minkowski distances

## Definition

For  $n \in \mathbb{N}$ ,  $p \in [1, \infty]$ , the *Minkowski distance of order  $p$*  is given by  $d_p : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ , where

$$d_p(x, y) := \|x - y\|_p.$$

## Example

If  $n = 2$ ,  $p = 2$ , then we have:

$$d_2((x_1, y_1), (x_2, y_2)) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$



# Minkowski distances

## Definition

For  $n \in \mathbb{N}$ ,  $p \in [1, \infty]$ , the *Minkowski distance of order  $p$*  is given by  $d_p : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ , where

$$d_p(x, y) := \|x - y\|_p.$$

## Example

If  $n = 2$ ,  $p = 2$ , then we have:

$$d_2((x_1, y_1), (x_2, y_2)) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

## Corollary

For  $p \in [1, \infty]$ ,  $d_p : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  is a metric.

# Unit spheres in Minkowski metric spaces

- For  $n \in \mathbb{N}$ ,  $p \in [1, \infty]$ , let  $(\mathbb{R}^n, d_p)$  be the Minkowski metric space of order  $p$ . The *unit sphere* is defined as:


$$\{x \in \mathbb{R}^n \mid d_p(x, 0) = 1\}$$

# Unit spheres in Minkowski metric spaces

- For  $n \in \mathbb{N}$ ,  $p \in [1, \infty]$ , let  $(\mathbb{R}^n, d_p)$  be the Minkowski metric space of order  $p$ . The *unit sphere* is defined as:

$$\{x \in \mathbb{R}^n \mid d_p(x, 0) = 1\}$$

- Unit sphere with  $n = 2$ ,  $p = 1$ :




# Unit spheres in Minkowski metric spaces

- For  $n \in \mathbb{N}$ ,  $p \in [1, \infty]$ , let  $(\mathbb{R}^n, d_p)$  be the Minkowski metric space of order  $p$ . The *unit sphere* is defined as:

$$\{x \in \mathbb{R}^n \mid d_p(x, 0) = 1\}$$

- Unit sphere with  $n = 2$ ,  $p = 2$ :




# Unit spheres in Minkowski metric spaces

- For  $n \in \mathbb{N}$ ,  $p \in [1, \infty]$ , let  $(\mathbb{R}^n, d_p)$  be the Minkowski metric space of order  $p$ . The *unit sphere* is defined as:

$$\{x \in \mathbb{R}^n \mid d_p(x, 0) = 1\}$$

- Unit sphere with  $n = 2$ ,  $p = \infty$ :



# Modified definitions

There are some special types of spaces similar to metric spaces:

## Definition

A *pseudometric space* has  $d$  satisfy symmetry and the triangle inequality, but allows for  $x \neq y$ , with  $d(x, y) = 0$ .

# Modified definitions

There are some special types of spaces similar to metric spaces:

## Definition

An *extended metric space* allows  $d$  to take on the value  $d(x, y) = \infty$ . All other properties of a metric remain, under the conventions  $\infty = \infty$ ,  $\infty \leq \infty$ ,  $\infty + \infty = \infty$ , and for  $c \in \mathbb{R}$ ,  $c < \infty$  and  $c + \infty = \infty$ .

# Modified definitions

There are some special types of spaces similar to metric spaces:

## Definition

An *ultrametric space* satisfies conditions (i) and (ii) but replaces the triangle inequality with the *strong triangle inequality*:

$$\forall x, y, z \in X, \quad d(x, z) \leq \max(d(x, y), d(y, z))$$



## Part 2:

Part 2 video: Metric Spaces in Practice!