

MATH 287 HOMEWORK 2

ANDREW MOORE

Date: September 10, 2021.

Exercise 1. Proposition 2.7(iv). Let $m, n, p, q \in \mathbf{Z}$. If $m < n$ and $p < 0$ then

$np < mp$.

Proof. -enter your proof here-

□

Exercise 2. Proposition 2.12(iii). For all $m, n, p \in \mathbf{Z}$, if $p < 0$ and $mp < np$ then $n < m$.

Proof. We are told we can assume the following: $0 - p \in \mathbf{N}$, and $np - mp \in \mathbf{N}$.

We must show that $n < m$, i.e. $m - n \in \mathbf{N}$. Rearranging $np - mp \in \mathbf{N}$, we have:

$$\begin{aligned} np - mp &= p(n - m) \\ &= p(n(-1)(-1) + m(-1)) \\ &= p(-1)(n(-1) + m) \\ &= -p(-n + m) \\ &= -p(m - n) \in \mathbf{N}. \end{aligned}$$

We can reexpress $0 - p \in \mathbf{N}$ as $-p \in \mathbf{N}$ (which we know from Axiom 2.1.4).

From the series of equations above, we've seen that $np - mp \in \mathbf{N}$ can be expressed as $-p(m - n) \in \mathbf{N}$. Axiom 2.1.2 tells us that the product of any two numbers in \mathbf{N} is also in \mathbf{N} . Therefore, $(m - n)$ must be in \mathbf{N} , i.e. $n < m$.

□

Exercise 3. Proposition 2.26. In this problem, the textbook gives a proof.

Your homework is to rewrite the proof in more detail.

Imagine a student in the class is confused by the proof. Rewrite the proof in a way that would make sense and be clear for a confused student.

Exercise 4. Project 2.28. Determine for which natural numbers $k^2 - 3k \geq 4$ and prove your answer.

Answer.

Claim 4.1. $k^2 - 3k \geq 4$ for $k \geq 4$.

Proof. We are given the inequality, $k^2 - 3k \geq 4$, and are asked to justify which values of k the inequality is true. We will prove the claim $k^2 - 3k \geq 4$ for $k \geq 4$ using induction. First, note that $k \leq 3$ does not satisfy the inequality.

For $k = 3$:

$$k^2 - 3k \geq 4 \Rightarrow (3)^2 - 3(3) \geq 4 \Rightarrow 9 - 9 \geq 4 \Rightarrow 0 \geq 4$$

The statement $0 \geq 4$ is *false*, because $0 \neq 4$ and $0 - 4 \notin \mathbf{N}$. It is trivial to demonstrate the same result for $k = 1$ and $k = 2$, and this is left to the reader. Assuming that k can't be ≤ 3 , let us redefine k to be $k = 3 + j$, such that $j \in \mathbf{N}$ and $j \geq 1$. We will now prove the base case.

For $k = 3 + 1 = 4$ ($j = 1$):

$$k^2 - 3k \geq 4 \Rightarrow (4)^2 - 3(4) \geq 4 \Rightarrow 16 - 12 \geq 4 \Rightarrow 4 \geq 4$$

This last statement is true ($4 = 4$). We will now show that the results apply for all j , by proving this is true for $k = 4 + (j + 1)$.

For $k = 4 + (j + 1) = 5 + j$:

$$(k + 1)^2 - 3(k + 1) \geq 4$$

$$(k^2 + 2k + 1) - 3k - 3 \geq 4$$

$$k^2 - k - 2 \geq 4$$

$$(5 + j)^2 - (5 + j) - 2 \geq 4$$

$$25 + 10j + j^2 - j - 7 \geq 4$$

$$j^2 + 9j + 18 \geq 4$$

The final statement $j^2 + 9j + 18 - 4 \in \mathbf{N}$ is true because there is no value of j in \mathbf{N} that could cause the value of the statement to be ≤ 0 . \square

\diamond