

# MATH 287 HOMEWORK 3

ANDREW MOORE

---

*Date:* September 17, 2021.

*Exercise 1.* Project 3.1. Express each of the following statements using quantifiers.

(iii) Every integer is the product of two integers.

(iv) The equation  $x^2 - 2y^2 = 3$  has an integer solution.

*Answer.*

(iii)  $\forall m \in \mathbf{Z}, \exists a \in \mathbf{Z} \text{ and } \exists b \in \mathbf{Z}, \text{ such that } m = a \cdot b.$

(iv) There exist integers  $x$  and  $y$  in  $\mathbf{Z}$ , such that the equation  $x^2 - 2y^2 = 3$  is true.

◇

*Exercise 2.* Project 3.2. In each of the following cases explain what is meant by the statement and decide whether it is true or false.

(iii) For each  $x \in \mathbf{Z}$  there exists  $y \in \mathbf{Z}$  such that  $xy = x$ .

(iv) There exists  $y \in \mathbf{Z}$  such that for each  $x \in \mathbf{Z}$ ,  $xy = x$ .

*Answer.*

(iii) This case states that for every integer  $m$  in the set of  $\mathbf{Z}$ , another integer  $y$  exists that satisfies the equation  $xy = x$ . I would say that it is true, but it is not specific enough. We know there exists only one  $y$  that satisfies the equation  $xy = x$ , which is  $y = 1$  (Axiom 1.3).

(iv) This case states that there is an integer  $y$ , and for each integer  $m$  in  $\mathbf{Z}$ ,  $y$  satisfies the equation  $xy = x$ . As above, I would say this statement is true, but should be made stronger by saying "there exists a unique integer  $y = 1$ , which satisfies the equation  $xy = x$ ". Alternatively, the (strong) statement could be expressed as  $\exists!y \in \mathbf{Z}: xy = x$ .

◇

*Exercise 3.* Project 3.7. Negate the following statements.

- (iv) The newspaper article was neither accurate nor entertaining. (Phrased differently: "The newspaper article was not accurate, and not entertaining.")
- (v) If  $\gcd(m, n)$  is odd, then  $m$  or  $n$  is odd.
- (vi)  $H/N$  is a normal subgroup of  $G/N$  if and only if  $H$  is a normal subgroup of  $G$ .
- (vii) For each  $\varepsilon > 0$  there exists  $N \in \mathbf{N}$  such that for all  $n \geq N$ ,  $|a_n - L| < \varepsilon$ .

*Answer.*

- (iv) The newspaper article was either accurate, or entertaining.
- (v) If  $\gcd(m, n)$  is even, then  $m$  and  $n$  are even.
- (vi)  $H/N$  is a normal subgroup of  $G/N$  and  $H$  is not a normal subgroup of  $G$ , or  $H$  is a normal subgroup of  $G$  and  $H/N$  is not a normal subgroup of  $G/N$ .

(vii) There exists some  $\varepsilon > 0$ , for all  $N$  in  $\mathbf{N}$ , such that an  $n \geq N$  exists

where  $|a_n - L|$  is greater than or equal to  $\varepsilon$ .

◇

*Exercise 4.* For all  $k \in \mathbf{N}$ ,  $5^k + 3$  is divisible by 4.

- (1) Write what this statement says for  $k = 1$ . Is it true or false? Explain.
- (2) Write what this statement says for  $k = 2$ . Is it true or false? Explain.
- (3) Write what this statement says for  $k = 3$ . Is it true or false? Explain.

Now, prove the statement for all  $k \in \mathbf{N}$ , using induction.

- (1) For  $k = 1$ , the statement says that  $5^1 + 3$  is divisible by 4. It's true  
because  $5 + 3 = 8$ , and  $8 = 4 \cdot 2$ .
- (2) For  $k = 2$ , the statement says that  $5^2 + 3$  is divisible by 4. It's true  
because  $25 + 3 = 28$ , and  $28 = 4 \cdot 7$ .
- (3) For  $k = 3$  the statement says that  $5^3 + 3$  is divisible by 4. It's true  
because  $125 + 3 = 128$ , and  $128 = 4 \cdot 32$ .

*Proof.* We are asked to prove that for all  $k \in \mathbf{N}$ ,  $5^k + 3$  is divisible by 4. We will show this is true using induction.

Base case ( $k = 1$ ):

$$5^1 + 3 = 5 + 3$$

$$= 8$$

$$= 4 \cdot 2.$$

We have shown that there is at least one  $k \in \mathbf{N}$  that proves the statement is true. Assuming that  $k \in \mathbf{N}$  is true, we will now show that  $(k + 1) \in \mathbf{N}$ :

$$5^{k+1} + 3 = (5^k \cdot 5^1) + 3$$

$$= 5^k \cdot 5 + 3$$

$$= \dots$$

□