

Global HIV Trends and Treatment Effectiveness Analysis

Data storytelling presentation based on the Power BI dashboard

Maxine Ndudzo-Sibiya

GLOBAL HIV TRENDS AND TREATMENT EFFECTIVENESS ANALYSIS

692M

People Living with HIV

13M






New Infections

26M

HIV Deaths

44.82

Average ART Coverage %



1. The global HIV picture is improving, but the burden remains large

692M

People living with HIV

13M

New infections


26M

HIV deaths


44.82%

Average ART coverage

Storyline: the dashboard shows progress over time, but the scale of people living with HIV means prevention, testing, and treatment must remain a global priority.




2. New HIV infections are trending downward



The line chart shows new infections falling from the early 2000s to 2024. This suggests that prevention, awareness, and testing programmes are having an impact.

However, the trend has not reached zero. Continued prevention campaigns are still needed, especially in regions where new infections remain high.



3. HIV deaths have reduced, showing the value of treatment access



Deaths decline over time alongside new infections. This may reflect improved access to ART, stronger HIV care, and better follow-up for people already diagnosed.


The priority is to protect these gains by expanding treatment coverage and reducing late diagnosis.

4. Africa carries the highest HIV burden



Both continent charts tell the same story: Africa has the largest number of people living with HIV and the highest HIV-related deaths. This makes it the most important region for targeted action.

5. ART coverage differs by country



The top ART coverage countries are mainly high-income European countries. This shows that treatment access is not equal across countries.

Countries with lower ART coverage should be prioritised because weaker treatment access can increase deaths and reduce quality of life.

6. Actionable recommendations

1

Prioritise high-burden regions

Increase funding for HIV prevention, testing, and ART programmes, especially in Africa.

2

Improve ART access

Target countries with lower treatment coverage to reduce HIV-related deaths.

3

Strengthen prevention campaigns

Focus education, testing, and awareness programmes where new infections remain high.

4

Use country-level monitoring

Track weak areas and target support based on data rather than broad assumptions.

Conclusion: Global HIV outcomes are improving, but resources should be directed to the regions and countries where the dashboard shows the greatest burden and weakest treatment access.