

GLOBAL HIV TRENDS AND TREATMENT EFFECTIVENESS ANALYSIS

692M

People Living with HIV

13M

New Infections


26M

HIV Deaths


44.82

Average ART Coverage %


New HIV Infections Over time




New Infections vs HIV Deaths Over Time




Top 10 Countries by Average ART Coverage



People Living with HIV by Continent



HIV Deaths by Continent



Interactive Filters

Year

2000

2024

Continent

Africa

Country

All

Use the filters above to explore the dashboard by year, continent and country

KEY INSIGHTS

1. Africa carries the highest HIV burden, with the largest number of people living with HIV and the highest HIV-related deaths.
2. New HIV infections show a downward trend over time, suggesting progress in prevention and awareness efforts.
3. HIV deaths have also reduced over time, which may indicate improved access to treatment and care.
4. ART coverage varies by country, showing that some countries have stronger treatment access than others.

KEY RECOMMENDATIONS

1. Prioritize high-burden regions, especially Africa, with increased funding for HIV prevention, testing, and treatment programmes.
2. Improve ART access in countries with lower treatment coverage to reduce HIV-related deaths and improve survival.
3. Strengthen prevention campaigns in countries or regions where new infections remain high.
4. Use country-level monitoring to identify areas where treatment access is weak and target support more effectively.