



Neurocognitive profiles of people with borderline personality disorder

Arthur D.P. Mak and Linda C.W. Lam

Purpose of review

This review summarizes recent neurocognitive research to better delineate the nosology, prognostication and cause underlying borderline personality disorder (BPD).

Recent findings

BPD had marked clinical heterogeneity with high comorbidity. Executive dysfunction in this disorder was linked to suicidality and treatment adherence, and may serve as an endophenotype. BPD was also characterized by cognitive distortions such as risky decision-making, deficient feedback processing, dichotomous thinking, jumping to conclusion, monocausal attribution and paranoid cognitive style. Social cognition deficits recently described in BPD include altered social inference and emotional empathy, hypermentalization, poorer facial emotional recognition and facial expressions. In electrophysiological studies, BPD was found to have predominantly right hemispheric deficit in high-order cortical inhibition. Reduced left orbitofrontal activity by visual evoked potential and magnetoencephalography correlated with depressive symptoms and functional deterioration. Brain structures implicated in BPD include the hippocampus, dorsolateral prefrontal cortex and anterior cingulate cortex. Abnormal anatomy and functioning of frontolimbic circuitry appear to correlate with cognitive deficits.

Summary

Frontolimbic structural and functional abnormalities underlie the broad array of cognitive abnormalities in BPD. Further research should espouse broader considerations of effects of comorbidity and clinical heterogeneity, and include community samples and, possibly, longitudinal designs.

Keywords

borderline personality disorder, emotions, frontal lobe, limbic system, neuropsychological tests

INTRODUCTION

Borderline personality disorder (BPD) is a complex and chronic mental disorder characterized by impulsivity, affective instability, cognitive distortions and unstable interpersonal relationships. It is the commonest of personality disorders, estimated to affect 0.5–5.9% of the general population [1,2]. People with BPD are characterized by complex comorbidities and high suicide rates [3]. Neurocognitive research may better delineate the nosology, prognostication and cause underlying BPD, but existing studies have yielded conflicting results.

This article reviews the recent findings on neuropsychological, electrophysiological and imaging studies related to BPD. Pubmed and Psycinfo search was done using keywords 'Borderline personality', 'Borderline personality disorder', 'cognition', 'executive function', 'attention', 'memory', 'visual perception', 'neuropsychological tests' and 'psychomotor performance'. Relevant articles were then selected.

CLINICAL PHENOMENON

Confirmatory factor analysis based on the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria established a three-factor model (Disturbed Relatedness, Affective Instability and Behavioral Dyscontrol) [4]. Another recent study of a mixed clinical sample found two latent classes (asymptomatic and symptomatic), as well as a single severity dimension [5]. These supported a hybrid categorical dimensional model underlying BPD. One single DSM-IV BPD criterion was found to be reliable in distinguishing individuals with more

Department of Psychiatry, The Chinese University of Hong Kong, Hong Kong SAR

Correspondence to Arthur D.P. Mak, Department of Psychiatry, The Chinese University of Hong Kong, Tai Po Hospital, Chuen On Road, Tai Po, New Territories, Hong Kong SAR. Tel: +852 26076027; e-mail: arthurdpmak@gmail.com

Curr Opin Psychiatry 2013, 26:90–96

DOI:10.1097/YCO.0b013e32835b57a9

KEY POINTS

- BPD was associated with executive dysfunction, social cognition deficits and cognitive distortions.
- Executive dysfunction could predict suicide, though could be amenable to treatment and predict better treatment adherence.
- Electrophysiological findings from BPD reflected deficient high-order inhibition of attentional processes.
- Anatomically and functionally, frontolimbic networks are implicated in the neuropsychological dysfunctions in BPD.

axis-I disorder, suicidality, hospitalizations and impairment [6]. Dimensional scoring effectively identified individuals with subthreshold borderline personality features [7]. These supported the potential of espousing dimensionality in diagnosis.

The upcoming DSM-5 classification proposes a prototype matching system that defines personality disorders with narrative descriptions. Expert consensus ratings on a broad array of personality traits showed that ratings of a prototypical case of BPD based on the DSM-5 underemphasized constructs of antagonism and disinhibition compared with DSM-IV BPD criteria, with more reliance placed on interpersonal dependency. On diagnostic ratings of psychiatric outpatients with personality disorders, analysis of mediator models using bootstrapping technique found that antagonism and disinhibition mediated the correlation between the DSM-IV BPD diagnosis and impairment (overall, social, marital, occupational, distress to others) [8]. This called for caution on the heuristic and clinical impact of this change of criteria.

COMORBIDITY

A total of 84.5% of BPD patients had one or more 12-month comorbid axis I disorders, including post-traumatic stress disorder (PTSD), mood and anxiety disorders [1]. In particular, 39.2% of BPD patients also suffered from PTSD. Impulsivity and emotional dysregulation are shared with attention deficit hyperactivity disorder (ADHD), which was found in 11% of a tertiary-care BPD sample, and predicted a higher prevalence of disruptive disorders [9]. Bipolar disorder and BPD have in common the features of affective instability, impulsivity and elevated suicide rates. BPD showed more prominent maladaptive self-schemas, more motor impulsivity and lack of planning, whereas bipolar patients have more deficits in sustained attention. Childhood

trauma was less common in bipolar disorder except for rapid-cycling bipolar disorder, which had similar rates of history of abuse to BPD [10].

NEUROPSYCHOLOGICAL FINDINGS: EXECUTIVE DYSFUNCTION

BPD has been associated with deficits in executive control, working memory and long-term memory consolidation [11], which are related to the core constructs of affective dysregulation and impulsivity.

Completed suicide occurs in 10% of BPD patients [3]. Both suicide attempters and BPD patients were associated with executive dysfunction, especially on trail-making and decision-making tests. It had been proposed that cognitive rigidity and aberrant decision-making processes may contribute to suicidal behavior [12]. In a more recent study, patients with BPD were found to have weaker Stroop interference control than healthy controls. Multivariate analysis found weak Stroop interference and Beck Depressive Inventory score to be the only significant predictors of suicide risk and lifetime suicide attempts [13], both having likelihood ratios of similar sizes.

In addition, better baseline executive control (trail-making test) and visual memory performance (Benton visual retention) predicted higher treatment adherence in a year-long treatment program consisting of antidepressant drug treatment and dialectic behavioral therapy for BPD [14].

Executive function deficits may improve with BPD treatment and could serve as treatment targets and efficacy indicators. Dialectical behavioral therapy-mindfulness module [15] was found to result in significantly improved scores in continuous performance test (CPT-II; commissions, hit reaction time, detectability scores) in BPD patients, and duration of mindfulness practice was associated with greater improvement in clinical symptomatology.

Executive dysfunction also showed promise as an endophenotype for BPD. First-degree relatives of BPD patients [16] had been found to have response inhibition deficits (Conner's CPT-II-elevated commission errors; atypically fast response times), with a moderately high rate of recurrence among siblings, as well as nonredundance with diagnostic status, supporting its use as an endophenotype.

NEUROPSYCHOLOGICAL FINDINGS: 'COLD' COGNITIVE DISTORTIONS: DECISION-MAKING AND FEEDBACK PROCESSING

Patients with BPD tended to make risky choices and were unable to improve their performance [17]. A reduced feedback-related negativity predicted

inability to learn from negative feedback. These decision-making deficits were correlated with BPD symptom severity and impulsivity, suggesting risky decision-making to be a perpetuating factor for BPD.

NEUROPSYCHOLOGICAL FINDINGS: 'COLD' COGNITIVE DISTORTIONS: COGNITIVE BIASES

Dichotomous thinking has commonly been identified in BPD. This, along with other cognitive biases more akin to schizophrenic thinking styles, including monocausal attribution, jumping to conclusion and paranoid cognitive style such as mistrust and sensory irritations, has recently been replicated in BPD [18²²]. These biases were found to be associated with BPD symptom severity. In contrast to the tendency of schizophrenic patients to externalize events [19], BPD patients showed a tendency to internalize both positive and negative events without self-serving bias.

NEUROPSYCHOLOGICAL FINDINGS: SOCIAL COGNITION: EMPATHY

Empathy, comprising cognitive and emotional components, may contribute to interpersonal dysfunction in BPD. Using the Multifaceted Empathy test, BPD patients [20²³] were found to have significant impairments in cognitive and emotional empathy. Altered social inference and emotional empathy could contribute to dysfunctional emotional and social responses in BPD.

NEUROPSYCHOLOGICAL FINDINGS: SOCIAL COGNITION: MENTALIZATION

Mentalization, defined as a form of imaginative mental activity for perception of interpretation of human behavior in terms of intentional mental states [21²⁴], has been linked to the social cognitive difficulties in BPD.

On a multi-round virtual trust game [22²⁵], BPD patients were better at adjusting their investment to the fairness of their virtual partner and assessing relevant emotional cues from unfair trustees, and at using objective fairness of their counterparts to guide responses. This superior 'theory of mind' may actually represent that a 'hypermentalizing' tendency, which refers to the deployment of unusual alternative strategies, was associated with BPD traits [21²⁶]. This may lead to socially inappropriate reactions in BPD.

BPD patients showed poorer social perspective coordination, that is, capacity to differentiate and integrate the perspectives of self with those of others

[23²⁷]. BPD patients had a biased perception of participation. They more easily felt excluded even when included [24²⁸], in turn having increase in other-focused negative emotions.

NEUROPSYCHOLOGICAL FINDINGS: SOCIAL COGNITION: FACIAL EMOTION RECOGNITION

Facial emotion recognition may contribute to social cognitive responses. BPD patients were found to exhibit more overall errors in recognizing facial expressions on facial morph tasks, in particular those with negative valence with fearful and surprised expressions [25]. Errors for expressions with positive valence were less consistently elevated [26²⁹]. A higher facial emotion recognition threshold in BPD was, in turn, associated with 'difficulty identifying feelings', a facet of alexithymia [27³⁰].

In another study with young BPD patients, this error-proneness in emotional facial recognition was not replicated [28³¹]. Instead, a later study found that [29³²], on a modified dot-probe task, BPD youth had faster response to congruent fear stimuli, and were slower in responding to incongruent than paired neutral trials. This indicated an attentional bias for fearful faces characterized by difficulty in disengaging attention from threatening stimuli.

NEUROPSYCHOLOGICAL FINDINGS: SOCIAL COGNITION: FACIAL EMOTIONAL EXPRESSION

In the above-mentioned study mimicking social participation situations, BPD patients reacted with fewer positive facial expressions and more mixed emotional expressions in response to social exclusion [24³³]. Ambiguous facial emotional expressions may account for relationship disturbances in BPD.

ELECTROPHYSIOLOGICAL STUDIES: ELECTROENCEPHALOGRAPHY

Electroencephalography (EEG) abnormalities have been found in BPD patients, including diffuse slow activity, random or semi-rhythmic theta and delta waves in the absence of focal or epileptiform features. Under a Bayesian statistical network [30³⁴], such EEG abnormalities were found to be interdependent, in BPD patients, with myriad other neurophysiological abnormalities, including sleep EEG (abnormal slow-wave sleep, rapid eye movement latency and duration), thyroid function tests, dexamethasone suppression test and neurological soft signs. This interrelated pathophysiological

mechanism underlying BPD may further be evaluated and broadened to help explore BPD nosology and cause.

ELECTROPHYSIOLOGICAL STUDIES: EVENT-RELATED POTENTIAL

The p3a event-related potential was studied in BPD patients to assess deficits in high-order inhibition in attention [31²²]. Right-handed BPD patients showed larger amplitude and delayed habituation of p3a in the right hemisphere, especially the frontomedial aspect, compatible with deficient high-order inhibition in attentional processes. Previous research had shown right hemisphere volume to be associated with maternal behavior in infancy and to be less genetically determined than the left hemisphere [32,33]. It is, thus, plausible that the right hemispheric predominance in deficit in BPD may reflect environmental or maturational influences.

ELECTROPHYSIOLOGICAL STUDIES: VISUAL EVOKED POTENTIAL AND MAGNETOENCEPHALOGRAPHY

Depressive symptomatology and global functional deterioration [34²²] were found to be significantly correlated with reduced left orbitofrontal activity in BPD patients, as measured with minimum normal estimate of steady-state visual evoked fields on magnetoencephalography.

ELECTROPHYSIOLOGICAL STUDIES: TRANSCRANIAL MAGNETIC STIMULATION

Cortical silent period (CSP) indicates cortical activation of inhibitory gamma-aminobutyric acid (GABA) neurotransmitter circuits. A recent transcranial magnetic stimulation study found BPD patients [35²²] to have reduced CSP. Alexithymia in BPD was not significantly associated with CSP after controlling for psychopathologic factors, but left CSP predicted 'difficulty identifying feelings'. Although this study highlighted the significance of GABA-mediated cortical inhibition and alexithymia in BPD patients, the independence of alexithymia from BPD as phenomenological constructs needs further assessment.

ANATOMICAL STUDIES: DORSOLATERAL PREFRONTAL CORTEX, HIPPOCAMPUS AND TRAUMA

Structural MRI found reduced hippocampal size and smaller dorsolateral prefrontal cortex (DLPFC) gray matter volumes in BPD [36,37²²]. Abuse in

childhood was associated with bilaterally reduced hippocampal volume, whereas DLPFC gray matter volumes inversely varied with impulsiveness [37²²].

In a recent meta-analysis, BPD patients had bilateral reductions in hippocampal volumes. In the four studies including PTSD comorbidity, the association was most prominent in comorbid BPD–PTSD, but rather mixed in BPD patients without PTSD [38²²]. PTSD may contribute to hippocampal reduction, while traumatic experience could also mediate volumetric reductions in the hippocampus in BPD.

ANATOMICAL STUDIES: ANTERIOR CINGULATE CORTEX

BPD adults were found to have smaller anterior cingulate cortex (ACC) [39–41]. Results from adolescents have been conflicting [42,43], probably attributed to cohort effects accentuated by proximity to onset time, heterogeneity in comorbidity and sample restrictions. A recent morphometric study on adolescents with BPD and major depression [44²²] showed significantly reduced BA 24 gray matter volume, which predicted increased symptom severity and suicide attempts.

ANATOMICAL STUDIES: AMYGDALA

A previous meta-analysis reporting significant reductions in hippocampal and amygdala volumes in BPD [45] was limited by omitting studies that did not show amygdala volume reductions [46]. One recent meta-analysis [47²²], including only one [48] of the negative studies omitted in the previous meta-analysis, found modest volume reductions of amygdala and hippocampus bilaterally, independent of illness state or comorbidity.

ANATOMICAL STUDIES: NEUROANATOMY AND SUICIDE

Apart from the association between ACC volume and suicidality [44²²], a recent structural MRI brain study [49²²] found that, of BPD patients, high lethality attempters had smaller right middle superior temporal gyrus, middle-inferior orbitofrontal gyrus, parahippocampal gyrus and insular cortex, and left fusiform gyrus and lingual gyrus, compared with low lethality attempters. This implies broad areas of deficits associated with executive dysfunction, facial emotion perception and memory processes. These observations are consistent with a model of diffuse neurobiological vulnerability to suicidal behavior with accentuated reactions at times of increased stress.

FUNCTIONAL STUDIES: POSITRON EMISSION TOMOGRAPHY

In resting state, euthymic BPD patients, matched to controls for impulsivity, were reported to have hypometabolism in frontal lobe and hypermetabolism in motor cortical regions, medial and anterior cingulus, occipital and temporal lobes, left superior parietal gyrus and right superior frontal gyrus [50^{***}].

FUNCTIONAL STUDIES: FUNCTIONAL MRI

BPD patients had enhanced activation of left amygdala and right insula after viewing aversive stimuli [51^{***}]. When being prompted to decrease the initial emotional reaction, BPD patients were less able to voluntarily decrease aversive emotions by cognitive reappraisal, correlated with reduced activation of left orbitofrontal cortex and increased activation of both insulae. Opiate-dependent BPD patients were found to have reduced activation to negative stimuli in amygdala and anterior cingulate when responding to emotional cues [52^{***}], but the effect could not be teased out from the fact that all the opiate-dependent patients were on suboxone while functional MRI (fMRI) was conducted, and a substantial proportion had active substance use.

During cognitive empathy, BPD patients had significantly reduced brain responses in the left superior temporal sulcus and gyrus. The level of reduction was associated with levels of intrusive symptomatology. During emotional empathy, BPD patients showed greater brain activity in the right middle insular cortex, in association with skin conductance response, which denoted hyperarousal [20^{***}].

CONNECTIVITY: WHITE MATTER CONNECTIVITY

Previous Diffusion Tensor Imaging studies had shown abnormal prefrontal functioning in BPD patients, but most included samples of BPD patients with comorbidities such as ADHD [53] or particular features such as severe self-harm or dissociative symptoms [54]. A recent study on medication-free noncomorbid BPD found significant decrease of fractional anisotropy in the genu and rostral areas of the corpus callosum as well as prefrontal white matter fasciculi bilaterally [55^{***}]. This showed similarity to people with major depressive disorder who had decreased fractional anisotropy in dorsolateral prefrontal white matter [56], and bipolar disorder patients with a lower fractional anisotropy in the genu [57,58].

CONNECTIVITY: FUNCTIONAL CONNECTIVITY

A recent resting-state fMRI study found functional connectivity differences in the default mode network (DMN) and a right frontoparietal network in BPD, with regions of abnormal connectivity within DMN [cuneus, insula and frontoparietal cortex (FPC)] related to BPD symptoms [59^{***}]. These findings, though limited by nonexclusion of axis-I comorbidity and psychotropic medication use, were the first to suggest the role of abnormal functional connectivity of prefrontal and insula resting-state networks in core BPD symptom clusters of impulsivity and dissociation. The study confirmed the previous task-based abnormalities in left FPC physiology, but did not find abnormalities in executive function and language-related regions in the resting state, which may show up better in task-based studies.

Another fMRI study [60^{***}] found that limbic and paralimbic regions had enhanced negative coupling with prefrontal regions when BPD patients experienced pain while receiving an emotion arousing stimulus, whereas positive connectivity was found in BPD between limbic and basal ganglia, as well as the precuneus and posterior cingulate, when seeing neutral pictures combined with painful heat sensation. This illustrated brain correlates in alterations of emotional regulation processes in BPD patients when experiencing pain.

CONCLUSION

In summary, increased evidence found frontolimbic structural and functional abnormalities to underlie the broad array of cognitive abnormalities in BPD.

The recent literature confirmed the conceptual coherence of BPD but accentuated clinical heterogeneity and overlap with other diagnostic entities. The effects of these issues are felt in all modalities in neurocognitive research in BPD. Only a few studies were able to identify homogeneous samples, while it may be argued that BPD may in itself be so complex and heterogeneous that, rather than distilling study samples into 'pure' form, 'natural' comorbid and heterogeneous samples may be more representative. Also, most studies had low sample size. This hampers the power of the studies given the infinite, overlapping and nonexclusive nature of borderline psychopathology. In addition, community-based data on BPD are sparse, which raises the question of whether the high comorbidity of BPD is inflated by referral filter bias. Nonetheless, the question of comorbidity in BPD may be even better understood in the perspectives of multimorbidity in terms of affective, behavioral and cognitive disorders across

all overlapping conditions, which would be an audacious undertaking. Longitudinal data are similarly lacking, which could be a result of the difficulty in retaining and recruiting BPD patients in research. Future research on neurocognitive domains of BPD should, therefore, be designed with explicit inclusion and comparison of various subgroups of BPD with different symptomatology and comorbidities.

Acknowledgements

None.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (pp. 129–131).

1. Grant BF, Chou SP, Goldstein RB, *et al.* Prevalence, correlates, disability, and comorbidity of DSM-IV borderline personality disorder: results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry* 2008; 69:533–545.
 2. Lenzenweger MF, Lane MC, Loranger AW, *et al.* DSM-IV personality disorders in the National Comorbidity Survey Replication. *Biol Psychiatry* 2007; 62:553–564.
 3. Paris J. Chronic suicidality among patients with borderline personality disorder. *Psychiatr Serv* 2002; 53:738–742.
 4. Calvo N, Andion O, Gancedo B, *et al.* Borderline personality disorder (BPD) ■ diagnosis with the self-report Personality Diagnostic Questionnaire-4+ (PDQ-4+): confirmation of the 3-factor structure. *Actas Esp Psiquiatr* 2012; 40:57–62.
- This study was the first to use confirmatory factor analysis based on a self-report measure on borderline psychopathology.
5. Hallquist MN, Pilkonis PA. Refining the phenotype of borderline personality disorder: diagnostic criteria and beyond. *Personal Disord* 2012; 3:228–246.
 6. Zimmerman M, Chelminski I, Young D, *et al.* Does the presence of one feature of borderline personality disorder have clinical significance? Implications for dimensional ratings of personality disorders. *J Clin Psychiatry* 2012; 73:8–12.
 7. Zimmerman M, Chelminski I, Young D, *et al.* Is dimensional scoring of borderline personality disorder important only for subthreshold levels of severity? *J Pers Disord* 2012. [Epub ahead of print]
 8. Samuel DB, Miller JD, Widiger TA, *et al.* Conceptual changes to the definition of borderline personality disorder proposed for DSM-5. *J Abnorm Psychol* 2012; 121:467–476.
 9. Speranza M, Revah-Levy A, Cortese S, *et al.* ADHD in adolescents with borderline personality disorder. *BMC Psychiatry* 2011; 11:158.
 10. Coulston CM, Tanious M, Mulder RT, *et al.* Bordering on bipolar: the overlap ■ between borderline personality and bipolarity. *Aust N Z J Psychiatry* 2012; 46:506–521.
- This is a comprehensive and objectively presented review encompassing the broad similarities and contrasts between BPD and bipolar disorder in their clinical, anatomical and functional aspects.
11. Ruocco AC. The neuropsychology of borderline personality disorder: a meta-analysis and review. *Psychiatry Res* 2005; 137:191–202.
 12. LeGris J, van Reekum R. The neuropsychological correlates of borderline personality disorder and suicidal behaviour. *Can J Psychiatry* 2006; 51:131–142.
 13. Legris J, Links PS, van Reekum R, *et al.* Executive function and suicidal risk in ■ women with borderline personality disorder. *Psychiatry Res* 2012; 196:101–108.
- This is a preliminary study comparing BPD patients with healthy controls on executive functions, wherein weak Stroop interference control was associated with suicide risk and lifetime suicide attempts, with nearly equal contribution to that of depression. Executive function and depression together explained 34% of the variance in total suicide risk.
14. Fertuck EA, Keilp J, Song I, *et al.* Higher executive control and visual memory ■ performance predict treatment completion in borderline personality disorder. *Psychother Psychosom* 2012; 81:38–43.
- Derived from data from a randomized controlled trial of a treatment program with antidepressant drug treatment and dialectic behavioral therapy, strengths include medication-free status of patients at baseline and comprehensive neuropsychological assessment.
15. Soler J, Valdeperez A, Feliu-Soler A, *et al.* Effects of the dialectical behavioral ■ therapy-mindfulness module on attention in patients with borderline personality disorder. *Behav Res Ther* 2012; 50:150–157.
- The first study to show the attentional processes underlying mindfulness skills.
16. Ruocco AC, Laporte L, Russell J, *et al.* Response inhibition deficits in ■ unaffected first-degree relatives of patients with borderline personality disorder. *Neuropsychology* 2012; 26:473–482.
- Response inhibition deficit, a known heritable attribute, showed promise as an endophenotype of BPD.
17. Schuermann B, Kathmann N, Stiglmayr C, *et al.* Impaired decision making and ■ feedback evaluation in borderline personality disorder. *Psychol Med* 2011; 41:1917–1927.
- Astounding clinical significance from this finding about feedback evaluation deficit, which characterizes much of the social and decisional difficulties in BPD. Sample size limitations hamper power.
18. Moritz S, Schilling L, Wingenfeld K, *et al.* Psychotic-like cognitive biases in ■ borderline personality disorder. *J Behav Ther Exp Psychiatry* 2011; 42:349–354.
- The significant findings from this study should be replicated in larger samples including both sexes.
19. Lincoln TM, Mehl S, Exner C, *et al.* Attributional style and persecutory ■ delusions. Evidence for an event independent and state specific external-personal attribution bias for social situations. *Cogn Ther Res* 2010; 34:297–302.
 20. Dziobek I, Preissler S, Grodzanovic Z, *et al.* Neuronal correlates of altered ■ empathy and social cognition in borderline personality disorder. *Neuroimage* 2011; 57:539–548.
- The superior temporal sulcus was implicated in social cognitive processes, which may mediate dysfunctional emotional responses.
21. Sharp C, Pane H, Ha C, *et al.* Theory of mind and emotion regulation ■ difficulties in adolescents with borderline traits. *J Am Acad Child Adolesc Psychiatry* 2011; 50:563–573; e1.
- This study well illustrates the notion that mentalizing may influence borderline disorder through the mediating role of emotion dysregulation.
22. Franzen N, Hagenhoff M, Baer N, *et al.* Superior 'theory of mind' in borderline ■ personality disorder: an analysis of interaction behavior in a virtual trust game. *Psychiatry Res* 2011; 187:224–233.
- What appeared to be 'superior' theory of mind may in fact represent unusual and inappropriate mentalization.
23. Jennings TC, Hulbert CA, Jackson HJ, *et al.* Social perspective coordination in ■ youth with borderline personality pathology. *J Pers Disord* 2012; 26:126–140.
- This study described the social cognitive impairment in BPD, but inclusion of subthreshold condition may raise concern of increased heterogeneity.
24. Staebler K, Renneberg B, Stopsack M, *et al.* Facial emotional expression in ■ reaction to social exclusion in borderline personality disorder. *Psychol Med* 2011; 41:1929–1938.
- This study did not include a control group, and only women were recruited.
25. Domes G, Czeschnek D, Weidler F, *et al.* Recognition of facial affect in ■ borderline personality disorder. *J Pers Disord* 2008; 22:135–147.
 26. Unoka Z, Fogd D, Fuzy M, *et al.* Misreading the facial signs: specific ■ impairments and error patterns in recognition of facial emotions with negative valence in borderline personality disorder. *Psychiatry Res* 2011; 189:419–425.
- Lack of control group as comparison and inclusion of medicated participants limited the findings of this study, which should be replicated.
27. Domes G, Grabe HJ, Czeschnek D, *et al.* Alexithymic traits and facial emotion ■ recognition in borderline personality disorder. *Psychother Psychosom* 2011; 80:383–385.
- Small sample size, exclusion of male participants and lack of a control group were the main limitations, yet the results indicated the role of alexithymic traits in social cognitive functions.
28. Jovev M, Chanan A, Green M, *et al.* Emotional sensitivity in youth with ■ borderline personality pathology. *Psychiatry Res* 2011; 187:234–240.
- Sample size restrictions may have contributed to the negative findings in this study.
29. Jovev M, Green M, Chanan A, *et al.* Attentional processes and responding to ■ affective faces in youth with borderline personality features. *Psychiatry Res* 2012. [Epub ahead of print]
- This study focused on and located deficient attentional processes in responding to facial affect in BPD.
30. De la Fuente JM, Bengoetxea E, Navarro F, *et al.* Interconnection between ■ biological abnormalities in borderline personality disorder: use of the Bayesian networks model. *Psychiatry Res* 2011; 186:315–319.
- The use of a Bayesian network statistical method to explore the complex interdependent dynamics in causation of BPD should be considered for other disorders too.

31. Meares R, Schore A, Melkonian D. Is borderline personality a particularly right hemispheric disorder? A study of P3a using single trial analysis. *Aust N Z J Psychiatry* 2011; 45:131–139.
Right hemispheric deficits could reflect environmental effects and should be replicated in younger cohorts.
32. Lyons DM, Afarian H, Schatzberg AF, *et al.* Experience-dependent asymmetric variation in primate prefrontal morphology. *Behav Brain Res* 2002; 136:51–59.
33. Chanen AM, Velakoulis D, Carison K, *et al.* Orbitofrontal, amygdala and hippocampal volumes in teenagers with first-presentation borderline personality disorder. *Psychiatry Res* 2008; 163:116–125.
34. Diaz-Marsa M, Carrasco JL, Lopez-Ibor M, *et al.* Orbitofrontal dysfunction related to depressive symptomatology in subjects with borderline personality disorder. *J Affect Disord* 2011; 134:410–415.
Small sample size and restriction to the study of cortex by the method chosen were the main limitations. Control group was also lacking.
35. Lang S, Stopsack M, Kotchoubey B, *et al.* Cortical inhibition in alexithymic patients with borderline personality disorder. *Biol Psychol* 2011; 88:227–232.
Alexithymia was related to cortical inhibition in BPD. Larger samples should be used to explore the interaction between alexithymic traits and borderline disorder.
36. Zetsche T, Preuss UW, Frodl T, *et al.* Hippocampal volume reduction and history of aggressive behaviour in patients with borderline personality disorder. *Psychiatry Res* 2007; 154:157–170.
37. Sala M, Caverzasi E, Lazzaretti M, *et al.* Dorsolateral prefrontal cortex and hippocampus sustain impulsivity and aggressiveness in borderline personality disorder. *J Affect Disord* 2011; 131:417–421.
This study may benefit from larger sample size, but highlights the disparate role of the hippocampus and DLPFC.
38. Rodrigues E, Wenzel A, Ribeiro MP, *et al.* Hippocampal volume in borderline personality disorder with and without comorbid posttraumatic stress disorder: a meta-analysis. *Eur Psychiatry* 2011; 26:452–456.
The effect of PTSD on brain volume in BPD patients suggested a mediating effect of either trauma or the PTSD syndrome.
39. Hazlett EA, New AS, Newmark R, *et al.* Reduced anterior and posterior cingulate gray matter in borderline personality disorder. *Biol Psychiatry* 2005; 58:614–623.
40. Tebartz van Elst L, Hesslinger B, Thiel T, *et al.* Frontolimbic brain abnormalities in patients with borderline personality disorder: a volumetric magnetic resonance imaging study. *Biol Psychiatry* 2003; 54:163–171.
41. Minzenberg MJ, Fan J, New AS, *et al.* Frontolimbic structural changes in borderline personality disorder. *J Psychiatr Res* 2008; 42:727–733.
42. Whittle S, Chanen AM, Fornito A, *et al.* Anterior cingulate volume in adolescents with first-presentation borderline personality disorder. *Psychiatry Res* 2009; 172:155–160.
43. Brunner R, Henze R, Parzer P, *et al.* Reduced prefrontal and orbitofrontal gray matter in female adolescents with borderline personality disorder: is it disorder specific? *Neuroimage* 2010; 49:114–120.
44. Goodman M, Hazlett EA, Avedon JB, *et al.* Anterior cingulate volume reduction in adolescents with borderline personality disorder and co-morbid major depression. *J Psychiatr Res* 2011; 45:803–807.
There was no control group, but findings indicated anterior cingulate to be associated with suicidality.
45. Nunes PM, Wenzel A, Borges KT, *et al.* Volumes of the hippocampus and amygdala in patients with borderline personality disorder: a meta-analysis. *J Pers Disord* 2009; 23:333–345.
46. Leichsenring F, Leibing E, Kruse J, *et al.* Borderline personality disorder. *Lancet* 2011; 377:74–84.
47. Ruocco AC, Amirthavasagam S, Zakzanis KK. Amygdala and hippocampal volume reductions as candidate endophenotypes for borderline personality disorder: a meta-analysis of magnetic resonance imaging studies. *Psychiatry Res* 2012; 201:245–252.
BPD patients had significantly smaller amygdala in this MRI study. In contrast to a previous meta-analysis, more negative studies were included, but only one of those four excluded in the study by Nunes *et al.*
48. Schmahl C, Berne K, Krause A, *et al.* Hippocampus and amygdala volumes in patients with borderline personality disorder with or without posttraumatic stress disorder. *J Psychiatry Neurosci* 2009; 34:289–295.
49. Soloff PH, Pruitt P, Sharma M, *et al.* Structural brain abnormalities and suicidal behavior in borderline personality disorder. *J Psychiatr Res* 2012; 46:516–525.
A broad range of anatomical correlates were found predicting suicide risk, but medical sequelae of suicide attempts were not accounted for in the data.
50. Salavert J, Gasol M, Vieta E, *et al.* Fronto-limbic dysfunction in borderline personality disorder: a 18F-FDG positron emission tomography study. *J Affect Disord* 2011; 131:260–267.
The sample was small, with only eight BPD patients.
51. Schulze L, Domes G, Kruger A, *et al.* Neuronal correlates of cognitive reappraisal in borderline patients with affective instability. *Biol Psychiatry* 2011; 69:564–573.
A small sample size again limits power of this study.
52. Smoski MJ, Salsman N, Wang L, *et al.* Functional imaging of emotion reactivity in opiate-dependent borderline personality disorder. *Personal Disord* 2011; 2:230–241.
A proportion of patients had active substance use and all were using suboxone, which may account for the reduced amygdala activation.
53. Rusch N, Weber M, Il'yasov KA, *et al.* Inferior frontal white matter microstructure and patterns of psychopathology in women with borderline personality disorder and comorbid attention-deficit hyperactivity disorder. *Neuroimage* 2007; 35:738–747.
54. Grant JE, Correia S, Brennan-Krohn T, *et al.* Frontal white matter integrity in borderline personality disorder with self-injurious behavior. *J Neuropsychiatry Clin Neurosci* 2007; 19:383–390.
55. Carrasco JL, Tajima-Pozo K, Diaz-Marsa M, *et al.* Microstructural white matter damage at orbitofrontal areas in borderline personality disorder. *J Affect Disord* 2012; 139:149–153.
This study looked at the white matter connectivity in BPD. Sample size was limited.
56. Wu F, Tang Y, Xu K, *et al.* White matter abnormalities in medication-naïve subjects with a single short-duration episode of major depressive disorder. *Psychiatry Res* 2011; 191:80–83.
57. Adler CM, Holland SK, Schmithorst V, *et al.* Abnormal frontal white matter tracts in bipolar disorder: a diffusion tensor imaging study. *Bipolar Disord* 2004; 6:197–203.
58. Yurgelun-Todd DA, Silveri MM, Gruber SA, *et al.* White matter abnormalities observed in bipolar disorder: a diffusion tensor imaging study. *Bipolar Disord* 2007; 9:504–512.
59. Wolf RC, Sambataro F, Vasic N, *et al.* Aberrant connectivity of resting-state networks in borderline personality disorder. *J Psychiatry Neurosci* 2011; 36:402–411.
Resting-state MRI showed a 'DMN' pattern of deficit, with abnormal functional connectivity of prefrontal and insula and resting-state networks in core BPD symptom clusters of impulsivity and dissociation.
60. Niedtfeld I, Kirsch P, Schulze L, *et al.* Functional connectivity of pain-mediated affect regulation in borderline personality disorder. *PLoS One* 2012; 7:e33293.
This MRI study nicely illustrated the changes in functional connectivity to pain in people with BPD.