
Databases, search engines and repositories
Citation searching
Identification
Screening
Included
Records identified (n = 29,171):
PubMed (n = 1,258)
Europe PMC (n = 1,454)
OpenAlex (n = 11,027)
Semantic Scholar (n = 6,683)
arXiv (n = 3,074)
PsyArXiv (OSF) (n = 186)
Crossref (n = 306)
DataCite (n = 4,934)
CORE (n = 249)
Records removed before screening:
Duplicate records (n = 16,385)
Non−study records (n = 753)
No exact search−term match (n = 5,181)
Records identified from citation searching (not in the database set; n = 9,248)
Removed by the same automatic rules (before 2020, non−study, no exact term match; n = 9,023)
Records screened (n = 6,852)
Records excluded (n = 5,596)
Records screened (n = 225)
Records excluded (n = 155)
Reports sought for retrieval (n = 1,256)
Reports not retrieved (no open full text; n = 490)
Reports sought for retrieval (n = 70)
Reports not retrieved (no open full text; n = 53)
Reports assessed for eligibility (n = 766)
Reports excluded (n = 355):
No psychological or clinical construct (n = 133)
No human coding of the same units (n = 79)
No generative LLM (n = 41)
External criterion, not coders (n = 33)
Other (n = 28)
Theme−level agreement only (n = 21)
Humans rate LLM output (n = 8)
Design (review, protocol) (n = 8)
Duplicate report (n = 4)
Reports assessed for eligibility (n = 17)
Reports excluded (n = 10):
External criterion, not coders (n = 3)
Humans rate LLM output (n = 2)
No psychological or clinical construct (n = 2)
No human coding of the same units (n = 2)
No generative LLM (n = 1)
Reports included (n = 7)
Studies included in review (n = 377)
Reports of included studies (n = 416)
Excluded at data extraction: 2 studies
Studies with kappa−family estimates (n = 111)
Studies with ICC or correlations (n = 99)