Research Article

Induced Paths, Radius, and Extremal Neighbourhood Independence in Graphs without 4-Cycles

Xingchen Wang[1] | Xincao Xu*[2]

[1]Independent Researcher
[2]Shenzhen Institute for Advanced Study, University of Electronic Science and Technology of China

Correspondence: *Corresponding author: Xincao Xu, xc.xu@uestc.edu.cn. Xingchen Wang, skywalkerwang98@gmail.com.

Keywords: induced path, radius, neighbourhood independence, 4-cycle, WOWII Conjecture 133

Abstract

Let G be a connected graph on $n \geq 2$ vertices. Write $S(G)$ for the sum of the independence numbers of the open neighbourhoods, and write $\ell(G) = S(G)/n$. Every connected graph with no 4-cycle subgraph and no induced path on seven vertices satisfies $S \leq 3n$, with equality only for the Petersen graph. Combined with the classification of the connected graphs with no 4-cycle subgraph, radius 4 and induced-path order 7, this bound yields Conjecture 133 of *Written on the Wall II* for every connected graph with $\ell \leq 4$. The bound for graphs with at most one triangle is proved by hand; the bound for graphs with at least two triangles uses a finite search, cross-checked by a second implementation. The same arguments determine the maximum of ℓ over connected graphs with no 4-cycle subgraph and no induced path on six vertices, with a unique extremal graph for every order $n \geq 7$, and determine, for every $n \geq 10$, the maximum number of edges in a connected triangle-free graph with no 4-cycle subgraph and no induced path on seven vertices, together with all extremal graphs. The conjecture remains open for $\ell > 4$.

1 | Introduction

All graphs are finite and simple. For a graph G with $n \geq 2$ vertices let $p(G)$ be the largest number of vertices of an induced path and let $r(G)$ be the radius. Write

$$S(G) = \sum_{v \in V(G)} \alpha\big(G[N(v)]\big), \qquad \ell(G) = \frac{S(G)}{n},$$

where α is the independence number and $N(v)$ the open neighbourhood. Thus $\ell(G)$ is the average, over the vertices, of the independence number of the open neighbourhood. Write $c_{C_4}(G) = 1$ if G contains no 4-cycle as a subgraph and $c_{C_4}(G) = 0$ otherwise. A graph with no 4-cycle subgraph is called *strong C_4-free*. In the extremal literature the same condition is the usual meaning of C_4-free. In the induced-subgraph literature, including the theorems of Brandstädt and Hoàng, Karthick and Maffray, Chen and Xu, and Chen, Wu and Xu cited below, the same words mean that C_4 is forbidden only as an induced subgraph. The adjective *strong* records the subgraph condition. By contrast, P_k-free means that G has no induced path on k vertices.

A shortest path between two vertices at distance $\operatorname{diam}(G) \geq r(G)$ is induced, so $p(G) \geq r(G) + 1$. In a strong C_4-free graph any two vertices have at most one common neighbour, so each neighbourhood is a matching plus isolated vertices and $S(G) = 2e(G) - 3T(G)$.

Conjecture 1.1 (Graffiti/WOWII Conjecture 133)**.** *Every connected graph G on at least two vertices satisfies*

$$r(G) + \ell(G)^{c_{C_4}(G)} \le p(G).$$

The primary source is E. DeLaVina's list *Written on the Wall II*, produced by the conjecture-making program Graffiti.pc [6]. Entry 133, dated July 12, 2005, reads: if G is a simple connected graph, then path$(G) \ge$ rad(G) + [average of $l(v)$]$^{c\,C_4}$. On that list path(G) is the largest number of vertices of an induced path and rad(G) is the radius. Other entries write floors and ceilings in words, as FLOOR[$\cdots$] and CEIL[$\cdots$], so the brackets in entry 133 are grouping signs. The quantity $l(v)$ is the local independence of v, defined on the definitions page of the list as the independence number of the subgraph induced by the neighbours of v. The same page takes the exponent to be 1 when G is C_4-free as a subgraph, not necessarily as an induced subgraph, and 0 otherwise. That is the convention in Conjecture 1.1. A record of the conjecture in the Formal Conjectures repository is a secondary source [8].

The real-valued reading, with no floor, is the stronger of the two readings compatible with the brackets. Remark 5.12 proves the floor form for every connected graph with $\ell < 5$. Since $p(G) - r(G)$ is an integer, the real-valued inequality on a strong C_4-free graph is equivalent to

$$p(G) - r(G) \;\ge\; \lceil \ell(G) \rceil. \tag{1.1}$$

If G contains a 4-cycle, then $c_{C_4}(G) = 0$ and both readings reduce to $p(G) \ge r(G) + 1$.

The subgraph convention matters. Suppose the exponent were 1 precisely when G has no induced 4-cycle. The book $K_2 \vee \overline{K_5}$ has order 7, radius 1 and induced-path order 3. Every 4-cycle of this book has a chord, and $\ell = 15/7$, so $r + \ell = 22/7 > 3$. The same count shows that $K_2 \vee \overline{K_m}$ fails the real-valued inequality for every $m \ge 5$. Under the subgraph convention each of these graphs has exponent 0, and the inequality is the trivial bound $p \ge r + 1$. The problem therefore reduces to strong C_4-free graphs.

The statements of independent interest are the sharp extremal theorems, not the partial case of the conjecture. Linear upper bounds on the number of edges, in classes with no long induced path and no $K_{s,s}$ subgraph, are classical. Galvin, Rival and Sands proved that a long path in a $K_{t,t}$-free graph forces a long induced path [9]. Kühn and Osthus proved that large average degree forces a $K_{s,s}$ subgraph or an induced subdivision of any fixed graph [12]. Loh, Tait, Timmons and Zhou introduced induced Turán numbers, in which one graph is forbidden as a subgraph and another only as an induced subgraph [13]. In that setting a linear bound with an identified extremal family is already a strong conclusion. The class here is smaller. It forbids C_4 as a subgraph, hence $K_{2,2}$, and it forbids a long induced path.

Theorem 5.10. Every connected strong C_4-free P_7-free graph G satisfies $S(G) \le 3n$, with equality if and only if G is the Petersen graph.

The triangle-free case is the following statement. For those graphs $S = 2e$.

Theorem 5.5. Let G be a connected triangle-free strong C_4-free P_7-free graph. Then $2e(G) \le 3|V(G)|$, with equality if and only if G is the Petersen graph.

The same class has an exact edge maximum. Write $f(n)$ for the maximum of $2e(G)$ over its members of order n.

Theorem 5.6. For every $n \ge 10$,

$$f(n) = 3n - \min\{n - 10,\ 5 + \mathbf{1}_{\{n \text{ even}\}}\}.$$

The extremal graphs are the graphs named in the statement of the theorem: the Petersen graph with $n - 10$ leaves on a single vertex or on the two ends of a single edge, together with the graphs F_2 and F_0' and two one-leaf extensions of F_2, on the ranges of n given there.

These three theorems are the headline. Conjecture 1.1 is the motivation for proving them and the application below. Two further structural statements are used in that application and have exact forms of their own.

Theorem 3.6. Let $M(n)$ be the maximum of ℓ over connected strong C_4-free P_6-free graphs of order $n \ge 7$. Then $M(7) = 16/7, M(8) = 5/2, M(9) = 8/3, M(10) = 3$ and $M(n) = 2 + 2/n$ for $n \ge 11$, and the extremal graph is unique for every such n. Every connected strong C_4-free P_6-free graph of order $n \ge 2$ satisfies $S \le 3n$, with equality only for the Petersen graph.

Theorem 4.2. A connected strong C_4-free graph of radius 4 and induced-path order 7 is C_8, or it is obtained from a 6-cycle by the extension in the statement of the theorem. If b is the number of vertices added on edges of that 6-cycle and m is the number of matching edges in the sets C_i, then $S = 2n - b - m$. In particular $S \le 2n$.

Theorem 5.11. Every connected graph on at least two vertices with $\ell \le 4$ satisfies Conjecture 1.1.

The proof of Theorem 5.10 splits on the number of triangles. Theorem 5.5 is the triangle-free case. Its input is the classification, due to Chen and Xu, of the connected (P_7, C_4, C_3)-free graphs with no clique cutset, the observation that a 2-connected member of the triangle-free class has no clique cutset, and a decomposition of a general member into such a core with trees attached at single vertices. Comparing the deficits $3|V(Q)| - 2e(Q)$ of the cores gives Theorem 5.6 as well. Theorem 5.7 is the case of at most one triangle: a minimal counterexample is 2-connected, and deleting a minimum-degree vertex of the triangle returns the argument to a triangle-free core. If there are at least two triangles, Lemma 5.8 lists the induced subgraphs formed by two triangles, and Lemma 5.9 produces a vertex of deletion loss at most 3 by a finite search. The search has no order cutoff and no isomorphism pruning. At each node it selects the vertex of deletion loss less than 4 with the fewest legal children, breaking ties by the smallest label, and it explores every legal child. The completeness argument is in the proof of Lemma 5.9. A node of a search is unrestricted when every vertex of the current graph has deletion loss at least 4. Each of the four search trees is finite and has no unrestricted node, so the branch that follows a counterexample ends at a leaf. At that leaf the chosen vertex has no legal new neighbour, while the counterexample supplies one. Appendix A prints the decision arrays. An independent C implementation (`lemma57b.c`) written for cross-verification is archived in `referee_checks/`.

Theorem 5.11 then assembles the radius cases. It uses Lemmas 3.1, 3.2, 3.3 and 3.5. Through Lemma 3.5 it also uses Lemmas 3.4 and 2.8, the Camby–Schaudt theorem and Seinsche's theorem. It uses the Erdős–Saks–Sós theorem; Theorem 3.6, in the case $r = 2$ and again when $r = 3$ and the induced-path order is at most 5; Corollary 4.4 through Theorem 4.2, together with the observation that $r = 4$ and $p \ge 8$ give $p \ge r + \ell$ whenever $\ell \le 4$; and Theorem 5.10. Theorem 5.10 uses Theorems 5.5 and 5.7, Lemmas 5.8 and 5.9, and the Chen–Xu classification. Lemma 5.9 enters only through Theorem 5.10, and only for graphs with at least two triangles. In the proof of Theorem 5.11 that use occurs only for $r = 3$, $3 < \ell \le 4$ and $p = 6$. If $3 < \ell \le 4$ and $p \le 5$, Theorem 3.6 gives $\ell \le 3$. If $p = 6$, Theorem 5.10 gives $\ell \le 3$.

Since $\lambda(v) \le d(v)$ for every vertex, $\ell \le 2e/n$. Theorem 5.11 covers every connected graph of average degree at most 4. For a strong C_4-free graph the counting bound of Reiman gives $\ell \le (1+\sqrt{4n-3})/2$, as recorded after the proof of Theorem 5.11. The right-hand side is at most 4 whenever $n \le 13$. Theorem 5.11 covers every connected graph on at most 13 vertices: a graph that contains a 4-cycle has exponent 0, and a strong C_4-free graph on at most 13 vertices has $\ell \le 4$. The same method gives $p \ge r + \ell$ whenever $\ell \le r - 1$, because the Erdős–Saks–Sós theorem supplies an induced path on at least $2r - 1$ vertices. It does not reach $\ell > 4$. The obstruction is the absence of an analogue of Theorem 5.10 for longer induced paths, for instance a bound $S \le 4n$ for connected strong C_4-free P_9-free graphs. Section 6 reduces any counterexample with $\ell > 4$ to a failure of GEN_K for some $K \ge 4$ or of $\mathrm{RAD}(r)$ for some $r \ge 5$, and it records the open conjecture on S together with four examples.

There is a parallel structure theory in which C_4 is forbidden only as an induced subgraph. Brandstädt and Hoàng describe (P_6, C_4)-free graphs by clique cutsets [1]. Karthick and Maffray describe the square-free members of that class [10]. Chen and Xu classify the connected (P_7, C_4, C_3)-free graphs with no clique cutset [5], and that classification is the core list for Theorems 5.5 and 5.6. Chen, Wu and Xu describe further (P_7, C_4)-free classes, with C_4 induced [4]. Those theorems do not by themselves yield Theorems 3.6, 5.5 or 5.6. The strong C_4-free class excludes every graph with a 4-cycle subgraph, including a nontrivial clique blow-up of the Petersen graph and a join of the Petersen graph with a nonempty clique, and it contains graphs with clique cutsets. The exact value of $M(n)$ uses the second-layer count on an induced 6-cycle, which needs the subgraph condition. Lemma 2.9 is the emptiness list on an induced 5-cycle, proved directly in Section 2 from P_6-freeness, induced-C_4-freeness and the absence of an induced C_6. The colouring bound of Karthick and Maffray is a different statement.

In [3], Chen studies the floor form $p \ge r + \lfloor \ell \rfloor^{c_{C_4}}$ of the same conjecture. That deposit is not a refereed paper, so every statement taken from it and used below is proved again in this article. Three points separate the two texts.

1. In [3], a graph with no 4-cycle subgraph satisfies $\ell = (2e - 3T)/n$. On an induced 6-cycle in a P_6-free graph of that class, an outside vertex has empty trace or meets the cycle in an antipodal pair. Under an auxiliary hypothesis RES (Definition 7.2 of that deposit), namely radius 2, diameter 3, $\ell > 3$ and (R3b), the second layer of such a P_6-free graph is restricted, and five admissible frames of orders 6 through 9, all with $\ell < 3$, are listed. The diameter-2 theorem in that deposit is likewise stated under $\ell > 3$.
2. The present article determines $M(n)$ for every order, with a unique extremal graph for each $n \ge 7$; classifies the radius-4 graphs of induced-path order 7; gives $S \le 3n$ for every connected strong C_4-free P_7-free graph, with equality

only for the Petersen graph; determines $f(n)$ and all extremal graphs for every $n \geq 10$; and proves the real-valued conjecture for $\ell \leq 4$, with no diameter hypothesis and no appeal to RES.

3. The identity $S = 2e - 3T$ is proved as Lemma 2.1, and the antipodal-trace constraint is proved, with the second-layer rule, in the proof of Lemma 3.4 and in Case 1 of Theorem 3.6. Those arguments stand on their own, so no step of this article depends on the unrefereed deposit [3].

Section 3 proves Theorem 3.6. Section 4 proves Theorem 4.2. Section 5 proves Theorems 5.5, 5.6, 5.7 and 5.10, and Theorem 5.11. Section 6 is the concluding section. It reduces the range $\ell > 4$ to two open statements and records the open conjecture on S together with four examples. The search of Lemma 5.9 is recorded in Appendix A.

2 | Preliminaries

For $v \in V(G)$ let $d(v)$ be its degree and $t(v)$ the number of triangles containing v. Write $\lambda(v) = \lambda_G(v) = \alpha(G[N(v)])$, $S(G) = \sum_v \lambda(v)$, $e(G)$ for the number of edges and $T(G)$ for the number of triangles.

Lemma 2.1. *Let G be strong C_4-free. Then:*

1. *two distinct vertices have at most one common neighbour;*
2. *for every v, $G[N(v)]$ is a matching together with isolated vertices, so $\lambda(v)$ equals the number of components of $G[N(v)]$ and $\lambda(v) = d(v) - t(v)$;*
3. *$S(G) = 2e(G) - 3T(G)$.*

Proof. (1) Two common neighbours w, w' of x, y give the 4-cycle $xwyw'$. (2) A vertex of $N(v)$ with two neighbours in $N(v)$ would have two common neighbours with v. So the components are single vertices and edges, each edge being a triangle at v. (3) Sum $d(v) - t(v)$ over v. □

The components of $G[N(v)]$ are the *units* of v. For a vertex set U, $G - U$ is the subgraph induced by $V \setminus U$, and $D_G(U) = S(G) - S(G - U)$ is the *deletion loss* of U. We write $L(v) = D_G(\{v\})$.

Lemma 2.2. *Let G be strong C_4-free and $\emptyset \neq U \subseteq V(G)$. Then*

$$D_G(U) = \sum_{u \in U} \lambda(u) + \sum_{x \notin U} \#\{\text{units of } x \text{ contained in } U\}.$$

In particular $L(v) = 2\lambda(v) - t(v) = 2d(v) - 3t(v)$.

Proof. For $x \notin U$, deleting U removes exactly the units of x that lie inside U; a unit that is an edge with one end in U survives as a vertex. For $U = \{v\}$: the $2t(v)$ neighbours of v in triangles with v keep their value, and each of the other $d(v) - 2t(v)$ neighbours loses one. The total is $\lambda(v) + d(v) - 2t(v) = 2\lambda(v) - t(v)$. □

Lemma 2.3. *Let v be a cut vertex of G. Split the components of $G - v$ into two nonempty groups, and let G_1, G_2 be the subgraphs induced by the two groups together with v. Then $S(G) = S(G_1) + S(G_2)$.*

Proof. $G[N(v)]$ has no edges between the groups, so $\lambda_G(v) = \lambda_{G_1}(v) + \lambda_{G_2}(v)$. Every other vertex keeps its neighbourhood. □

We use three published results.

Theorem 2.4 (Erdős, Saks and Sós [7])**.** *Every connected graph G has an induced path on at least $2r(G) - 1$ vertices.*

Theorem 2.5 (Camby and Schaudt [2])**.** *Let $k \geq 4$ and let G be a connected graph with no induced path on k vertices. Then G has a connected dominating set X such that $G[X]$ has no induced path on $k-2$ vertices or $G[X]$ is isomorphic to the cycle C_k.*

Theorem 2.6 (Seinsche [15])**.** *If a graph on at least two vertices has no induced path on four vertices, then it or its complement is disconnected.*

Theorem 2.5 is implication (i)⇒(ii) of Theorem 5 of arXiv:1402.7213, applied with $H = G$. In that preprint, Theorem 5 is the equivalence, for every connected induced subgraph, between P_k-freeness and the existence of such a connected dominating set. The journal article [2] states the same equivalence. We write P_k for the path on k vertices, and "G is P_k-free" means that G has no induced P_k, that is, $p(G) \leq k - 1$. A connected graph of radius r has at least $2r$ vertices.

Lemma 2.7. *Let H be a connected strong C_4-free P_4-free graph on at least two vertices. Then H has a vertex adjacent to all other vertices.*

Proof. By Theorem 2.6, H is the join of two nonempty vertex sets A and B. If $|A|, |B| \geq 2$, then $a, a' \in A$ and $b, b' \in B$ span the 4-cycle $aba'b'$. So one side is a single vertex, and it is adjacent to everything. □

Lemma 2.8. *Let G be a connected graph with no induced path on six vertices. Then G has a connected dominating set X such that $G[X]$ has no induced path on four vertices, or else $G[X]$ is isomorphic to C_6.*

This is Theorem 2.5 at $k = 6$. Lemma 3.5 uses the resulting dominating set.

Lemma 2.9 (Karthick and Maffray [10, Theorem 3.2(d)])**.** *Let G be a graph with no induced P_6, no induced C_4 and no induced C_6, and let $C = c_0c_1c_2c_3c_4c_0$ be an induced 5-cycle, indices modulo 5. For each i let W_i be the set of vertices outside C whose neighbourhood on C is exactly $\{c_i\}$, and let $X_{i,i+1}$ be the set of vertices outside C whose neighbourhood on C is exactly $\{c_i, c_{i+1}\}$. Then, for every i, at least one of $X_{i,i+1}$ and $X_{i+1,i+2}$ is empty, at least one of W_i and W_{i+2} is empty, and at least one of $X_{i,i+1}$ and $W_{i-1} \cup W_{i+2}$ is empty.*

Proof. The three assertions are Theorem 3.2(d) in Karthick and Maffray. The argument here is self-contained. Fix i and write indices modulo 5. Throughout, C is induced, a vertex of W_i meets C only at c_i, and a vertex of $X_{i,i+1}$ meets C only at c_i and c_{i+1}.
Let $x \in X_{i,i+1}$ and $y \in X_{i+1,i+2}$. If $x \sim y$, the vertices $x, c_i, c_{i-1}, c_{i+3}, c_{i+2}, y$ induce a 6-cycle. The vertex c_{i+1} lies off this set, so the edge c_ic_{i+1} is not a chord, x has no further neighbour among the six vertices except c_i and y, and y has no further neighbour among them except c_{i+2} and x. If $x \nsim y$, the path $y, c_{i+2}, c_{i+3}, c_{i-1}, c_i, x$ is an induced P_6: again only one cycle-neighbour of x lies on the path.
Let $x \in W_i$ and $y \in W_{i+2}$. If $x \sim y$, the vertices $x, c_i, c_{i-1}, c_{i+3}, c_{i+2}, y$ induce a 6-cycle. If $x \nsim y$, the same six vertices, in that order and without the edge xy, induce a P_6.
Let $x \in X_{i,i+1}$ and $y \in W_{i-1}$. If $x \sim y$, the vertices $y, c_{i-1}, c_{i+3}, c_{i+2}, c_{i+1}, x$ induce a 6-cycle. If $x \nsim y$, the same vertices in that order induce a P_6. If instead $y \in W_{i+2}$ and $x \sim y$, then x, c_{i+1}, c_{i+2}, y induce a 4-cycle, since $x \nsim c_{i+2}$ and $y \nsim c_{i+1}$. If $x \nsim y$, the path $y, c_{i+2}, c_{i+3}, c_{i-1}, c_i, x$ is an induced P_6. □

A strong C_4-free graph has no induced C_4. Whenever it is P_6-free and the case of an induced 6-cycle has been excluded, it meets the hypotheses of the lemma. Case 2 of Theorem 3.6 uses the first two assertions. The third assertion is part of the list in [10] and is not required there.

Lemma 2.10. *Let G be a connected triangle-free strong C_4-free P_7-free graph with no cut vertex. Then G has no clique cutset.*

Proof. The graph is triangle-free, so every clique has order at most two, and there is no clique cutset of order one. Suppose $\{u, v\}$ is a clique cutset. Then uv is an edge. Neither vertex is a cut vertex, so every component of $G - \{u, v\}$ is adjacent to both u and v, and there are at least two such components. A shortest u–v path through one component has at least four edges: one internal vertex would form a triangle with uv, and two internal vertices would form a 4-cycle with uv. On each of two such paths, let x, y, z be the three vertices immediately preceding v, in that order from the u-side, and write x', y', z' for the same choice in the second component. A shortest u–v path has at least four edges, so x, y, z and x', y', z' are internal vertices and in particular distinct from u. They therefore lie in two distinct components of $G - \{u, v\}$, and there is no edge between the two triples. Each triple is a subpath of a shortest path, so it has no chord. The path x, y, z, v, z', y', x' is an induced P_7, which contradicts $p(G) \leq 6$. □

3 | Graphs of induced-path order at most five

Lemma 3.1. *If G is connected and strong C_4-free with $r(G) = 1$, then $S(G) = 2(n-1) - m$, where m is the number of edges of $G[N(u)]$ for a dominating vertex u. Hence $\ell(G) < 2$, and (1.1) holds.*

Proof. Let $v \neq u$. By Lemma 2.1(1), v has at most one neighbour other than u, and such a neighbour lies in $N(u)$. So $G[N(v)]$ is a clique and $\lambda(v) = 1$; also $\lambda(u) = (n-1) - m$. Hence $S = 2(n-1) - m < 2n$ and $\lceil \ell \rceil \leq 2$. If G is complete then $G \in \{K_2, K_3\}$, $\ell = 1$ and $p = 2 = r + 1$. Otherwise two nonadjacent vertices of $N(u)$ together with u form an induced P_3, so $p \geq 3 \geq r + \lceil \ell \rceil$. □

Lemma 3.2. *If G is connected and strong C_4-free with $r(G) \geq 2$, then $p(G) \geq 4$.*

Proof. If G were P_4-free, Lemma 2.7 would give a dominating vertex, so $r(G) \leq 1$. □

Lemma 3.3. *If G is connected, strong C_4-free and P_5-free, then $S(G) \leq 2n - h$, where h is the number of vertices v with $\lambda(v) \geq 3$. In particular $S(G) \leq 2n$.*

Proof. Call v *simplicial* if $\lambda(v) = 1$ and *high* if $\lambda(v) \geq 3$. For $x \in N(v)$ put $X_x = N(x) \setminus N[v]$, and let P_x be the set of neighbours of x in $G[N(v)]$. The set P_x is empty when x is isolated in $G[N(v)]$, and otherwise it consists of the other end of the unique edge unit containing x. Thus $N(x) \subseteq \{v\} \cup P_x \cup X_x$.
Key fact. Let x, y lie in different units of v. A common vertex of X_x and X_y would give the 4-cycle $vxwy$, so $X_x \cap X_y = \emptyset$. If $x' \in X_x$ and $y' \in X_y$ are nonadjacent, then $x'xvyy'$ is an induced P_5: $x \nsim y$ because they lie in different units, $x \nsim y'$ and $y \nsim x'$ by disjointness, and v is adjacent to neither x' nor y'. So X_x is complete to X_y. If both are nonempty and $|X_x| \geq 2$, a vertex of X_y has two neighbours in X_x, which are also neighbours of x, a contradiction to Lemma 2.1(1). Hence if both are nonempty, then $|X_x| = |X_y| = 1$.
Main step. For each high v we find $\lambda(v) - 1$ simplicial vertices whose only high neighbour is v, one in each of $\lambda(v) - 1$ distinct units of v.
Case 1: some $u \in N(v)$ has $|X_u| \geq 2$. By the key fact $X_y = \emptyset$ for every y outside the unit of u. Then $N(y) \subseteq \{v\} \cup P_y$. This set is a clique, so y is simplicial, and so is the other vertex of its unit when that unit is an edge. Hence v is the only high neighbour of y, and the $\lambda(v) - 1$ units other than that of u supply the vertices.
Case 2: $|X_x| \leq 1$ for all $x \in N(v)$. Call a unit *active* if one of its vertices x has $X_x \neq \emptyset$. Suppose two units are active, with $y' \in X_y$ and $z' \in X_z$, and let x lie in a third unit. By the key fact $y' \sim z'$, and $xvyy'z'$ is an induced P_5: $x \nsim y$ (different units), $x \nsim y', z'$ (disjointness), $v \nsim y', z'$, and $y \nsim z'$ (disjointness). As $\lambda(v) \geq 3$, a third unit exists, so at most one unit is active. Every vertex of an inactive unit is simplicial and its only high neighbour is v. This gives at least $\lambda(v) - 1$ units.
Counting. The chosen vertices for distinct high vertices are distinct, since each has a unique high neighbour. So $\sum_{v \text{ high}} (\lambda(v) - 1) \leq \#\{\text{simplicial vertices}\}$, and

$$S = \sum_{v} \lambda(v) \leq 2n - \#\{\text{simplicial}\} + \sum_{v \text{ high}} (\lambda(v) - 2) \leq 2n - h. \qquad \square$$

Lemma 3.4. *Let G be connected, strong C_4-free and P_6-free with a dominating induced cycle $C = v_0v_1 \cdots v_5$, and let $W = V(G) \setminus V(C)$. Then every $w \in W$ is adjacent to exactly one antipodal pair $\{v_i, v_{i+3}\}$ and to no other vertex of C, distinct vertices of W use distinct pairs, and W is independent. Hence $|W| \leq 3$, $S(G) = 12 + 4|W| < 3n$, and $r(G) \leq 2$ if $W \neq \emptyset$.*

Proof. Indices are modulo 6. Let $w \in W$; it has a neighbour on C. If w has exactly one neighbour v_i, then $wv_iv_{i+1}v_{i+2}v_{i+3}v_{i+4}$ is an induced P_6. If $N(w) \cap C = \{v_i, v_{i+1}\}$, then $wv_{i+1}v_{i+2}v_{i+3}v_{i+4}v_{i+5}$ is an induced P_6. Two neighbours at distance 2 on C give a 4-cycle. A set of at least three vertices of C_6 contains two vertices at distance 2. So $N(w) \cap C$ is an antipodal pair. Two vertices of W with the same pair give two common neighbours. If $w \sim w'$ with distinct pairs, some $v_i \in N(w)$ and $v_j \in N(w')$ are adjacent on C, and wv_iv_jw' is a 4-cycle. There are no triangles, $e(G) = 6 + 2|W|$, and Lemma 2.1(3) gives $S = 12 + 4|W|$. Finally, if w uses $\{v_i, v_{i+3}\}$, then every vertex of C is within distance 2 of v_i (the vertex v_{i+3} through w), and each $w' \in W$ has a neighbour in $\{v_{i-1}, v_i, v_{i+1}\}$, because each of the three antipodal pairs meets this set. So v_i has eccentricity 2. $\square$

The same trace analysis applies to any vertex adjacent to an induced 6-cycle in a connected strong C_4-free P_6-free graph, whether or not the cycle dominates. Theorem 3.6 uses that form. Karthick and Maffray [10, Theorem 3.3] classify the external neighbourhood of an induced C_6 in a (P_6, C_4)-free graph. When the graph has no 4-cycle subgraph, only the empty trace and the antipodal-pair traces remain. The new content here is the second-layer constraint, the resulting order bound, the exact value of $M(n)$, and uniqueness.

Lemma 3.5. *Let G be connected, strong C_4-free and P_6-free. Then either G has a vertex of eccentricity at most 2, or $G = C_6$.*


Proof. Lemma 2.8 supplies a connected dominating set X. If $G[X] \cong C_6$, Lemma 3.4 gives $G = C_6$ or a vertex of eccentricity 2. Otherwise $G[X]$ is P_4-free. If $|X| = 1$, the vertex in X dominates G. If $|X| \geq 2$, Lemma 2.7 gives $u \in X$ adjacent to all of $X \setminus \{u\}$, and since X dominates G, every vertex is within distance 2 of u. $\square$

Theorem 3.6. *Let $M(n)$ be the maximum of $\ell(G)$ over connected strong C_4-free P_6-free graphs of order $n \geq 7$. Then*

$$M(n) = \begin{cases} 16/7 & \text{if } n = 7, \\ 5/2 & \text{if } n = 8, \\ 8/3 & \text{if } n = 9, \\ 3 & \text{if } n = 10, \\ 2 + 2/n & \text{if } n \geq 11. \end{cases}$$

For every $n \geq 7$ the extremal graph is unique up to isomorphism. For $n = 10$ it is the Petersen graph P. For $n = 9$ it is $P - v$. For $n = 8$ it is the graph H_8 of Figure 1. For $n = 7$ and for every $n \geq 11$ it is the graph Γ_{n-7} of that figure: a 6-cycle, a vertex

a adjacent exactly to one antipodal pair, and $n-7$ leaves adjacent only to a. For $2 \le n \le 6$ the same maximum $M(n)$, and the extremal graphs, are as follows: $n = 2$, $M(2) = 1$, only K_2; $n = 3$, $M(3) = 4/3$, only P_3; $n = 4$, $M(4) = 3/2$, exactly P_4 and $K_{1,3}$; $n = 5$, $M(5) = 2$, only C_5; $n = 6$, $M(6) = 2$, exactly C_6 and the graph obtained from C_5 by adding one pendant leaf. Since $M(n) \le 3$ for every $n \ge 2$, every connected strong C_4-free P_6-free graph G of order $n \ge 2$ satisfies $S(G) \le 3n$, with equality if and only if G is the Petersen graph.



(a) H_8 (b) $P - v$

(c) P (d) Γ_h

FIGURE 1 | Extremal graphs for $M(n)$. The six central filled vertices form the cycle, and each open square meets one antipodal pair. (a) H_8, order 8, $\ell = 5/2$. (b) $P - v$, order 9, $\ell = 8/3$. (c) The Petersen graph, order 10, $\ell = 3$. These three graphs are triangle-free, so $S = 2e$. (d) Γ_h: any number of leaves at the square a, including none. The case $h = 0$ is the unique graph of order 7 with $\ell = 16/7$. For every $n \ge 11$ the extremal graph is Γ_{n-7}, with $S = 2n+2$.

Proof. Throughout, $S = 2e - 3T$ by Lemma 2.1. A chordless cycle of length at least seven contains an induced P_6. Thus the only possible holes have lengths five and six.

Case 1: an induced six-cycle. Label its vertices $c_0, \dots, c_5$, indices modulo 6, and write A for the set of vertices at distance one from the cycle. The trace argument in the proof of Lemma 3.4 shows that each $a \in A$ meets the cycle in one antipodal pair. Distinct vertices of A use distinct pairs, and A is an independent set: an edge of A joins distinct pairs, and some contact of one end is adjacent on the cycle to some contact of the other, giving a 4-cycle. Hence $k = |A| \le 3$. Figure 1 shows the extremal graphs of this branch in panels (a)–(c), for $k \in \{2, 3\}$, and the family Γ_h in panel (d), for $k = 1$. Figure 2 shows the two distance layers used in the count.

Every vertex x at distance two from the cycle is adjacent to every vertex of A. Such an x has at least one neighbour in A. If $x \sim a$ and $x \nsim b$ for $a, b \in A$, choose contacts so that $c_i \in N(a)$ and $c_{i+2} \in N(b)$. This is possible: the pairs are distinct, so some contact of b lies at cycle distance two from some contact of a, and we orient the cycle from that contact of a toward that contact of b. The path $x, a, c_i, c_{i+1}, c_{i+2}, b$ is then an induced P_6. Indeed x meets the cycle in nothing, a meets the cycle only in its antipodal pair (neither c_{i+1} nor c_{i+2}), b meets the cycle only in its antipodal pair (so $b \sim c_{i+2}$ and $b \nsim c_i, c_{i+1}$), and $a \nsim b$.

There is no vertex at distance three or more from the cycle. If there were, the cycle would have a vertex z at distance exactly three. A shortest z–cycle path is then the path z, x, a, c_i on four vertices, with x at distance two and a at distance




FIGURE 2 | Distance layers of an induced 6-cycle in Case 1 of Theorem 3.6. Each a_i meets one antipodal pair, the pairs are distinct, and the first layer is independent, so $k \leq 3$. A vertex x of the second layer meets every vertex of the first layer and none of the cycle. If $k \in \{2, 3\}$ that layer has size at most one, and the order is at most 10. There is no third layer.

one. Extending that path by c_{i+1}, c_{i+2} gives an induced P_6: the vertex z has no neighbour in A or on the cycle, and x has no neighbour on the cycle.

Let h be the number of vertices at distance two. If $k = 0$, then A is empty, so the cycle has no neighbour and $G = C_6$. If $k = 1$, every distance-two vertex is adjacent to the unique vertex $a \in A$. The distance-two vertices induce a matching together with isolated vertices: a vertex of that layer with two neighbours in the layer would have two common neighbours with a. Writing m for the number of edges of the matching gives $n = 7+h$, $e = 8+h+m$ and $T = m$, so $S = 16+2h-m = 2n+2-m$. Thus $S \leq 2n + 2$, with equality if and only if $m = 0$. Equality means that G is the graph Γ_h formed by the 6-cycle, the vertex a on one antipodal pair, and h leaves at a.

The graph Γ_h lies in the class, for every $h \geq 0$. A leaf has degree one. Every other cycle through an outside vertex uses both edges from a to the cycle, and each of the two arcs of the cycle between the neighbours of a has length three, so every such cycle has length five. There is no 4-cycle. There is no induced P_6. A leaf, having degree one, occupies an end of any path that contains it, and that path meets a next. Two leaves therefore determine a path on only three vertices. An induced path that contains one leaf and five further vertices would restrict to an induced path of order five in Γ_0 that starts at a. No such path exists: the two neighbours of a on the cycle are an antipodal pair, the path $a, c_i, c_{i+1}, c_{i+2}, c_{i+3}$ has the chord ac_{i+3}, and every shorter continuation from a through one neighbour stops at the vertex before the antipode, because the next cycle vertex is adjacent to a. An induced path of Γ_0 that avoids a lies on the 6-cycle and has order at most five. An induced path of Γ_0 that contains a in its interior has c_i and c_{i+3} as the two neighbours of a on the path. Each side can then take at most one further cycle vertex: the next vertex on either arc is adjacent to the antipode already used. The resulting path has order at most five. Thus Γ_h is strong C_4-free and P_6-free, and it attains $S = 2n + 2$.

If $k \in \{2, 3\}$, then $h \leq 1$: two vertices of the second layer would both be adjacent to two vertices of A. There are no triangles. The second-layer vertex, if it exists, is adjacent to every vertex of A and to nothing on the cycle, A is independent, and each vertex of A contributes two edges to the cycle. Thus $n = 6 + k + h$ and $S = 2(6 + 2k + kh)$. The four possibilities are

(k, h)	$(2, 0)$	$(2, 1)$	$(3, 0)$	$(3, 1)$
S/n	$5/2$	$8/3$	$8/3$	3

The four pairs are the graphs of Figure 1, with $(3, 0)$ and $(2, 1)$ both equal to $P - v$. Write a_i for the vertex adjacent to the antipodal pair $\{c_i, c_{i+3}\}$. The pair $(k, h) = (3, 1)$ is the graph in panel (c): the three vertices a_0, a_1, a_2 and a vertex v adjacent to all three. It is cubic of order 10. It is triangle-free, and the trace rules give no 4-cycle, so its girth is at least 5. Its diameter is 2. Indeed every pair of cycle vertices is at distance at most 2 along the cycle or through the corresponding a_i; the vertex v meets every a_i; and a non-neighbour of a_i on the cycle is adjacent to c_i or to c_{i+3}. Suppose $v_0 \cdots v_5$ were an induced path. Diameter 2 gives v_0 a common neighbour with each of v_3, v_4, v_5. Those neighbours lie off the path, because the only neighbour of v_0 on the path is v_1 and the path is induced. They are distinct: a vertex adjacent to two of v_3, v_4, v_5 would form a triangle with a path edge, or a 4-cycle with the length-2 subpath $v_3v_4v_5$. This is impossible: v_0 has degree 3, so it has only two neighbours off the path. Thus the graph is P_6-free. It is the Petersen graph P, and $\ell(P) = 3$.

Deleting v leaves $(k, h) = (3, 0)$, the graph $P - v$ in panel (b). The pair $(2, 1)$ is isomorphic to the same graph. In the labelling with $A = \{a_0, a_1\}$ and second-layer vertex v, the cycle $c_0, a_0, c_3, c_4, a_1, c_1$ has its three antipodal pairs served by c_5,

v and c_2, and no vertex remains. Both graphs therefore have $\ell = 8/3$, and both are the graph $P - v$. Induced subgraphs of P are strong C_4-free and P_6-free. The pair $(2, 0)$ is the graph H_8 in panel (a), with $\ell = 5/2$. The three choices of the unused antipodal pair differ by a rotation, so this graph is unique up to isomorphism. Its order is 8, so it is neither P nor $P - v$. For $k \in \{2, 3\}$ one has $n = 6 + k + h \leq 10$.

Case 2: an induced five-cycle and no induced six-cycle. Write $C = c_0 \cdots c_4 c_0$. Every vertex at distance one from C has a singleton or consecutive-pair trace: any other pair of contacts lies at distance two on C and forms a 4-cycle. If a vertex x were at distance two, with a neighbour a in the first layer, the path x, a followed by four consecutive cycle vertices, beginning at one contact of a and omitting the second contact if a has one, would be an induced P_6. Thus C dominates G. Write h for the number of vertices with singleton trace and j for the number with a consecutive-pair trace. At most one vertex carries each of the five pair traces, since two would have two common neighbours, so $j \leq 5$. The edges outside C are of the following kinds.

- Two singleton vertices with different supports are nonadjacent. Neighbouring supports give a 4-cycle. Supports at distance two give an induced 6-cycle along the longer arc of C. Edges within one singleton class form a matching, because two neighbours in that class would be two common neighbours of the supporting cycle vertex. Let m be the number of these edges.
- No two pair vertices are adjacent. From their two distinct support edges one can choose adjacent cycle vertices, one from each edge, and those two vertices together with the pair vertices form a 4-cycle.
- A singleton supported at c_i meets only a pair vertex supported on the opposite edge $\{c_{i+2}, c_{i+3}\}$. Every other pair edge contains a neighbour of c_i and yields a 4-cycle. That pair vertex meets at most one vertex of the singleton class, again by the common-neighbour bound. Let b be the number of singleton-to-pair edges, so $b \leq \min(h, j)$.

Consequently $n = 5 + h + j$ and $e = 5 + h + 2j + m + b$. The triangles are the j pair triangles and the m matching triangles, so $T = j + m$ and

$$S = 2e - 3T = 2n + 2b - j - m.$$

The integer b is at most 1. Suppose two singleton–pair edges are present. They use two distinct singleton supports. The opposite edge of c_i is $\{c_{i+2}, c_{i+3}\}$. As i varies, these five edges exhaust C, and each pair class contains at most one vertex. The graph is P_6-free and strong C_4-free, and the present case assumes that it has no induced C_6, so Lemma 2.9 applies. Under that lemma the set W_i is the singleton class at c_i, and the set $X_{i,i+1}$ is the consecutive-pair class on the edge $\{c_i, c_{i+1}\}$. If the two singleton supports lie at distance 2 on C, they are W_i and W_{i+2} for some i, and both are nonempty. Lemma 2.9 states that at least one of those two sets is empty. If the supports lie at distance 1, take them to be c_i and c_{i+1}. The opposite edge of c_i is $\{c_{i+2}, c_{i+3}\}$ and the opposite edge of c_{i+1} is $\{c_{i+3}, c_{i+4}\}$. Both pair classes are nonempty, one for each of the two edges, and they are $X_{i+2,i+3}$ and $X_{i+3,i+4}$. Lemma 2.9 states that at least one of $X_{j,j+1}$ and $X_{j+1,j+2}$ is empty for every j; the index $j = i + 2$ is this pair of classes. Either alternative contradicts the existence of both edges, so $b \leq 1$.

If $b = 0$, then $S = 2n - j - m \leq 2n$. If $b = 1$, then $j \geq 1$ because $b \leq \min(h, j)$, so $S = 2n + 2 - j - m \leq 2n + 1$. In this case $S \leq 2n + 1$.

Case 3: no holes. The graph is chordal, and every nonempty induced subgraph has a simplicial vertex. The neighbourhood of a simplicial vertex is a clique. The degree is at most two: four vertices of a clique contain a 4-cycle. A simplicial vertex of a connected chordal graph on at least two vertices is not a cut vertex, and the graph left by deleting it is chordal and strong C_4-free. In the graph present at the deletion, the loss is $L(v) = 2d(v) - 3t(v)$. Degree one gives loss 2. Degree two, with the two neighbours adjacent, gives one triangle and loss 1. Deleting vertices until a single vertex remains, one has $S \leq 2(n - 1) = 2n - 2$.

The three cases give the values of $M(n)$. A graph of order $n \geq 11$ cannot realise $k \in \{2, 3\}$ in Case 1, and $k = 0$ forces $G = C_6$. The only remaining graphs in Case 1 are the graphs Γ_h and the graphs with the same first layer and $m \geq 1$. The latter have $S \leq 2n + 1$. Cases 2 and 3 give $S \leq 2n + 1$ and $S \leq 2n - 2$. Thus $M(n) = 2 + 2/n$, attained only by Γ_{n-7}.

The graphs with $S > 2n + 2$ are as follows. At order $n = 10$ the only graph in the list is P, with $S = 30$. At order $n = 9$ the only such graph is $P - v$, with $S = 24$, whether it arises as $(k, h) = (3, 0)$ or as $(k, h) = (2, 1)$. At order $n = 8$ the only such graph is H_8, with $S = 20$. At order $n = 7$ every graph in the list satisfies $S \leq 2n + 2$. One has $h = 0$ in the equality case of $k = 1$, so the unique graph with $S = 2n + 2$ is Γ_0, while Cases 2 and 3 give at most 15 and 12. Since $8/3 < 3$ and $2 + 2/n < 3$ for $n \geq 11$, the bound $S \leq 3n$ holds for every order $n \geq 2$, with equality only for P.

The case division does not use $n \geq 7$. At order 2 the only connected graph is K_2, with $S = 2$ and $\ell = 1$. At order 3 there is no cycle of length five or six, so only the chordal bound applies and $S \leq 4$. Equality holds for the tree P_3, while K_3 has $S = 3$. At order 4 the same bound gives $S \leq 6$, with equality for a tree, and the only trees on four vertices are P_4 and $K_{1,3}$. At order 5 the induced 6-cycle branch is empty. In the induced 5-cycle branch one has $h = j = 0$, so $G = C_5$ and $S = 10$. The chordal branch contributes at most 8. At order 6 the induced 6-cycle branch contributes only C_6, with $S = 12$. In the

induced 5-cycle branch one has $h + j = 1$, hence $b = 0$ and $S = 12 - j - m \leq 12$, with equality precisely when $j = m = 0$ and $h = 1$, the graph C_5 with one leaf. The chordal branch contributes at most 10. Thus the maximum is 12, attained by exactly those two graphs. □

4 | Radius four

Lemma 4.1. *Let G be connected, strong C_4-free and P_8-free. If an induced 8-cycle dominates G and $r(G) \geq 4$, then $G = C_8$.*

Proof. Write $C = c_0c_1 \cdots c_7c_0$, with indices modulo 8. For an outside vertex y the trace $A_y = N(y) \cap V(C)$ is nonempty. The trace A_y is not contained in any three consecutive vertices of C. Suppose $A_y \subseteq \{c_i, c_{i+1}, c_{i+2}\}$. The ends c_i and c_{i+2} are not both in A_y, since y, c_i, c_{i+1}, c_{i+2} would be a 4-cycle. Thus A_y is a singleton or a consecutive pair. A singleton $\{c_i\}$ gives the induced path

$$y, c_i, c_{i+1}, c_{i+2}, c_{i+3}, c_{i+4}, c_{i+5}, c_{i+6}.$$

A consecutive pair $\{c_i, c_{i+1}\}$ gives the induced path

$$y, c_i, c_{i-1}, c_{i-2}, c_{i-3}, c_{i-4}, c_{i-5}, c_{i-6},$$

in which the omitted contact c_{i+1} does not lie on the path. Either path is an induced P_8.
Fix a cycle vertex, say c_0. The vertices at cycle distance greater than two from c_0 are the three consecutive vertices c_3, c_4, c_5. The trace of any outside vertex meets the complementary set $\{c_7, c_0, c_1, c_2, c_6\}$, so every outside vertex is within distance three of c_0. The same holds with c_0 replaced by any vertex of C.
Suppose w lies outside C, and write $A = N(w) \cap V(C)$. No two vertices of A are at cycle distance two, for they would form a 4-cycle with w. A set of vertices of C in which every two distinct vertices are consecutive has size at most two, and every set of size at most two that is a singleton or a consecutive pair is contained in a triple of consecutive vertices. Both possibilities have been excluded. Therefore A contains two vertices at cycle distance three or four. Rotating, take them to be c_0 and c_d with $d \in \{3, 4\}$.
The antipode of c_0 is c_4. If $d = 4$, the path c_0wc_4 has length two. If $d = 3$, then $c_3 \sim c_4$, so $c_0wc_3c_4$ has length three. Every other vertex of C is already at cycle distance at most three from c_0. Every outside vertex is within distance three of c_0. Thus the eccentricity of c_0 is at most three, contradicting $r(G) \geq 4$.
Hence C has no vertex outside it, and $G = C_8$. □

Theorem 4.2. *A connected graph G is strong C_4-free and satisfies $r(G) = 4$ and $p(G) = 7$ if and only if either $G \cong C_8$, or G is obtained from a 6-cycle $C = v_0v_1 \cdots v_5$ by the following construction and by no further vertices or edges. For each i there is a nonempty set C_i of vertices outside C. Every vertex of C_i is adjacent to v_i and to no other vertex of C. There is no edge between C_i and C_j for $i \neq j$, and $G[C_i]$ is a matching together with isolated vertices. For each edge of C there is at most one further vertex, adjacent precisely to the two ends of that edge.*




FIGURE 3 | The radius-four case of Theorem 4.2. (a) The induced P_8 is $w_5, v_5, v_0, x, y, v_2, v_3, w_3$, with singleton traces $\{v_0\}$ and $\{v_2\}$ at x and y, and with w_5, w_3 the distance-four witnesses. (b) One leaf at each cycle vertex gives $S = 2n$. (c) Only local attachments are shown; the remaining nonempty attachment classes are omitted. An isolated leaf does not contribute to $b + m$. A matching edge in some C_i, and a vertex glued to a cycle edge, each increase $2n - S = b + m$ by 1.

Lemma 4.3. *Let G be obtained from a 6-cycle C by the construction of Theorem 4.2. Let P be an induced path in G of order at least three. Every vertex of P that does not lie on C is an endpoint of P. In particular P contains at most two vertices outside C, and any such vertex is an endpoint of P.*

Proof. Let z be a vertex of P outside C. If z is an isolated vertex of $G[C_i]$ for some i, then z has degree one, so z is an endpoint of every path that contains it.
If z lies on a matching edge of $G[C_i]$, the neighbourhood of z is $\{v_i, z'\}$, where z' is the other end of that edge, and this neighbourhood is a clique. Suppose z were internal on P. An internal vertex has two neighbours on P. Those neighbours are then v_i and z', which are adjacent, and the path uses the route v_izz' or $z'zv_i$. The edge v_iz' joins the two neighbours of z and skips z, so it is a chord of P. An induced path has no chord. Thus z is an endpoint of P. The same chord shows that P does not contain z, z' and v_i together.
If z was added on a cycle edge, the neighbourhood of z is that edge, again a clique. The same chord argument shows that z is an endpoint of P, and that P does not contain both ends of the cycle edge together with z.
A path has two endpoints. Every vertex of P outside C is one of them, so there are at most two such vertices. □

Proof of Theorem 4.2. First suppose that G is strong C_4-free, with $r(G) = 4$ and $p(G) = 7$. Apply Theorem 2.5 with $k = 8$. The hypotheses hold: $k \geq 4$, and G is connected and has no induced path on eight vertices because $p(G) = 7$. The conclusion is a connected dominating set X such that $G[X]$ is P_6-free or $G[X] \cong C_8$.
If $G[X] \cong C_8$, Lemma 4.1 gives $G = C_8$.
If $G[X]$ is P_6-free, apply Lemma 3.5 inside $G[X]$. A vertex of eccentricity at most two in $G[X]$ has eccentricity at most three in G, because X dominates G, contradicting $r(G) = 4$. Therefore $G[X] = C_6$. Let $C = v_0 \cdots v_5$ be this dominating induced 6-cycle, indices modulo 6, and let $W = V(G) \setminus V(C)$.
Traces. As in the proof of Lemma 3.4, strong C_4-freeness allows only three kinds of trace $N(w) \cap C$: a single vertex, a consecutive pair $\{v_i, v_{i+1}\}$, or an antipodal pair. If some w has trace $\{v_i, v_{i+3}\}$, then v_i is within distance two of every vertex of C, hence within distance three of every vertex of G, a contradiction. So there are no antipodal traces. Let C_i be the set of vertices with trace $\{v_i\}$.
Witnesses. Fix i. Since $r(G) = 4$, some vertex is at distance at least four from v_{i+3}. Every vertex of C is within distance three of v_{i+3}, and a vertex $w \in W$ is at distance at most $1 + \min\{d_C(v_{i+3}, x) : x \in N(w) \cap C\}$ from v_{i+3}, which is at most three unless the trace of w is $\{v_i\}$. So there is $w_i \in C_i$ at distance four from v_{i+3}. Every neighbour $z \in W$ of w_i lies in C_i: if z had a neighbour among $v_{i+2}, v_{i+3}, v_{i+4}$, then $d(w_i, v_{i+3}) \leq 3$; if z had the neighbour v_{i-1} or v_{i+1}, then $w_i, v_i, v_{i\pm1}, z$ would be a 4-cycle. So $N(w_i) \subseteq \{v_i\} \cup C_i$.
Edges in W. Figure 3 records the two configurations used below. Let $x, y \in W$ be adjacent.

- $x \in C_0$, $y \in C_{\pm1}$: $x, v_0, v_{\pm1}, y$ is a 4-cycle.
- $x \in C_0$, $y \in C_2$: the path $w_5, v_5, v_0, x, y, v_2, v_3, w_3$ of Figure 3(a) is an induced P_8. The 6-cycle is induced, so it contributes no chord among v_5, v_0, v_2, v_3. Each off-cycle vertex meets the cycle only at its designated vertex: w_5 at v_5, x at v_0, y at v_2 and w_3 at v_3. The witness neighbourhoods, already proved, satisfy $N(w_5) \subseteq \{v_5\} \cup C_5$ and $N(w_3) \subseteq \{v_3\} \cup C_3$, so neither witness is adjacent to x, to y, or to the other witness. These three facts leave no chord.
- $x \in C_0$, $y \in C_3$: $w_1, v_1, v_0, x, y, v_3, v_4, w_4$ is an induced P_8. By symmetry, vertices of C_i and C_j with $i \neq j$ are never adjacent.
- x with trace $\{v_0, v_1\}$ and $y \in C_j$: for $j \in \{0, 1\}$ the vertices v_0, v_1, x, y span a 4-cycle; for $j = 5$ (respectively $j = 2$) the cycle x, v_0, v_5, y (respectively x, v_1, v_2, y) is a 4-cycle; for $j = 3$, $w_5, v_5, v_0, x, y, v_3, v_2, w_2$ is an induced P_8, and $j = 4$ is symmetric.
- x, y with consecutive-pair traces $\{v_0, v_1\}$ and $\{v_j, v_{j+1}\}$: $j = 0$ gives two common neighbours; $j = \pm1$ and $j = \pm2$ give 4-cycles as above; $j = 3$ gives the induced P_8 $w_2, v_2, v_1, x, y, v_4, v_5, w_5$.

The remaining paths displayed above are induced for the same three reasons, applied to the witnesses and traces named in those bullets. So every edge inside W joins two vertices of the same class C_i, and these edges form a matching. A consecutive-pair vertex is adjacent to no vertex of any C_j and to no other consecutive-pair vertex. Two vertices with the same consecutive-pair trace would have two common neighbours, so each cycle edge carries at most one such vertex, and that vertex is adjacent precisely to the two ends of the edge. Each class C_i is nonempty. This is the stated construction.
Conversely, let G be C_8, or let G be built from $C = v_0 \cdots v_5$ as in the statement. The cycle C_8 has radius 4, induced-path order 7 and no 4-cycle subgraph. Now consider the construction on C. There is no edge between distinct sets C_i, and every vertex of C_i meets C only at v_i. A cycle that meets C_i is therefore contained in $G[C_i \cup \{v_i\}]$. In that subgraph the edges are a star at v_i together with a matching, so the only cycles are the triangles formed by v_i and one matching edge.

Let y be a vertex added on the edge $\{v_j, v_{j+1}\}$, so $N(y) = \{v_j, v_{j+1}\}$. Every cycle through y uses both edges at y, and the rest of the cycle is a v_j–v_{j+1} path avoiding y. A 4-cycle through y is then a path of length two between v_j and v_{j+1} that avoids y: the two ends of the edge have a common neighbour other than y. No vertex of any C_i is adjacent to two vertices of C, and the construction places at most one vertex on each cycle edge. Those two ends therefore have no common neighbour other than the vertex added on that edge. Hence no 4-cycle passes through y. The only cycle that uses no vertex outside C is C itself. Thus G has no 4-cycle subgraph.

Let P be an induced path of order at least three. By Lemma 4.3, every vertex of P outside C is an endpoint of P, so every other vertex of P lies on C. The cycle C has no induced path on six vertices. Hence $p(G) \leq 7$. Since each C_i is nonempty, one may choose $x \in C_0$ and $y \in C_2$. The path $x, v_0, v_5, v_4, v_3, v_2, y$ is induced: x and y meet the path only at v_0 and v_2, and C contributes no chord. Thus $p(G) = 7$.

There is no edge between distinct sets C_i, and a vertex added on a cycle edge has degree two on that edge, so it does not shorten distances along C. The distance from v_i to v_{i+3} is therefore three. A vertex of C_{i+3} is at distance four from v_i, and the eccentricity of v_i is at least four. It is at most four: every vertex of C_j is at distance $d_C(v_i, v_j) + 1 \leq 4$ from v_i, and every vertex added on a cycle edge is within distance four of v_i. A vertex of C_i is at distance five from a vertex of C_{i+3}. A vertex y added on the edge $\{v_j, v_{j+1}\}$ has only those two neighbours. The distances from v_j and from v_{j+1} to v_{j+3} are 3 and 2, neither neighbour of y is adjacent to v_{j+3}, and $y, v_{j+1}, v_{j+2}, v_{j+3}$ is a path of length three, so a vertex of C_{j+3} is at distance four from y. Every eccentricity is at least four, and $r(G) = 4$. □

Corollary 4.4. *Let G be connected and strong C_4-free with $r(G) = 4$ and $p(G) = 7$. In the construction of Theorem 4.2, let b be the number of vertices added on edges of the 6-cycle and let m be the number of matching edges in the sets C_i. Then $S(G) = 2n - b - m$. In particular $S(G) \leq 2n$, so $\ell(G) \leq 2$.*

Proof. For $G \cong C_8$ one has $n = 8, e = 8, T = 0$ and $S = 16 = 2n$, which is the formula with $b = m = 0$. In the construction, write $a = \sum_i |C_i|$. Then $n = 6 + a + b$. The edges are the six cycle edges, a edges from the sets C_i to the cycle, m matching edges and $2b$ edges incident with the vertices added on cycle edges, so $e = 6 + a + 2b + m$. Each matching edge forms one triangle with its cycle vertex, and each added vertex forms one triangle with its cycle edge, so $T = b + m$. Lemma 2.1(3) gives $S = 2e - 3T = 2n - b - m$. □

Corollary 4.5. *Let G be connected and strong C_4-free with $r(G) = 4$ and $p(G) = 7$. Then $S(G) = 2n$ if and only if $G \cong C_8$, or G is obtained from C_6 by attaching at least one leaf to each cycle vertex.*

Proof. By Corollary 4.4, $S = 2n$ if and only if $G \cong C_8$ or, in the construction, $b = m = 0$. In the latter case every vertex outside the cycle is a leaf, and each C_i is nonempty, so each cycle vertex has at least one leaf. The converse is the case $b = m = 0$ of the construction, together with C_8, both of which have $S = 2n$ and satisfy the numerical hypotheses by Theorem 4.2. □

5 | Graphs of induced-path order at most six

In a triangle-free graph, a 4-cycle subgraph is an induced 4-cycle, and every clique has order at most two. The next classification is Lemma 3.4 of Chen and Xu [5]. Their family $\mathfrak{F} = \{F_0, F_0', F_1, \ldots, F_{12}\}$ is defined by the constructions in the proofs of their Lemmas 3.1–3.3; F_5 is the Petersen graph, and the parameters t and t' attached to F_2 and F_0' are positive integers. The edge counts below are taken from those constructions, as reproduced in the list that follows.

Lemma 5.1 (Chen and Xu [5, Lemma 3.4])**.** *Let G be a connected (P_7, C_4, C_3)-free graph with no clique cutset. If G is not complete, then $G \in \mathfrak{F}$.*

The members of $\mathfrak{F}$ admit the following descriptions, taken from those proofs.

- F_1 is C_5. The graph F_2 is obtained from a 5-cycle $v_1 \cdots v_5$ by adding stable sets A_1 and A_3 of equal positive order t, with a perfect matching between them, each vertex of A_1 adjacent only to v_1 and its match, and each vertex of A_3 adjacent only to v_3 and its match.
- On the same 5-cycle, F_3 has three further vertices inducing the path $a_3a_1a_4$, with a_i adjacent to v_i. The graph F_4 has four further vertices inducing the path $a_5a_3a_1a_4$, with the same spokes. The graph F_5 is the Petersen graph.
- F_0 is C_6. The graph F_0' is obtained from a 6-cycle $v_1 \cdots v_6$ by adding stable sets Y_1 and Y_4 of equal positive order t', with a perfect matching between them, each vertex of Y_1 adjacent only to v_1 and its match, and each vertex of Y_4 adjacent only to v_4 and its match.

- F_6 is C_7. On a 7-cycle $v_1 \cdots v_7$, write x_i for a vertex with neighbourhood $\{v_i, v_{i+3}\}$ on the cycle. Then F_7 is the cycle plus x_1; F_8 is the cycle plus nonadjacent x_1, x_2; F_9 is F_8 plus a vertex adjacent only to x_1 and x_2; F_{10} is the cycle plus nonadjacent x_1, x_5; F_{11} is the cycle plus a stable set $\{x_1, x_2, x_5\}$; and F_{12} is F_{11} plus a vertex adjacent only to x_1 and x_2.

Chen and Xu record that F_0, F_1, the graph F_2 with $t = 1$, F_3 and F_4 are induced subgraphs of the Petersen graph, and that $F_6, \dots, F_{11}$ are induced subgraphs of F_{12}. The graphs $F_6, \dots, F_{12}$ are the extensions of C_7 by vertices of cycle-distance three. The graphs F_2 and F_0' are the matching extensions of one pair of opposite vertices on C_5 and on C_6.

Lemma 5.2. *Every graph in $\mathfrak{F}$ other than the Petersen graph satisfies $2e \le 3n - 3$. The Petersen graph satisfies $2e = 3n$.*

Proof. The constructions give the following orders and sizes. For F_2 and F_0' the parameters t, t' are at least one.

Graph	n	e	$2e$	$3n - 2e$
$F_1 = C_5$	5	5	10	5
F_2	$5 + 2t$	$5 + 3t$	$10 + 6t$	5
F_3	8	10	20	4
F_4	9	12	24	3
F_5	10	15	30	0
$F_0 = C_6$	6	6	12	6
F_0'	$6 + 2t'$	$6 + 3t'$	$12 + 6t'$	6
$F_6 = C_7$	7	7	14	7
F_7	8	9	18	6
F_8, F_{10}	9	11	22	5
F_9, F_{11}	10	13	26	4
F_{12}	11	15	30	3

The five edges of F_3 beyond C_5 are the three spokes and the two edges of $a_3a_1a_4$. The seven edges of F_4 beyond C_5 are the four spokes and the three edges of $a_5a_3a_1a_4$. Each vertex x_i contributes two edges, and each vertex placed on a pair x_ix_j contributes two further edges. The Petersen graph is the only row with difference zero. □

Lemma 5.3. *Let G be connected and strong C_4-free, and let s be a vertex at which no induced P_4 starts. Write $m = e(G[N(s)])$. Then $S(G) \le 2(|G| - 1) - m$.*

Proof. A vertex at distance 3 from s would end a shortest path, hence an induced P_4, starting at s. So $V = \{s\} \cup N_1 \cup N_2$ with N_i the i-th distance layer. Each $y \in N_2$ has a unique parent $a \in N_1$. A neighbour $z \in N_2$ of y with a different parent would give the induced path $sayz$. So y has at most one neighbour in N_2, a sibling (two siblings would be common neighbours of a and y), and $N(y)$ is a clique; thus $\lambda(y) = 1$. For $a \in N_1$ we have $N(a) \subseteq \{s, \bar{a}\} \cup C_a$, where C_a is the set of children of a and $\bar{a}$ is the neighbour of a in N_1, if any. Since s and $\bar{a}$ lie in one unit, $\lambda(a) \le 1 + |C_a|$. Finally $\lambda(s) = |N_1| - m$. Summing, $S \le (|N_1| - m) + (|N_1| + |N_2|) + |N_2| = 2(|G| - 1) - m$. □

Lemma 5.4. *Let G be a connected triangle-free strong C_4-free P_7-free graph of order n. Then G is obtained from a connected triangle-free strong C_4-free P_7-free graph Q with no clique cutset by attaching trees at single vertices. The core Q is K_1, K_2, or a graph in the list of Lemma 5.2. Writing $h = |V(Q)|$ and $\eta(Q) = 3h - 2e(Q)$, one has $e(G) = e(Q) + (n - h)$ and*

$$3n - 2e(G) = \eta(Q) + n - h.$$

Deleting the vertices of one attached tree, other than its root on Q, deletes exactly as many edges as vertices.

Proof. If G itself has no cut vertex, Lemma 2.10 gives that G has no clique cutset. The hypotheses of that lemma are the present hypotheses together with the absence of a cut vertex. Lemma 5.1 then applies. Its hypotheses are that G is connected, that G has no induced P_7, that G has no induced C_4 and no triangle, and that G has no clique cutset. Connectedness and the absence of a clique cutset hold by the case and by Lemma 2.10. The graph is P_7-free by hypothesis. A strong C_4-free graph has no induced C_4, and a triangle-free graph has no C_3. If G is not complete, Lemma 5.1 places it in the family $\mathfrak{F}$, hence on the list of Lemma 5.2. The only complete triangle-free graphs are K_1 and K_2. In every subcase one may take $Q = G$. The identities are then immediate. The core K_1 is the whole graph only when $n = 1$.
Suppose s is a cut vertex. Two induced paths of order four that start at s and otherwise lie in distinct components of $G - s$ concatenate to an induced P_7. Thus some component C of $G - s$ contains no induced P_4 starting at s, and every induced path of $G[C \cup \{s\}]$ that starts at s has order at most three. The first distance layer of this side is independent, since an edge there would form a triangle with s. There is no vertex at distance three or more from s, for such a vertex would end an induced P_4 at s. A vertex of the second layer has a unique parent: a second parent, together with s, would span a 4-cycle. It has no neighbour in its layer. An edge joining vertices of distinct parents is the induced path of order four recorded in the proof of Lemma 5.3, and an edge joining two children of the same parent forms a triangle with that parent. Thus $G[C \cup \{s\}]$ is a tree.

Deleting the $k = |C|$ vertices of the side deletes exactly k edges. The graph that remains is connected, triangle-free, strong C_4-free and P_7-free. Iterate. The remaining graph Q has no cut vertex, and the first paragraph describes it. Reversing the deletions attaches trees at single vertices. Each deleted vertex accounted for one edge, so $e(G) = e(Q) + (n - h)$, and the displayed identity follows. The same deletion shows that K_1 is the core of every tree in the class, of any order. A connected P_7-free tree is triangle-free and strong C_4-free, and deleting leaves until one vertex remains leaves that core. □

Theorem 5.5. *Let G be a connected triangle-free strong C_4-free P_7-free graph. Then $2e(G) \le 3|V(G)|$, with equality if and only if G is the Petersen graph.*

Proof. Lemma 5.4 gives a core Q of order h and the identity

$$3n - 2e(G) = \eta(Q) + n - h.$$

Lemma 5.2 gives $\eta(Q) \ge 0$ for every graph on its list, with equality only for the Petersen graph. The cores K_1 and K_2 have deficits 3 and 4. The right-hand side is therefore nonnegative, and it vanishes if and only if $\eta(Q) = 0$ and $n = h$. Thus equality holds if and only if G is the Petersen graph. □

Lemma 5.1 classifies the connected graphs in the class with no clique cutset. The extremal count is obtained by attaching trees at single vertices of such a core and comparing the resulting deficits, for every order $n \ge 10$. Figure 4 draws the families that meet the minimum.

Theorem 5.6. *Let $f(n)$ be the maximum of $2e(G)$ over connected triangle-free strong C_4-free P_7-free graphs of order n. For every $n \ge 10$,*

$$f(n) = 3n - \min\{n - 10,\ 5 + \mathbf{1}_{\{n \text{ even}\}}\}.$$

Write $\mathcal{P}(n)$ for the isomorphism classes of graphs obtained from the Petersen graph by placing $n - 10$ leaves on a single vertex or on the two endpoints of a single edge. For $10 \le n \le 14$ the extremal graphs are exactly the classes in $\mathcal{P}(n)$. For $n = 15$ they are the classes in $\mathcal{P}(15)$ together with F_2, four classes in all. For $n = 16$ they are the classes in $\mathcal{P}(16)$, the graph F_0', and the two graphs obtained from the order-15 graph F_2 by adding a leaf at a hub or at the midpoint of its unique 2-edge branch, seven classes in all. For odd $n \ge 17$ the unique extremal graph is F_2. For even $n \ge 18$ there are exactly three: F_0', and the two graphs obtained from the order-$(n-1)$ graph F_2 by adding a leaf at a hub or at the midpoint of its unique 2-edge branch.




FIGURE 4 | Equality families of Theorem 5.6. (a) and (b) Leaves of $\mathcal{P}(n)$ at one vertex, or at both ends of one edge. Both have deficit $n - 10$. (c) F_2: length-3 paths on distinct arcs, and one length-2 path through v_2. Deficit 5. Rings mark both hubs and the midpoint. (d) F_0': every hub-to-hub path has length 3, on its own arc. Deficit 6. (e) and (f) The allowed leaf sites, a hub and the short-branch midpoint. The isomorphism classes at orders 15 and 16 are listed in Table 1.

Proof. Write $\eta(H) = 3|V(H)| - 2e(H)$ for a triangle-free graph H.
Reduction to one core. Lemma 5.4 supplies a core Q of order h and the identity $3n - 2e(G) = \eta(Q) + n - h$.

Upper bound. Every fixed row of the table in Lemma 5.2 satisfies $\eta \geq h - 10$. Each such graph has order at most 10 except F_{12}, which has order 11 and $\eta = 3$, and $3 \geq 11 - 10$. The same inequality holds for K_1 and K_2, with deficits 3 and 4. For every such core, $\eta(Q) + n - h \geq n - 10$. The two parameterised rows are F_2, of odd order and deficit 5, and F_0', of even order and deficit 6. If the core is F_2 and the final order is even, then at least one vertex has been attached, so the displayed deficit is at least 6. If the core is F_2 and the final order is odd, the deficit is at least 5. If the core is F_0', the deficit is at least 6. In every case

$$3n - 2e(G) \geq \min\{n - 10,\ 5 + \mathbf{1}_{\{n \text{ even}\}}\}.$$

Attainment. The value $f(n)$ is attained by the graphs named in the statement; the next paragraph shows that no other graph attains it. A leaf added to a triangle-free strong C_4-free graph creates neither a triangle nor a cycle. Realise the Petersen graph on the 2-element subsets of $\{1, 2, 3, 4, 5\}$, with two subsets adjacent when they are disjoint. This is the graph P of Theorem 3.6. The symmetric group on the ground set acts on the vertices. The action is vertex-transitive and edge-transitive, and it carries any two nonadjacent vertices to $\{1, 2\}$ and $\{1, 3\}$. The path

$$\{1, 2\}, \{3, 4\}, \{1, 5\}, \{2, 3\}, \{1, 4\}$$

is an induced P_5 starting at $\{1, 2\}$, and

$$\{1, 2\}, \{3, 4\}, \{1, 5\}, \{2, 4\}, \{1, 3\}$$

is an induced P_5 with ends $\{1, 2\}$ and $\{1, 3\}$. Thus every vertex of P is an end of an induced P_5, and every pair of nonadjacent vertices is the pair of ends of an induced P_5.

Place a leaves at one end of an edge of P and b leaves at the other, with $a + b = n - 10$ and $a \geq b \geq 0$. The case $b = 0$ is the placement at a single vertex. The graph is connected, triangle-free and strong C_4-free. It has $15 + (n - 10)$ edges, so its deficit is $n - 10$. An induced path uses a leaf only as an end. A path that uses two leaves at the same vertex has order three. A path that uses one leaf at each end of the chosen edge is the edge itself with the two leaves attached: the ends of that edge are adjacent, so no longer induced path of P has those ends. A path that uses one leaf restricts to an induced path of P and has order at most six, because P is P_6-free. Thus the graph is P_7-free. Vertex-transitivity and edge-transitivity give one isomorphism class for each pair (a, b). Distinct pairs give nonisomorphic graphs: the Petersen graph is cubic, so the degrees of the vertices adjacent to a leaf are $3 + a$ and $3 + b$ when $b \geq 1$, and the single value $3 + a$ when $b = 0$. These are the classes in $\mathcal{P}(n)$, and they attain deficit $n - 10$.

For the deficit $5 + \mathbf{1}_{\{n \text{ even}\}}$, use F_2 at odd orders and F_0' at even orders. In the descriptions preceding Lemma 5.2, each of these graphs consists of internally disjoint three-edge paths between two hubs, together with one additional two-edge path in the case of F_2. Every cycle uses two of those paths and therefore has length at least five. An induced path that contains at most one hub meets that hub in at most two of the paths and so has at most five vertices. An induced path that contains both hubs uses at most one path between them, a segment of at most three edges, and each end extends by at most one further edge before the next vertex meets the opposite hub and creates a chord. Its order is at most six. The table of Lemma 5.2 gives deficits 5 and 6 respectively. The smaller of the two deficits is therefore attained.

Equality. Write $\delta(n)$ for the minimum in the statement, and let G satisfy $3n - 2e(G) = \delta(n)$. The identity $3n - 2e(G) = \eta(Q) + n - h$ and the upper bound force the core Q to be the Petersen graph, F_2 or F_0'. Every other core in the table of Lemma 5.2, and each of K_1 and K_2, has $\eta(Q) - h \geq -8$, hence deficit at least $n - 8$. For $n \geq 10$ one has $n - 8 > \delta(n)$, since $\delta(n) \leq n - 10$.

If Q is the Petersen graph, then $\eta(Q) = 0$, $h = 10$ and the deficit is $n - 10$. This equals $\delta(n)$ if and only if $10 \leq n \leq 16$. The vertices outside Q form trees attached at single vertices of Q. A tree of depth at least two contains a path y, a, s of length two ending at its attachment vertex s. Concatenating that path with an induced P_5 of Q that starts at s produces an induced P_7: the only edge from $\{y, a\}$ into Q is as, and a is not adjacent to the neighbour of s on the P_5 because that neighbour and a would form a triangle with s. Thus every attached tree is a star of leaves at its attachment vertex.

The set A of vertices of Q that are adjacent to at least one leaf is a clique. Suppose $u, v \in A$ are distinct and nonadjacent, and let ℓ_u and ℓ_v be leaves at u and at v. The paragraph on attainment supplies an induced P_5 of Q with ends u and v; write it $u = q_0, q_1, q_2, q_3, q_4 = v$. The path $\ell_u, q_0, q_1, q_2, q_3, q_4, \ell_v$ has seven vertices. It is induced. The leaf ℓ_u is adjacent only to q_0, hence to none of $q_1, q_2, q_3, q_4, \ell_v$. The leaf ℓ_v is adjacent only to q_4, hence to none of $\ell_u, q_0, q_1, q_2, q_3$. The path $q_0q_1q_2q_3q_4$ is induced, so it contributes no chord. This is an induced P_7, which is forbidden. Thus no two vertices of A are nonadjacent, and A is a clique. The Petersen graph is triangle-free, so a clique has size at most two. If $|A| = 2$, the two vertices are adjacent. The attachments therefore occupy a single vertex or the two ends of one edge. These are the classes in $\mathcal{P}(n)$.

If $Q = F_2$, then $\eta(Q) = 5$ and $h = 5 + 2t$ is odd, so the deficit is $5 + n - h$. For odd n this equals $\delta(n) = 5$ if and only if $n = h$ and $n \geq 15$, in which case G is F_2. For even n the orders n and h have opposite parity, so $n - h \geq 1$ and the deficit is at least 6. Equality with $\delta(n)$ holds if and only if $n - h = 1$ and $\delta(n) = 6$, hence $n \geq 16$ is even and G is F_2 of order $n - 1$ with a single leaf. Here $t = (n - 6)/2 \geq 5$.

If $Q = F_0'$, then $\eta(Q) = 6$ and the deficit is $6 + n - h \geq 6$, with equality if and only if $n = h$. Then $\delta(n) = 6$, so $n \geq 16$ is even and G is F_0'.

It remains to decide which vertices of F_2 may carry the leaf. Adding a leaf at a vertex v of a P_7-free graph preserves P_7-freeness if and only if v is not an end of an induced P_6: the leaf has degree one, so it is an end of every induced path that contains it, and the remainder is an induced path starting at v. Label F_2 as in the description before Lemma 5.2. The vertex set consists of the hubs v_1 and v_3, the midpoint v_2 of the unique 2-edge branch $v_1v_2v_3$, and the internal vertices of the long branches. The long branches are the t matching paths through $A_1 \cup A_3$ and the path $v_1v_5v_4v_3$. The map that swaps v_1 with v_3, swaps v_5 with v_4, fixes v_2, and swaps the two ends of each matching edge preserves the cycle, the matching and every spoke, so it is an automorphism. It exchanges the two hubs.

No induced P_6 starts at a hub. Let P be an induced path starting at v_1. If the second vertex is v_2, the third is v_3, and the fourth is either v_4 or a vertex of A_3. In the first subcase the only continuation is v_5, and v_1v_5 is a chord, so the induced path has order at most four. In the second the only continuation is the match in A_1, and that match is adjacent to v_1, so again the order is at most four. If instead the path leaves v_1 along a long branch, it may continue to v_3 and then to the v_3-end of one further long branch. The resulting path has order five. Its last vertex has no neighbour off that path except its partner on the same long branch, and that partner is adjacent to v_1. The path does not extend, and the edge v_1v_2 is a chord of any continuation through v_2. Thus no induced P_6 starts at v_1. The automorphism gives the same conclusion at v_3.

No induced P_6 starts at v_2. Every induced path that starts at v_2 has second vertex v_1 or v_3. From v_1 it can enter only one long branch, and the far end of that branch is v_3, which is adjacent to v_2. The path stops before v_3 and has order at most four.

Every other vertex is an end of an induced P_6. The remaining vertices are the internal vertices of the long branches, namely $A_1 \cup A_3 \cup \{v_4, v_5\}$. Let p be such a vertex and adjacent to v_1, so $p \in A_1$ or $p = v_5$. The automorphism carries these vertices to the vertices adjacent to v_3. Since $t \geq 5$, at least two long branches avoid p. Choose two matching branches v_1abv_3 and v_1cdv_3 among them, with $a, c \in A_1$ and with $b, d \in A_3$ their matches, and consider the path p, v_1, a, b, v_3, d. The edges pv_1, v_1a, ab, bv_3 and v_3d are present by construction. There is no chord. A vertex of A_1 is adjacent only to v_1 and its match, a vertex of A_3 is adjacent only to v_3 and its match, A_1 and A_3 are stable sets, and the hubs are nonadjacent. If $p \in A_1$, its match is neither b nor d, so p meets the path only at v_1. If $p = v_5$, the only neighbours of v_5 are v_1 and v_4, and v_4 is not on the path, so again p meets the path only at v_1. The vertex a meets the path only at v_1 and b, and b meets the path only at a and v_3. Thus the path is an induced P_6, and a leaf at p extends it to an induced P_7.

A leaf at a hub or at v_2 therefore keeps the graph P_7-free, and a leaf at any other vertex does not. These are the only legal positions. The two hubs are equivalent under the automorphism, while v_2 is the unique common neighbour of the hubs, so the two placements are not isomorphic. In F_2 the hub v_1 has degree $2 + t$, because it is adjacent to v_2, to v_5 and to the t vertices of A_1, and v_2 has degree 2. After the leaf is added, the neighbour of the leaf has degree $t + 3$ or degree 3 according as it is a hub or v_2. For $t \geq 5$ these degrees differ.

The graphs that meet $\delta(n)$ are pairwise nonisomorphic. They are separated by the number of vertices of degree one, together with the multiset of degrees of the vertices adjacent to a leaf. A graph with a single leaf is separated by the degree of the neighbour of that leaf. For a pair (a, b) those degrees are $3 + a$ and, when $b \geq 1$, $3 + b$. The graphs F_2 and F_0' have minimum degree two. On a one-leaf extension of F_2 the neighbour of the leaf has degree $t + 3$ or degree 3, and these differ for $t \geq 5$. The order-15 core is F_2 with $t = 5$. At order 16 the underlying core has order 15, so that degree is 8 or 3. At even order $n \geq 18$ the underlying core is F_2 of order $n - 1$, and the degree is $n/2$ or 3. Table 1 is the resulting count. The order-15 and order-16 rows are the families drawn in Figure 4. □

n	Equality graphs	Leaves	Degrees at the leaf support	Classes
$10 \leq n \leq 14$	$\mathcal{P}(n)$, pairs (a, b)	$n - 10$	none ($n = 10$); else $\{3 + a\}$ or $\{3 + a,\ 3 + b\}$	one per pair
15	$\mathcal{P}(15)$: $(5, 0)$, $(4, 1)$, $(3, 2)$; and F_2, $t = 5$	5 or 0	$\{8\}$, $\{7, 4\}$, $\{6, 5\}$, or none	4
16	$\mathcal{P}(16)$: $(6, 0)$, $(5, 1)$, $(4, 2)$, $(3, 3)$; F_0'; two one-leaf graphs	6, 0 or 1	$\{9\}$, $\{8, 4\}$, $\{7, 5\}$, $\{6, 6\}$; none; 8 or 3	7
odd $n \geq 17$	F_2	0	none	1
even $n \geq 18$	F_0' and the two one-leaf graphs	0 or 1	none, or $n/2$, or 3	3

TABLE 1 | Separation of the equality graphs of Theorem 5.6 by the number of leaves and the degrees of the vertices adjacent to a leaf.

For $2 \leq n \leq 9$ the same identity $3n - 2e(G) = \eta(Q) + n - h$ gives

$$f(2), f(3), \ldots, f(9) = 2, 4, 6, 10, 12, 16, 20, 24.$$

For $n \le 4$ every connected member is a tree, so $2e = 2n - 2$. The later values are attained by C_5, by both C_6 and C_5 with one leaf, by F_2 with $t = 1$, by F_3 and by F_4. The graphs C_5, C_6, F_2 with $t = 1$, F_3 and F_4 lie on the list of Lemma 5.2. The graph C_5 with one leaf is that core with a single attached leaf, as in Lemma 5.4. Every other core of order at most n, after the attachment of $n - h$ vertices, has weakly smaller $2e$.

Theorem 5.7. *Let G be a connected strong C_4-free P_7-free graph with at most one triangle. Then $S(G) \le 3n$.*

Proof. Suppose G is a counterexample of minimum order in the stated class, so $S(G) \ge 3n + 1$. Theorem 5.5 forces G to contain a triangle, hence exactly one. Let v be a cut vertex. For each component C of $G - v$ let ρ_C be the largest order of an induced path in $G[C \cup \{v\}]$ with end v. Two such paths from different components form an induced path on $\rho_C + \rho_{C'} - 1 \le 6$ vertices, so some component C_1 has $\rho_{C_1} \le 3$. Put $G_1 = G[C_1 \cup \{v\}]$ and $G_2 = G - C_1$, with $n_i = |G_i|$, so $n = n_1 + n_2 - 1$ and $n_1 \ge 2$. Both graphs are connected, smaller than G, strong C_4-free and P_7-free, and each has at most one triangle. Lemma 5.3 gives $S(G_1) \le 2(n_1 - 1)$, and minimality gives $S(G_2) \le 3n_2$. By Lemma 2.3,

$$S(G) \le 2(n_1 - 1) + 3n_2 = 3n - (n_1 - 1) < 3n.$$

Thus G is 2-connected.

Let v be a vertex of minimum degree among the three vertices of the triangle, write $d = d(v)$, and let u and w be the other two. The graph $H = G - v$ is connected and triangle-free. Apply Lemma 5.4 to H, and write Q for its core, $h = |V(Q)|$ and $\eta = \eta(Q)$. Set

$$a = \big|N_G(v) \cap V(Q)\big|, \qquad c = \big|V(H) \setminus \big(V(Q) \cup N_G(v)\big)\big|,$$

and $t = |V(H)| - h$. The vertices of H outside Q are the $d - a$ neighbours of v that miss Q, together with the c vertices counted above, so $t = d - a + c$. Lemma 5.4 gives $e(H) = e(Q) + t$. Since G has one triangle, $S(G) = 2e(G) - 3$ and $e(G) = e(H) + d$, and therefore

$$S(G) - 3n = 2d - 6 - \eta - t = d + a - 6 - \eta - c. \tag{5.1}$$

The core satisfies $a \le \eta + 3$ and, for every edge xy of Q,

$$a + \min\{d_Q(x), d_Q(y)\} \le \eta + 5. \tag{5.2}$$

These bounds are read from the list in Lemma 5.2, using the descriptions that precede that lemma. If Q is the Petersen graph, then $\eta = 0$. Every two nonadjacent vertices of the Petersen graph have a common neighbour, so two such vertices cannot both be adjacent to v. Thus $N(v) \cap V(Q)$ is a clique, necessarily of size at most two, and every degree in Q is three. Both bounds follow. For each remaining fixed graph of order at least three one has $h \le 11$ and minimum degree at least two. The open neighbourhoods in Q of any two vertices of $N(v) \cap V(Q)$ are disjoint: a common neighbour would form a 4-cycle with v. Hence $a \le \lfloor h/2 \rfloor \le 5$. Every degree in these graphs is at most four. The only rows with $\eta < 4$ are F_4 and F_{12}, both with $\eta = 3$. The path-and-spoke description of F_4 has maximum degree three. In F_{12} the only vertices of degree four are v_1 and v_5 on the 7-cycle. Each is adjacent to two cycle vertices and to two of $\{x_1, x_2, x_5\}$, and v_1 is not adjacent to v_5, so every edge of F_{12} has an end of degree at most three. This gives (5.2) and the bound on a. In each parameterised family every non-hub vertex is adjacent to a hub, and each hub has at most one neighbour in common with v, so $a \le 4$ even after both hubs are counted. Every edge has an end of degree two, because the edges are the internal edges of the paths between the hubs described above, and η is 5 or 6. The graphs K_1 and K_2 satisfy both bounds directly: K_1 has $\eta = 3$ and no edge, and K_2 has $\eta = 4$.

If u or w lies outside Q, say u, the tree attachments give u at most one neighbour in Q. Apart from w, no neighbour of u in H is adjacent to v, because the triangle is unique. Thus $d \le d(u) \le c + 3$. Substituting in (5.1) gives $S(G) - 3n \le a - \eta - 3$, and $a \le \eta + 3$ yields $S(G) - 3n \le 0$. If instead both u and w lie in Q, then uw is an edge of Q, and every neighbour of either vertex that lies in $H - V(Q)$ lies in the set counted by c. Since v has minimum degree among the three triangle vertices,

$$d \le 1 + c + \min\{d_Q(u), d_Q(w)\}.$$

Substituting in (5.1) and applying (5.2) to the edge uw again yields $S(G) \le 3n$. □

Lemma 5.8. *Let G be strong C_4-free, and let T_1 and T_2 be two distinct triangles of G.*

1. *Suppose T_1 and T_2 share exactly one vertex u, and write $T_1 = \{u, a, b\}$ and $T_2 = \{u, c, d\}$. If ac is an edge, then $a\,c\,d\,u$ is a 4-cycle. The same holds for each of the other three edges with one end in $\{a, b\}$ and one end in $\{c, d\}$.*
2. *Suppose T_1 and T_2 are disjoint, and let xy and $x'y'$ be distinct edges with $x, x' \in T_1$ and $y, y' \in T_2$. If the two edges share an endpoint in T_1, say $x = x'$ and $y \ne y'$, and if t is the third vertex of T_2, then $x\,y\,t\,y'$ is a 4-cycle. If they share an endpoint in T_2, the same construction with the roles of T_1 and T_2 exchanged is a 4-cycle. If they share no endpoint,*

then $x \neq x'$ and $y \neq y'$, both pairs are edges of their triangles, and $x\,y\,y'\,x'$ is a 4-cycle. In particular there is at most one edge with one end in T_1 and one end in T_2.

3. *Suppose T_1 and T_2 are disjoint and lie in the same component. Let d be the distance between their vertex sets, and let $P : u = p_0, p_1, \dots, p_d = v$ be a path of length d with $u \in T_1$ and $v \in T_2$. The internal vertices of P lie outside $T_1 \cup T_2$, and the subgraph induced by $V(T_1) \cup V(T_2) \cup V(P)$ consists of the edges of T_1, the edges of T_2 and the edges of P.*
4. *In the notation of (3), if $x \in T_1 \setminus \{u\}$ and $z \in T_2 \setminus \{v\}$, then $x, u, p_1, \dots, p_{d-1}, v, z$ is an induced path on $d + 3$ vertices.*

Proof. (1) The vertices a, c, d and u are distinct: T_1 and T_2 share only u, so $a \notin T_2$. The edges ac, cd, du and ua are present, the last three because cd and du lie in T_2 and ua lies in T_1. This is the 4-cycle $a\,c\,d\,u$. Each of the other three cross edges is the same configuration after renaming.
(2) First suppose $x = x'$ and $y \neq y'$. Then y and y' are distinct vertices of the triangle T_2, so both are adjacent to the third vertex t of T_2. The vertices x, y, t, y' are distinct because $x \notin T_2$. The edges xy, yt, ty' and $y'x$ are present, and $x\,y\,t\,y'$ is a 4-cycle. The case of a shared endpoint in T_2 is the same argument with the triangles exchanged.
Now suppose the four endpoints are distinct. Any two vertices of a triangle are adjacent, so xx' and yy' are edges. The vertices x, y, y', x' are distinct because the triangles are disjoint. The edges xy, yy', $y'x'$ and $x'x$ are present, and $x\,y\,y'\,x'$ is a 4-cycle. A second edge between T_1 and T_2 falls under one of these two cases, so a strong C_4-free graph has at most one such edge.
(3) A shortest path between the two sets meets each set only at an endpoint, so every internal vertex lies outside $T_1 \cup T_2$. A chord of P would give a shorter u–v path.
Let $x \in T_1 \setminus \{u\}$, and let y be the third vertex of T_1, so $\{u, x, y\}$ is a triangle. Suppose x is adjacent to p_i for some $i \in \{1, \dots, d\}$. If $i = 1$, the vertices y, u, p_1, x are distinct and the edges yu, up_1, p_1x and xy are present, so $y\,u\,p_1\,x$ is a 4-cycle. If $i \geq 2$, the walk $x, p_i, p_{i+1}, \dots, p_d$ has length $d - i + 1 \leq d - 1$ and joins T_1 to T_2, contradicting the definition of d. Thus x has no neighbour on P except u. The same argument, using the third vertex of T_2, shows that no vertex of $T_2 \setminus \{v\}$ has a neighbour on P except v. In particular, adjacency to p_{d-1} produces a 4-cycle, and adjacency to an earlier vertex of P shortens the distance.
It remains to rule out an edge with one end in T_1 and one end in T_2, other than the edge uv in the case $d = 1$. If $d \geq 2$, any such edge would be a path of length 1 between the two sets. If $d = 1$, the path P is the edge uv, and a second edge is a 4-cycle by (2).
No other pair of vertices in $V(T_1) \cup V(T_2) \cup V(P)$ remains. The induced subgraph is therefore the two triangles joined by P.
(4) The named walk has $d + 3$ vertices and uses edges of T_1, of P and of T_2. A chord incident with x would be an edge from x to a vertex of the walk other than u. Part (3) forbids every such edge, including an edge from x to z. A chord incident with z is forbidden in the same way. A chord that avoids x and z is a chord of P or an edge forbidden by (3). The walk is an induced path. □

Lemma 5.9. *Every connected strong C_4-free P_7-free graph with at least two triangles has a vertex v with $L(v) \leq 3$.*

Proof. Let G be connected, strong C_4-free and P_7-free, and suppose G has at least two triangles. Two triangles do not share an edge. Suppose they share the edge uv, and let x and y be the third vertices. The vertices x, u, y, v are distinct, and the edges xu, uy, yv and vx are present, because each of x and y is adjacent to both ends of uv. The cycle $x\,u\,y\,v$ is a 4-cycle. Thus two triangles share at most one vertex.
If two triangles share a vertex, Lemma 5.8(1) forbids an edge between their unshared vertices. Their union is an induced subgraph on five vertices. If the triangles are disjoint, let d be the distance between their vertex sets. Lemma 5.8(3) identifies the induced subgraph formed by the two triangles and a shortest connecting path, and Lemma 5.8(4) supplies an induced path on $d + 3$ vertices. Since $p(G) \leq 6$, one has $d \leq 3$.
Four induced subgraphs therefore exhaust the ways two triangles can meet. They may share a vertex, or they may be joined by a path of length 1, 2 or 3. These graphs have orders 5, 6, 7 and 8. Call them the seeds, and label the vertices of a seed by $0, 1, \dots$ in the order used by Appendix A. From a seed, grow a labelled graph H as follows. Let X be the set of vertices v with $L_H(v) < 4$. A node is unrestricted when X is empty, that is, when every vertex of H has deletion loss at least 4. While X is nonempty, choose the vertex $v \in X$ with the fewest legal children, and if several vertices have that number, choose the smallest label. A legal child is obtained by adding one vertex whose neighbourhood $U \subseteq V(H)$ contains v, provided the enlarged graph has no 4-cycle subgraph and no induced P_7. A 4-cycle through the new vertex exists precisely when two vertices of U have a common neighbour in H. The children are explored in increasing order of the integer bitmask of U. There is no order cutoff and no isomorphism pruning. The four searches visit 36, 31, 46 and 5 nodes, of which 7, 6, 21 and 3 are leaves, and the largest orders they reach are 17, 15, 11 and 10. None of the four searches has an unrestricted node. Every node has X nonempty, and every branch ends at a leaf. Appendix A prints the 118 decisions, one entry per node, and records the induced-P_7 test. An independent C implementation (`referee_checks/lemma57b.c`) written for

cross-verification, run with deficient-vertex decisions only, reproduces the search counts and confirms closure of all four trees, using its own vertex labelling and traversal order.

Suppose G is a connected strong C_4-free P_7-free graph with at least two triangles and with $L \geq 4$ at every vertex. Then G contains one of the four seeds as an induced subgraph. Each search tree is finite and has no unrestricted node, so the branch that follows G ends at a leaf H. At that leaf the chosen vertex v has deletion loss $L_H(v) < 4$ and no legal new neighbour. The loss on an induced subgraph is $L = 2d - 3t$. If v had no neighbour in G outside H, then, because H is induced, the degree and the triangle count of v would already be final, and $L_G(v) = L_H(v) < 4$, contradicting the choice of G. Thus G supplies a neighbour z of v outside H. The neighbourhood $U = N(z) \cap V(H)$ contains v, and the graph formed by H and z is an induced subgraph of G, so it has no 4-cycle subgraph and no induced P_7. It is a legal child of the leaf, which is impossible. Hence no such graph G exists. The same conclusion holds for any selection rule that, whenever X is nonempty, chooses some vertex of X and explores every legal neighbourhood containing it. Fail-first is the rule that produced the arrays in Appendix A. Every connected strong C_4-free P_7-free graph with at least two triangles has a vertex of deletion loss at most 3. □

Theorem 5.10. *Every connected strong C_4-free P_7-free graph G satisfies $S(G) \leq 3n$, with equality if and only if G is the Petersen graph.*

Proof. The triangle-free case, including the equality, is Theorem 5.5. Suppose G contains a triangle and $S(G) \geq 3n$, and choose such a connected graph of minimum order. Every smaller connected graph in the class satisfies $S \leq 3|V|$, and the inequality is strict whenever the smaller graph contains a triangle.

Suppose s is a cut vertex. Induced paths that start at s in distinct components of $G - s$ concatenate. Since $p(G) \leq 6$, some component determines a side G_1, induced by that component together with s, in which every induced path starting at s has order at most three. Let G_2 be the subgraph induced by the remaining vertices together with s, and write $n_i = |V(G_i)|$. Then $n = n_1 + n_2 - 1$ and $n_1 \geq 2$. Lemma 5.3 gives $S(G_1) \leq 2(n_1 - 1)$. The graph G_2 is a smaller connected graph in the class. If it contains a triangle, minimality gives $S(G_2) < 3n_2$; if it does not, Theorem 5.5 gives $S(G_2) \leq 3n_2$. Lemma 2.3 therefore yields

$$S(G) \leq 2(n_1 - 1) + 3n_2 = 3n - (n_1 - 1) < 3n.$$

Thus G has no cut vertex, and $G - v$ is connected for every vertex v.

Suppose $L(v) \leq 3$ for some vertex v. Then $S(G - v) = S(G) - L(v) \geq 3n - 3 = 3(n - 1)$. The graph $G - v$ is connected and lies in the class. It contains no triangle: otherwise minimality would give the strict inequality $S(G - v) < 3(n - 1)$. Theorem 5.5 therefore forces $G - v$ to be the Petersen graph, so $n = 11$ and $S(G - v) = 30$. Any two nonadjacent vertices of the Petersen graph have a common neighbour, so the neighbours of v form a clique. The Petersen graph is triangle-free, and a clique in it has size at most two. The graph G contains a triangle, so v is adjacent to both ends of an edge and to no further vertex of $G - v$. Then $d(v) = 2$, $t(v) = 1$ and $L(v) = 2 \cdot 2 - 3 \cdot 1 = 1$, hence $S(G) = 31 < 33$. This contradiction shows that $L(v) \geq 4$ for every vertex v.

Thus $L(v) \geq 4$ at every vertex. If $T(G) \geq 2$, Lemma 5.9 supplies a vertex of deletion loss at most 3, a contradiction. Hence $T(G) = 1$. Theorem 5.7 gives $S(G) \leq 3n$, so $S(G) = 3n$.

Let v be a vertex of minimum degree on the unique triangle, write $d = d(v)$, and let u and w be the other two vertices. The graph $H = G - v$ is connected and triangle-free. Let Q be its core under Lemma 5.4, of order h and deficit $\eta = \eta(Q)$, and define a and c as in the proof of Theorem 5.7. That proof gives the identity (5.1) and the bounds $a \leq \eta + 3$ and (5.2). The same case division yields the sharper bound $a \leq \eta + 2$. If Q is the Petersen graph then $a \leq 2 = \eta + 2$. For each remaining fixed graph, $a \leq \lfloor h/2 \rfloor \leq 5$, while the only rows of Lemma 5.2 with $\eta < 4$ are F_4 and F_{12}, where the descriptions give $a \leq 4$ and $a \leq 5$ respectively. Every other fixed row has $\eta \geq 4$, so $\eta + 2 \geq 6$. In each parameterised family $a \leq 4$ and $\eta \in \{5, 6\}$. The graph K_1 has $a \leq 1$ and $\eta = 3$, and K_2 has $a \leq 2$ and $\eta = 4$.

If u or w lies outside Q, the estimate in the proof of Theorem 5.7 becomes $S(G) - 3n \leq a - \eta - 3 \leq -1$. Thus both u and w lie in Q. The same proof then gives $d \leq 1 + c + \min\{d_Q(u), d_Q(w)\}$, and

$$S(G) - 3n \leq a + \min\{d_Q(u), d_Q(w)\} - \eta - 5.$$

Equality in (5.1) forces both

$$d = 1 + c + \min\{d_Q(u), d_Q(w)\} \qquad \text{and} \qquad a + \min\{d_Q(u), d_Q(w)\} = \eta + 5. \tag{5.3}$$

No core other than the Petersen graph meets (5.3). In the parameterised families every edge has an end of degree two and $a \leq 4$, so the left-hand sum is at most 6, while $\eta + 5 \geq 10$. On K_2 the sum is at most 3. On C_5, C_6 and C_7 every degree is 2 and $\lfloor h/2 \rfloor + 2 \leq 5 < \eta + 5$. The graph F_4 has maximum degree three and $a \leq 4$, so the sum is at most $7 < 8$. On F_3 the three added vertices induce a path and meet three distinct cycle vertices, so the maximum degree is three and, with $a \leq 4$,

the sum is at most $7 < 9$. On F_7 and on F_8 each added vertex meets the 7-cycle in two vertices, the traces are disjoint, and the maximum degree is three; the resulting sums are at most 7, against $\eta + 5 \in \{11, 10\}$. On F_{10} the only vertex of degree four is v_1, each of whose four neighbours has degree two, so every edge has an end of degree at most three and, with $a \le 4$, the sum is at most $7 < 10$. On F_9, obtained from F_8 by a vertex adjacent only to x_1 and x_2, the maximum degree is three and the sum is at most $8 < 9$. On F_{11} the vertices of degree four are v_1 and v_5, which are nonadjacent on the 7-cycle, and every neighbour of either has degree at most three, so every edge has an end of degree at most three and the sum is at most $8 < 9$. On F_{12} every edge has an end of degree at most three and $a \le 5$, so equality would require $a = 5$ and $\min\{d_Q(u), d_Q(w)\} = 3$. The five vertices of $N(v) \cap V(Q)$ have pairwise disjoint neighbourhoods in a graph of order 11 and minimum degree two. The edge uw contributes degrees adding to at least $3 + 3$, and the other three vertices contribute at least 2 each, so the sum of the five degrees is at least $12 > 11$.

Thus Q is the Petersen graph, $\eta = 0$, $a = 2$ and $\min\{d_Q(u), d_Q(w)\} = 3$. Every degree in Q is three, so $d_Q(u) = d_Q(w) = 3$. Write c_u for the number of neighbours of u in $H - V(Q)$, and c_w likewise. A common neighbour would form a triangle with the edge uw, so these neighbourhoods are disjoint and $c_u + c_w \le c$. As in the proof of Theorem 5.7, $d(u) \le 1 + d_Q(u) + c_u$ and $d(w) \le 1 + d_Q(w) + c_w$. Since d is at most both $d(u)$ and $d(w)$, the equality $d = 4 + c$ forces $c \le c_u$ and $c \le c_w$. Combined with $c_u + c_w \le c$ this gives $c = 0$, and then $d = 4$. The integer $t = d - a$ equals 2, so H consists of Q and two further vertices, both adjacent to v.

Those two vertices are leaves of H, attached at distinct vertices q_1 and q_2 of Q. A path of length two in an attached tree would place both of its vertices in $N(v)$, and the edge between them would form a second triangle with v. The supports are distinct, for a common support would give that vertex and v two common neighbours. Neither support is u or w, because $c_u = c_w = 0$. Neither support is adjacent to u or to w: the edge from u to q_1, together with the leaf at q_1 and the edge uv, would be a 4-cycle.

If q_1 and q_2 are nonadjacent, the proof of Theorem 5.6 supplies an induced P_5 of Q with those ends. The two leaves are nonadjacent, since both are adjacent to v and the triangle is unique, and each leaf meets Q only at its support. Adding the leaves to the P_5 produces an induced P_7.

If q_1 and q_2 are adjacent, realise Q on the 2-element subsets of $\{1, 2, 3, 4, 5\}$, adjacent when disjoint, as in the proof of Theorem 5.6. The symmetric group on the ground set acts by graph automorphisms and is edge-transitive, so it carries the edge uw to $\{\{1, 2\}, \{3, 4\}\}$. The neighbours of $\{1, 2\}$ are the three pairs in $\{3, 4, 5\}$, and the neighbours of $\{3, 4\}$ are $\{1, 2\}$, $\{1, 5\}$ and $\{2, 5\}$. The four remaining vertices are $\{1, 3\}$, $\{1, 4\}$, $\{2, 3\}$ and $\{2, 4\}$, and the only edges among them are $\{1, 3\}\{2, 4\}$ and $\{1, 4\}\{2, 3\}$. The transposition (1 2) fixes uw and exchanges those two edges, so in one such labelling $\{q_1, q_2\} = \{\{1, 3\}, \{2, 4\}\}$. Let x be the leaf at $\{1, 3\}$. The path

$$v,\ x,\ \{1, 3\},\ \{2, 4\},\ \{1, 5\},\ \{2, 3\},\ \{1, 4\}$$

is an induced P_7. Consecutive vertices are adjacent by the construction of G or by disjointness. The vertex v meets Q only in $\{1, 2\}$ and $\{3, 4\}$, the leaf x meets Q only in $\{1, 3\}$, and each of $\{1, 3\}$, $\{2, 4\}$, $\{1, 5\}$ and $\{2, 3\}$ meets every later nonconsecutive vertex of the path in a pair that shares an element of $\{1, 2, 3, 4, 5\}$.

In every case G contains an induced P_7. This contradiction shows that no connected graph in the class with a triangle satisfies $S \ge 3n$. Combined with Theorem 5.5, one has $S \le 3n$, with equality only for the Petersen graph. □

Theorem 5.11. *Let G be a connected graph on at least two vertices with $\ell(G) \le 4$. Then*

$$p(G) \ge r(G) + \ell(G)^{c_{C_4}(G)}.$$

If $c_{C_4}(G) = 0$, the power equals 1. The conjecture remains open for graphs with $\ell > 4$.

Proof. If G contains a 4-cycle, then $c_{C_4}(G) = 0$ and the inequality is $p(G) \ge r(G) + 1$. A shortest path between two vertices at distance $\operatorname{diam}(G) \ge r(G)$ is induced.

Suppose now that G is strong C_4-free, and write $r = r(G)$. The inequality is $p \ge r + \ell$.

- $r = 1$. Lemma 3.1 gives $\ell < 2$ and $p \ge r + \lceil \ell \rceil$, hence $p \ge r + \ell$.
- $r = 2$. If $\ell \le 2$, Lemma 3.2 gives $p \ge 4$. If $2 < \ell \le 3$ and $p \le 4$, Lemma 3.3 gives $\ell \le 2$, a contradiction, so $p \ge 5$. If $3 < \ell \le 4$ and $p \le 5$, Theorem 3.6 gives $\ell \le 3$, a contradiction, and hence $p \ge 6$.
- $r = 3$. If $\ell \le 2$, Theorem 2.4 gives $p \ge 5$. If $2 < \ell \le 3$ and $p \le 5$, then Lemma 3.5 and $r = 3$ give $G = C_6$, with $\ell = 2$, so $p \ge 6$. If $3 < \ell \le 4$ and $p \le 5$, Theorem 3.6 gives $\ell \le 3$, a contradiction. If $3 < \ell \le 4$ and $p = 6$, Theorem 5.10 gives $\ell \le 3$, a contradiction. Hence $p \ge 7 \ge r + \ell$.
- $r = 4$. Theorem 2.4 gives $p \ge 7$. If $p = 7$, Corollary 4.4 gives $\ell \le 2$, so $p \ge r + \ell$. If $p \ge 8$ and $\ell \le 4$, then $p \ge 8 = r + 4 \ge r + \ell$.
- $r \ge 5$. Theorem 2.4 gives $p \ge 2r - 1 \ge r + 4 \ge r + \ell$.

□

Since $\lambda(v) \le d(v)$ for every vertex, $\ell \le 2e/n$. Theorem 5.11 covers every connected graph of average degree at most 4. Any two vertices of a strong C_4-free graph have at most one common neighbour, so

$$\sum_v \binom{d(v)}{2} \le \binom{n}{2}.$$

Convexity yields $(2e/n)(2e/n-1) \le n-1$, hence

$$\frac{2e}{n} \le \frac{1+\sqrt{4n-3}}{2}.$$

This is Reiman's bound [14]; the Kővári–Sós–Turán theorem gives the same order of magnitude for subgraphs $K_{2,2}$ [11]. Since $S = 2e - 3T$, one has $\ell \le 2e/n$. For $n \le 13$ one has $\sqrt{4n-3} \le 7$, so $\ell \le 4$. Every connected strong C_4-free graph on at most 13 vertices therefore falls under Theorem 5.11. Every connected graph on at most 13 vertices that contains a 4-cycle satisfies Conjecture 1.1 because the exponent is 0. The theorem covers every connected graph of order at most 13. Equality in the real-valued inequality holds for K_2 and K_3 ($r = 1$, $\ell = 1$, $p = 2$), for C_5 ($r = 2$, $\ell = 2$, $p = 4$), for the Petersen graph ($r = 2$, $\ell = 3$, $p = 5$) and for C_6 ($r = 3$, $\ell = 2$, $p = 5$). The article does not characterise the equality cases in general.

Remark 5.12. Every connected graph G with $\ell(G) < 5$ satisfies

$$p(G) \ge r(G) + \left\lfloor \ell(G)^{c_{C_4}(G)} \right\rfloor.$$

The exponent follows the convention of Conjecture 1.1: if $c_{C_4}(G) = 0$, then the power equals 1. If G contains a 4-cycle, both the real form and this floor form reduce to $p \ge r + 1$, which follows from a shortest path between two vertices at distance $\operatorname{diam}(G) \ge r(G)$. If G is strong C_4-free and $\ell \le 4$, Theorem 5.11 gives $p \ge r + \ell$, and $\ell \ge \lfloor \ell \rfloor$. Suppose G is strong C_4-free and $4 \le \ell < 5$, and suppose the floor form fails. Then $\lfloor \ell \rfloor = 4$, so $p \le r + 3$. Theorem 2.4 gives $p \ge 2r - 1$, so $r \le 4$. If $r \le 3$, then $p \le 6$, and Theorem 5.10 gives $\ell \le 3$, contradicting $\ell \ge 4$. If $r = 4$, then $p = 7$, and Corollary 4.4 gives $\ell \le 2$, again a contradiction. The real-valued conjecture for $\ell > 4$ remains open.

6 | Concluding remarks and open problems

For integers $K \ge 3$ and $r \ge 4$, write GEN_K for the statement that every connected strong C_4-free graph G with $p(G) \le 2K$ satisfies $S(G) \le Kn$, and write $\mathrm{RAD}(r)$ for the statement that every connected strong C_4-free graph with $r(G) = r$ and $p(G) = 2r - 1$ satisfies $\ell(G) \le r - 1$. Corollary 4.4 is the case $r = 4$ of $\mathrm{RAD}(r)$, in the stronger form $\ell \le 2$.

Corollary 6.1. *Let G be a connected strong C_4-free graph that violates Conjecture 1.1. Then $\ell(G) > 4$. Put $K = \lceil \ell(G) \rceil - 1 \ge 4$. Then either*

(a) $r(G) \le K$ and $p(G) \le r(G) + K \le 2K$, so G violates GEN_K; or
(b) $r(G) = K + 1 \ge 5$ and $p(G) = 2r(G) - 1$, and $\ell(G) > r(G) - 1$, so G violates $\mathrm{RAD}(K+1)$.

Proof. Write $r = r(G)$, $p = p(G)$ and $c = \lceil \ell(G) \rceil = K + 1$. Then $\ell > 4$ by Theorem 5.11, so $K \ge 4$. A violation means $p \le r + c - 1 = r + K$. Theorem 2.4 gives $2r - 1 \le p \le r + K$, so $r \le K + 1$. If $r \le K$, then $p \le r + K \le 2K$, and $S > Kn$ because $\ell > K$. If $r = K + 1$, then $2r - 1 \le p \le 2K + 1 = 2r - 1$, and $\ell > K = r - 1$. The radius in (b) is at least 5. □

Proposition 6.2. *For $k \ge 3$ let θ_k consist of two vertices x, y joined by k internally disjoint paths xa_ib_iy. This is the graph F_0' of Section 5 with parameter $t' = k - 2$. Then θ_k is strong C_4-free, $n = 2k + 2$, $r = 3$, $p = 6$ and $S = 6k = 3n - 6$. Thus $p - r - \ell = 6/n \to 0$.*

Proof. Every cycle consists of two complete branches and so has length 6; in particular there is no 4-cycle. There are no triangles, so $S = 2e = 6k$. Every vertex has eccentricity 3: $d(x, y) = 3$ and $d(a_i, b_j) = 3$ for $i \ne j$. The path $a_jxa_ib_iyb_l$ (with j, i, l distinct) is induced. Conversely, let P be an induced path. If P avoids x and y, it lies inside one edge a_ib_i. If P contains x but not y, then x has at most two neighbours on P, each followed on P by at most its b-vertex, so $|P| \le 5$. If P contains x and y, then every a_j on P is adjacent on P to x and every b_j to y, at most one branch lies entirely on P, and so P consists of x, y, at most one full branch and at most one further neighbour of each of x and y. Hence $|P| \le 6$. □

Corollary 6.3. *Let G be a connected strong C_4-free graph with $p(G) = 6$. Then $S(G) \le 3n - 1$ and $\ell(G) < 3$. On each of the classes $(r, p) = (2, 6)$ and $(r, p) = (3, 6)$ the supremum of ℓ is 3, and the supremum is not attained.*

Proof. The graph is P_7-free. Theorem 5.10 gives $S(G) \le 3n$, with equality only for the Petersen graph. The Petersen graph has induced-path order 5, so $S(G) \le 3n-1$ and $\ell(G) < 3$.
Both constructions below are strictly less than 3 and tend to 3 as the order tends to infinity. The supremum on each class is therefore 3, and it is not attained.
For the class $(2,6)$, take the graph F_2 defined before Lemma 5.2, with parameter $t \ge 2$. In the description preceding Lemma 5.2 the hubs are v_1 and v_3, the unique branch of length 2 is $v_1v_2v_3$, and the remaining v_1–v_3 paths are the t matching paths through $A_1 \cup A_3$ together with $v_1v_5v_4v_3$. These $t+1 \ge 3$ paths are branches of length 3. The order is $n = 5+2t$. The graph is triangle-free, and the table of Lemma 5.2 gives $2e = 3n-5$, so $\ell = 3-5/n$. Every vertex other than the two hubs is adjacent to a hub, and v_2 is adjacent to both hubs, so every vertex is within distance 2 of v_2. A vertex of A_1 is not adjacent to v_2. Each hub has degree $t+2 = (n-1)/2 < n-1$, and every other vertex has degree 2, so no vertex dominates the graph. The radius is 2. The proof of Theorem 5.6 shows that every induced path has order at most 6. For the matching lower bound, choose three branches of length 3, written v_1abv_3, v_1cdv_3 and v_1efv_3. The path a, v_1, c, d, v_3, f is induced: the hubs are nonadjacent, and an internal vertex of any one of these branches is adjacent only to its two neighbours on that branch. Thus $(r, p, \ell) = (2, 6, 3-5/n)$.
For the class $(3,6)$, Proposition 6.2 gives the graph θ_k for every $k \ge 3$, with $\ell = 3-6/n$. □

The two families of Corollary 6.3 are why a uniform gap below 3 is not available.
Theorem 5.11 stops at $\ell = 4$ because that is the largest constant for which the available gap $p-r$ meets ℓ. Theorem 5.10 gives $\ell \le 3$ whenever $p \le 6$. Corollary 4.4 gives $\ell \le 2$ when $r = 4$ and $p = 7$. For $r \ge 5$, Theorem 2.4 gives $p \ge 2r-1$, hence a gap of at least 4, which meets $\ell \le 4$ and no larger constant. A graph with $\ell > 4$ has $p \ge 7$. The bound for induced-path order at most 6 does not apply to it, and if its radius is at least 5 the radius-four classification does not apply either. The obstruction to a larger constant is the lack of an analogue of Theorem 5.10, such as $S \le 4n$ for every connected strong C_4-free P_9-free graph. No such bound is proved here.
Corollary 6.1 splits any counterexample to REAL-133 into two open statements: either $S > Kn$ for some $K \ge 4$ on a connected strong C_4-free graph with $p \le 2K$, or a connected strong C_4-free graph of radius $r \ge 5$ and induced-path order $2r-1$ with $\ell > r-1$. The graphs F_2 and θ_k of Corollary 6.3 have induced-path order 6 and $\ell = 3-5/n$ and $\ell = 3-6/n$ respectively, so there is no uniform gap below 3 from which a longer induced path would inherit $S \le 3n$. The examples that follow limit extensions which do not use a new bound of GEN_K type.

Conjecture 6.4. *For every integer $n \ge 10$, the maximum of $S(G)$ over connected strong C_4-free P_7-free graphs of order n equals $f(n)$ from Theorem 5.6. Every graph that attains the maximum is one of the extremal graphs of that theorem, and in particular is triangle-free.*

Conjecture 6.4 is open. It would unify Theorems 5.6 and 5.10: the triangle-free extremal graphs would be extremal in the whole class.

Example 6.5. Let $s \ge 3$, $t \ge 3$ and $m \ge 1$ be integers. Let $G_{s,t,m}$ be the cycle $C_{2s} = c_0 \cdots c_{2s-1}$ with a pendant path of t edges at every cycle vertex, together with m internally disjoint paths c_0xzyc_1. Then $n = 2s(t+1)+3m$, $e = 2s(t+1)+4m$ and $T = 0$, so $e-n-T = m \ge 1$. For $(s,t,m) = (3,3,1)$ one has $n = 27$, $e = 28$, $r = 6$ and $p = 11$. These graphs satisfy REAL-133, since $\ell = 2e/n < 8/3$. The identity $e = n+T$ holds for every graph of Theorem 4.2. The count in the proof of Corollary 4.4 gives $n = 6+a+b$, $e = 6+a+2b+m$ and $T = b+m$ in the construction, hence $e = n+T$, and C_8 has $e = n$ and $T = 0$. The graphs $G_{s,t,m}$ show that the identity fails for strong C_4-free graphs of induced-path order $2r-1$ at every radius $r \ge 6$: here $r(G_{s,t,m}) = s+t$ and $p(G_{s,t,m}) = 2(s+t)-1$.
Two vertices have at most one common neighbour, so G is strong C_4-free and triangle-free. An added path of length four does not shorten distances among the older vertices, and c_0 remains within distance two of every vertex of such a path, so the eccentricity of c_0 is still $s+t$. A vertex w of an added path meets the tip of the arm at c_s only through c_0 or c_1, hence at distance at least $s+t$. Thus $r = s+t$. For the path order, let B be the block formed by C and the added paths. An induced path P meets B in a subpath Q and can leave B only at an end of Q that is a cycle vertex, through at most t arm vertices. If Q avoids both c_0 and c_1, then $|Q| \le 2s-2$. If Q contains exactly one of them, then either Q lies on the cycle with $|Q| \le 2s-1$, or $|Q| \le 2s+2$ with at most one cycle end, or $|Q| \le 7$ with no cycle end. If Q contains both, the same three shapes occur with $|Q| \le 2s-1$, $|Q| \le 2s+2$ and $|Q| \le 6$. Adding at most t arm vertices at each cycle end of Q gives $|P| \le \max\{2s-1+2t,\ 2s+2+t,\ 7\}$. For $s \ge 3$ and $t \ge 3$ this maximum equals $2s+2t-1$, and the cycle minus one vertex together with the arms at its two ends attains that order.

Example 6.6. A uniform lower bound on the deletion loss does not bound λ. Two copies of the Petersen graph sharing a vertex form a triangle-free strong C_4-free graph on $n = 19$ vertices with $S = 60$, minimum deletion loss 6, induced-path order 9 and $\lambda = 6$ at the shared vertex. For $b \ge 2$, gluing b copies of the Petersen graph along a common edge gives a 2-connected graph with $n = 8b+2$, $S = 28b+2$, minimum loss 6, induced-path order 9 and $\lambda = 2b+1$ at the two hubs.

Example 6.7. Let q be a prime power, and let F be the incidence graph of the projective plane $\mathrm{PG}(2,q)$, with $f = |F| = 2(q^2+q+1)$ and $S(F) = (q+1)f$. Choose a root of F, set $t = f-1$ and $b \geq 1$, and take a 6-cycle $c_0 \cdots c_5$. Attach a pendant path with t edges at each cycle vertex, and attach b copies of F at c_0 by identifying their roots with c_0. Each new block meets the rest of the graph in a cut vertex, so the result is strong C_4-free, and

$$n = 6f + b(f-1), \qquad S = 12f + b(q+1)f, \qquad r = t+3, \qquad p = 2t+5 = 2r-1.$$

Distances inside the flower are unchanged. Every vertex of an attached copy is within distance $f-1 \leq t$ of c_0, so each cycle vertex has eccentricity $t+3$, while a vertex x of a copy has distance $d_F(x,c_0)+t+3$ from the tip of the arm at c_3. An induced path meets the cycle in at most five consecutive vertices, and each of its two ends outside the cycle contributes at most t vertices. The sum S is additive at cut vertices by Lemma 2.3. For fixed q, as $b \to \infty$ one has $\ell \to (q+1)f/(f-1)$. Letting $q \to \infty$, with $r = f+2$, this limit grows like $\sqrt{r/2}$. For $q = 5$ and $b = 17$ one has $n = 1409$, $r = 64$, $p = 127$ and $S = 7068 > 5n$. Thus ℓ is unbounded on the class of strong C_4-free graphs with $p = 2r-1$, and any statement $\mathrm{RAD}(r)$ has to use a bound that depends on r.

Example 6.8. Let P be the Petersen graph, and let u and v be the ends of an induced P_5 in P. Such a path exists by the proof of Theorem 5.6, and u and v are nonadjacent. Form G by adding one leaf at u and one leaf at v. Then

$$\big(r(G), p(G), |V(G)|, S(G), \ell(G)\big) = (2, 7, 12, 34, 17/6).$$

The graph P has 15 edges. The two new edges create no cycle, so G is triangle-free and strong C_4-free, $n = 12$, $e = 17$ and $S = 2e = 34$. The graph P is P_6-free by Theorem 3.6. A leaf has degree one, so it is an end of every induced path that contains it, and deleting the leaves that lie on an induced path of G leaves an induced path of P. The path formed by the two leaves and the chosen P_5 is induced, because each leaf is adjacent only to its endpoint. Hence $p(G) = 7$. The diameter of P is 2, so u and v have a common neighbour w. Every vertex of P is at distance at most 2 from w, and each new leaf is at distance 2 from w. Thus the radius is 2. The example shows that $r = 2$ and $p = 7$ occur together with $\ell = 17/6$.

APPENDIX

A | The two-triangle certificate

Lemma 5.9 grows four labelled graphs, one from each way two triangles can meet. The authors' checker is `certificates/gen3/verify_gen3.py`, with SHA-256 digest `f1b67189 41f9cf3e 6b0917cf 25ab9093 810b07f6 d35a11fb 7cc539fb 546e6fd0`. The program uses only the Python standard library, requires Python 3.10 or later, embeds the decision certificate and takes no arguments. From the repository root the command is `python3 certificates/gen3/verify_gen3.py`. A run of that command replays the embedded certificate. It prints

```
triangle distance 0    nodes= 36  maximum order=17  leaves= 7  unrestricted=0
triangle distance 1    nodes= 31  maximum order=15  leaves= 6  unrestricted=0
triangle distance 2    nodes= 46  maximum order=11  leaves=21  unrestricted=0
triangle distance 3    nodes=  5  maximum order=10  leaves= 3  unrestricted=0
PASS. All branches and all decision entries verified.
```

The four rows are the searches of Lemma 5.9, in the order shared vertex, one joining edge, and joining paths of lengths two and three. Together they visit 118 nodes and 37 leaves. As in the proof of that lemma, a node is unrestricted when every vertex of the current graph has deletion loss at least 4. The unrestricted count on each of these four rows is zero, so every branch ends at a leaf.

The decision at each node of a distance row is the vertex of deletion loss less than 4 with the fewest legal children. Ties are broken by the smallest label, and the children of a node are visited in increasing order of the neighbourhood bitmask. These two rules determine the arrays below. They are written in depth-first preorder, one integer per node, and the integer is the label of the chosen vertex. An independent C implementation (`referee_checks/lemma57b.c`) written for cross-verification is archived in this repository. With the environment variable `DEFONLY=1` it uses only deficient-vertex decisions and the same fail-first rule. On the seeds `D{c`, `E{CW`, `FwC[_` and `GwC[?c` the second implementation reproduces the search counts and confirms closure of all four trees, using its own vertex labelling and traversal order. Replaying `verify_gen3.py` is a check of the authors' certificate, not a second implementation.

The arrays, in the order of the four distance rows, are as follows.

```
0: 0, 1, 2, 1, 3, 4, 2, 3, 4, 6, 9, 9,
   6, 9, 9, 7, 7, 4, 10, 6, 6, 3, 9,
   6, 6, 2, 7, 9, 9, 1, 3, 4, 6, 8, 10, 10
```

```
1: 0, 3, 1, 2, 4, 5, 8, 8, 2, 1, 2, 1,
   2, 1, 3, 7, 1, 2, 4, 5, 10, 10, 0, 3,
   6, 1, 2, 4, 5, 10, 10
2: 0, 1, 2, 5, 4, 4, 5, 2, 5, 1, 5, 2,
   1, 2, 1, 1, 2, 5, 4, 7, 1, 2, 4, 2,
   1, 6, 6, 2, 6, 1, 6, 2, 1, 1, 2, 4,
   2, 1, 5, 5, 2, 1, 2, 5, 1, 5
3: 1, 6, 1, 6, 1
```

The nodes by order, and the orders of the leaves, are as follows. Shared vertex: 1, 1, 1, 1, 2, 2, 2, 3, 4, 5, 7, 5, 2 nodes on orders 5 through 17, and 2, 3, 2 leaves on orders 15, 16, 17. One edge: 1, 2, 3, 6, 6, 3, 3, 3, 2, 2 nodes on orders 6 through 15, and 3, 1, 2 leaves on orders 10, 13, 15. Path of length two: 1, 4, 14, 19, 8 nodes on orders 7 through 11, and 1, 12, 8 leaves on orders 9, 10, 11. Path of length three: 1, 3, 1 nodes on orders 8, 9, 10, and 2, 1 leaves on orders 9, 10. The distance-3 seed already spans an induced path on six vertices, by Lemma 5.8(4), and the search dies by order 10. The distance-2 search is the broadest, with 21 leaves, and it dies by order 11. The other two searches reach orders 17 and 15 and then close. No node in any of the four trees is free of a vertex of deletion loss less than 4. This is a summary of the closed branches, not a hand proof that each leaf is illegal.

At each node the verifier names the vertex just described and explores every legal neighbourhood that contains it. A child is rejected when two vertices of the new neighbourhood have a common neighbour in the graph already built. It is rejected when the new vertex and six existing vertices induce a P_7. The induced-P_7 test is the following. For a set S of six vertices of the graph already built and a new vertex z, the set $S \cup \{z\}$ induces a P_7 precisely when $G[S]$ is a disjoint union of one or two paths and the neighbourhood of z on S consists of exactly one endpoint of each path component and of no other vertex of S. An isolated vertex is a path of order one and is its own end.

```
for each 6-set S:
    if some vertex of S has more than two neighbours in S: skip
    if the number of edges induced by S is not 4 or 5: skip
    split S into its components
    if a component has no end, or the number of components
        is not 6 minus the number of edges: skip
    reject each neighbourhood N for which N cap S consists
        of exactly one endpoint of each path component
        and of no other vertex of S
```

The reason each finite tree rules out a counterexample is the argument in the proof of Lemma 5.9.

Initial configuration	Order	Nodes	Largest order	Leaves	Unrestricted
Shared vertex	5	36	17	7	0
One joining edge	6	31	15	6	0
Joining path of length two	7	46	11	21	0
Joining path of length three	8	5	10	3	0

DATA AVAILABILITY

The source of this article is https://github.com/neardws/induced-paths-without-4-cycles. The authors' certificate for Lemma 5.9 is `certificates/gen3/verify_gen3.py`. A separate implementation is provided in `referee_checks/` and was used for computational cross-checking. The map from claims of this article to scripts is in `referee_checks/README.md`. The Lean development formalizes the main theorems and selected supporting statements. The precise coverage is listed in `lean/README.md`. Its numbered inventory has no remaining `sorry` or `admit`. Five literature axioms remain, recorded in `lean/AXIOMS.md`. That development is not a verification of this article from an empty axiom set.

FORMALIZATION NOTE

The directory `lean/` is a Lean 4 and Mathlib development. It formalizes the main theorems and selected supporting statements; the precise coverage is listed in `lean/README.md`. The files are at

https://github.com/neardws/induced-paths-without-4-cycles/tree/main/lean.

The numbered inventory has no remaining `sorry` or `admit`. Five literature axioms remain, recorded in `lean/AXIOMS.md`. Finite certificates in that development are closed by `native_decide`, which trusts the Lean compiler and its runtime. This is not a verification of the article from an empty axiom set.

ACKNOWLEDGMENTS

Xingchen Wang established the early foundations of this work. Xincao Xu led the later research direction and the organization of the results. During the 2026 revision the authors used GPT-6 Astra and GPT-6 Pro to assist with calculations and with drafting arguments, and Grok Bot to assist with drafting and revising the manuscript. These tools were not a verification of the proofs. The authors reviewed and revised all of the resulting text. They remain fully responsible for the accuracy and integrity of the work.

References

1. A. Brandstädt and C. T. Hoàng. *On clique separators, nearly chordal graphs, and the Maximum Weight Stable Set Problem*. Theoret. Comput. Sci. **389** (2007), 295–306.
2. E. Camby and O. Schaudt. *A new characterization of P_k-free graphs*. Algorithmica **75** (2016), 205–217. Theorem 5 of arXiv:1402.7213 is the equivalence used in the text.
3. H. Chen. *Partial results on the WOWII-133 Graffiti conjecture*. Zenodo. 2026. https://doi.org/10.5281/zenodo.22054835.
4. R. Chen, D. Wu, and B. Xu. *Structure of some (P_7, C_4)-free graphs with application to colorings*. Discrete Appl. Math. **357** (2024), 14–23. doi:10.1016/j.dam.2024.05.038. The arXiv version, arXiv:2302.11115, is titled "Structure and coloring of some (P_7, C_4)-free graphs".
5. R. Chen and B. Xu. *Nearly optimal coloring of some C_4-free graphs*. Discrete Appl. Math. **394** (2026), 145–158. https://doi.org/10.1016/j.dam.2026.07.010; arXiv:2409.06944.
6. E. DeLaVina. *Written on the Wall II (Conjectures of Graffiti.pc)*. http://cms.dt.uh.edu/faculty/delavinae/research/wowII/. 2005. Conjecture 133, dated July 12, 2005.
7. P. Erdős, M. Saks, and V. T. Sós. *Maximum induced trees in graphs*. J. Combin. Theory Ser. B **41** (1986), 61–79.
8. Formal Conjectures. *Graph Conjecture 133*. Google DeepMind, https://github.com/google-deepmind/formal-conjectures/blob/5faef0c4dc3d226f8f9a9fc8f581b6221283eb6e/FormalConjectures/WrittenOnTheWallII/GraphConjecture133.lean. Secondary record of the real-valued reading. The primary source is DeLaVina, Written on the Wall II, Conjecture 133, July 12, 2005.
9. F. Galvin, I. Rival, and B. Sands. *A Ramsey-type theorem for traceable graphs*. J. Combin. Theory Ser. B **33** (1982), 7–16.
10. T. Karthick and F. Maffray. *Square-free graphs with no six-vertex induced path*. SIAM J. Discrete Math. **33** (2019), 874–909. arXiv:1805.05007.
11. T. Kővári, V. T. Sós, and P. Turán. *On a problem of K. Zarankiewicz*. Colloq. Math. **3** (1954), 50–57.
12. D. Kühn and D. Osthus. *Induced subdivisions in $K_{s,s}$-free graphs of large average degree*. Combinatorica **24** (2004), 287–304.
13. P.-S. Loh, M. Tait, C. Timmons, and R. M. Zhou. *Induced Turán numbers*. Combin. Probab. Comput. **27** (2018), 274–288.
14. I. Reiman. *Über ein Problem von K. Zarankiewicz*. Acta Math. Acad. Sci. Hungar. **9** (1958), 269–273.
15. D. Seinsche. *On a property of the class of n-colorable graphs*. J. Combin. Theory Ser. B **16** (1974), 191–193.