SUPPLEMENT TO INDUCED PATHS, RADIUS, AND EXTREMAL NEIGHBORHOOD INDEPENDENCE IN GRAPHS WITHOUT 4-CYCLES

XINGCHEN WANG AND NEIL XU

Abstract. This supplement contains the rooted bounds, the endpoint obstructions, the high-radius constructions, and the conditional reduction for GEN_3 removed from the article. Numbered references to results proved in the article point to that file. The article's Theorems on $M(n)$, radius four, the exact triangle-free function $f(n)$, and graphs with at most one triangle do not depend on these arguments.

1. Rooted bounds of orders three and four

The arguments of this section and of Sections 2 and 3 are not used for Theorems 3.6, 4.3, 5.4, 5.5 and 5.6 or for Corollary 5.11. The rooted bound that bears on the next open band is Theorem 1.2: it is the estimate which shows that a minimum counterexample to GEN_4 is 2-connected. The distance-layer form of the same estimate is omitted. The block-class bound and the Moore observation are kept because they delimit GEN_K, and Example 3.4 is kept because it is the explicit failure of the endpoint inequalities at order nine.

For a vertex s of a connected graph F let $p_s(F)$ be the largest order of an induced path starting at s. Write $D = d_F(s)$ and $m = e(F[N(s)])$.

Definition 1.1. For an integer $q \geq 3$, the rooted bound $\mathrm{ROOT}(q)$ asserts that every connected strong C_4-free graph F and every vertex s with $p_s(F) \leq q$ satisfy

$$S(F) \leq (q-1)(|F|-1) - (q-3)D - m.$$

The hypothesis $p_s(F) \leq q$ belongs in the statement. Without it the displayed inequality is false: if $q = 3$ and $F = C_5$, then $S(F) = 10$ and $m = 0$, while $2(|F|-1) = 8$. Lemma 5.3, already proved because the cut-vertex induction of Theorem 5.4 uses it, is the case $q = 3$.

Theorem 1.2. *Let F be connected and strong C_4-free, and let $s \in V(F)$ satisfy $p_s(F) \leq 4$. Write $D = d(s)$ and $m = e(F[N(s)])$. Then*

$$S(F) \leq 3(n-1) - D - m. \tag{1}$$


In particular $\mathrm{ROOT}(4)$ *holds. Equality holds for C_5 at any root, and for a graph formed by $k \geq 1$ copies of C_5 that meet only at the root.*

Proof. The proof is by induction on n. If $p_s(F) \leq 3$, Lemma 5.3 gives $S \leq 2(n-1) - m$. Since $D \leq n-1$, the right-hand side is at most $3(n-1) - D - m$.

Now suppose $p_s(F) = 4$, and let s, a, b, c be an induced path. The endpoint c is not a cut vertex: a neighbour of c in a component of $F - c$ that does not contain s would extend the path. Let $y \in N(c) \setminus \{b\}$. The path s, a, b, c, y is not induced, so y meets at least one of s, a, b. The edge ya would give the 4-cycle a, b, c, y. Thus y is adjacent to s or to b. The vertices s and c have at most one common neighbour, and so do b and c, hence $d(c) \leq 3$. If $d(c) = 3$, then c has two neighbours besides b, and one of them is adjacent to b. So c lies in a triangle, $t(c) \geq 1$ and $L(c) = 2d(c) - 3t(c) \leq 3$. Figure 1 shows the two deletions.


Date: November 2026.

(a) A longest induced path from s



Each extra neighbour y sees s or b (dashed edges). The edge ay would create a 4-cycle.

(b) Deleting two degree-2 endpoints



If $d(b) = d(c) = 2$, delete $U = \{b, c\}$. The path a, s, y survives and $D_G(U) = 6$.

FIGURE 1. Deletion at the end of a longest induced path of order four starting at s. (a) Each neighbor $y \neq b$ of c must be adjacent to s or b. Dashed lines indicate alternatives. (b) If neither terminal vertex c nor b can be deleted with loss at most three, the indicated 5-cycle is induced and both have degree two. The deletion of $\{b, c\}$ has loss six and preserves connectivity.

If $L(c) \leq 3$, delete c. The graph $F - c$ is connected, and $c \notin N[s]$, so D and m are unchanged. The inductive hypothesis gives $S(F - c) \leq 3(n-2) - D - m$, and therefore $S(F) \leq 3(n-2) - D - m + 3 = 3(n-1) - D - m$.

The remaining case is $d(c) = 2$ and $t(c) = 0$, so $L(c) = 4$. Let y be the neighbour of c other than b. Then y is not adjacent to b, hence ys is an edge, and s, a, b, c, y is an induced 5-cycle: every chord is excluded by the induced path s, a, b, c, by $t(c) = 0$, or by the 4-cycle that ay would create. The vertex b is the endpoint of the induced path s, y, c, b. If $L(b) \leq 3$, delete b and apply the inductive hypothesis as above.

Otherwise the same endpoint analysis gives $d(b) = 2$ and $t(b) = 0$. Delete $U = \{b, c\}$. A path that used the edges ab, bc, cy can be replaced along a, s, y, so $F - U$ is connected. The deleted edges are ab, bc and cy, and no triangle is deleted. Since $S = 2e - 3T$, one has $D_F(U) = 6 = 3|U|$. The vertices b and c lie outside $N[s]$, so D and m are unchanged. Induction gives $S(F - U) \leq 3(n-3) - D - m$, and $S(F) \leq 3(n-3) - D - m + 6 = 3(n-1) - D - m$.

For the equality examples, root C_5 at any vertex. Then $n = 5$, $D = 2$, $m = 0$ and $S = 10 = 3(n-1) - D - m$. If k copies of C_5 meet only at the root, then $n = 1 + 4k$, $D = 2k$ and $m = 0$, because the two neighbours of the root on each copy are nonadjacent, and $S = 10k = 3(n-1) - D - m$. □

Proposition 1.3. *Let F be connected and strong C_4-free, and let s satisfy $p_s(F) = 4$ and $\mathrm{ecc}(s) = 3$. Then $S(F) \leq 3(n-1) - D - m - 2$. Equality holds for P_4 rooted at an endpoint.*

Proof. The proof is by induction on n for instances with $p_s = 4$ and $\mathrm{ecc}(s) = 3$. Fix a vertex z at distance 3 from s. Whenever an endpoint other than z exists, the deletion in the proof of Theorem 1.2 applies to it. In the exceptional 5-cycle the vertices b and c are at distance 2 from s, so the deletion leaves z. The remaining graph is connected and strong C_4-free, D and m are unchanged, and distances do not decrease. Thus $d(s, z) \geq 3$. Since $p_s \leq 4$, no distance exceeds 3, so $p_s = 4$ and the eccentricity of s is still 3. A deletion of one vertex of loss at most 3 contributes 3, and the inductive bound on $n - 1$ vertices contributes $3(n-2) - D - m - 2$, which sum to $3(n-1) - D - m - 2$. Deleting $\{b, c\}$ contributes 6 against $3(n-3) - D - m - 2$.

If z is the unique endpoint, delete z. Then $p_s(F - z) \leq 3$. An extra neighbour of z is not adjacent to s, for that would give $d(s, z) \leq 2$. By the endpoint analysis it is adjacent to the predecessor of z and forms a triangle, so $L(z) \leq 2$. Lemma 5.3 gives $S(F - z) \leq 2(n-2) - m$,

hence $S(F) \le 2(n-1) - m$. A geodesic from s to z has at least two vertices outside $N[s]$, so $n - 1 - D \ge 2$ and $2(n-1) - m \le 3(n-1) - D - m - 2$.

For P_4 rooted at an endpoint one has $n = 4$, $D = 1$, $m = 0$ and $S = 6$, and the strengthened bound is $3 \cdot 3 - 1 - 2 = 6$. □

Proposition 1.4. *Let G be a counterexample to* GEN_4 *of minimum order. Then G is 2-connected. This does not decide* GEN_4.

Proof. Such a graph is connected and strong C_4-free, with $p(G) \le 8$ and $S(G) \ge 4n + 1$. Let v be a cut vertex, and let G_1, G_2 be the graphs of Lemma 2.3. Write ρ_i for the largest order of an induced path of G_i that starts at v. Paths from the two sides form an induced path of G on $\rho_1 + \rho_2 - 1 \le 8$ vertices, so some side, say G_1, has $\rho_1 \le 4$. Theorem 1.2 gives $S(G_1) \le 3(n_1 - 1)$. The graph G_2 is a smaller connected strong C_4-free graph of path order at most 8, so minimality gives $S(G_2) \le 4n_2$. Therefore $S(G) \le 3(n_1 - 1) + 4n_2$. Since $n = n_1 + n_2 - 1$,

$$3(n_1 - 1) + 4n_2 = 4n - n_1 + 1.$$

The hypothesis $n_1 \ge 2$ gives $4n - n_1 + 1 \le 4n - 1$, so $S(G) < 4n$, contradicting $S(G) \ge 4n + 1$. □

For $q \ge 5$ the bound $\mathrm{ROOT}(q)$ is not proved for arbitrary strong C_4-free graphs. Section 7 records the limitation. The next section proves every index for graphs whose blocks are bipartite or cycles. Conjecture 7.2 would imply $\mathrm{ROOT}(q)$ for the same q, by the attachment in Section 7. That implication is not a proof of the conjecture, and for $q \ge 5$ it does not prove $\mathrm{ROOT}(q)$.

2. Rooted bounds for every index

Let $\mathcal{B}$ be the class of connected strong C_4-free graphs in which every block is bipartite or a cycle. Edges are blocks, and cycles may be odd. The class contains every connected strong C_4-free bipartite graph and every connected strong C_4-free cactus. Theorem 2.2 keeps this block hypothesis. The graph $K_{2,3}$ does not show that the hypothesis is necessary. Its only block is itself, and that block is bipartite, but $K_{2,3}$ contains a 4-cycle subgraph. Rooted in its part of size two it has $p_s = 3$ and $S = 12 > 2(n-1) - m = 8$. The example tests the failure of $\mathrm{ROOT}(3)$ after the strong C_4-free hypothesis is removed. It does not test the block restriction inside the strong C_4-free class.

A rooted graph (G, s) with $p_s(G) = q$ and $\mathrm{ecc}(s) = q - 1$ is called *terminal*. For a terminal instance put $R = n - q + 1 - D \ge 0$, the number of vertices outside $N[s]$ and outside a root-to-farthest-vertex geodesic. Set $c_2 = 0$, $c_3 = 2$, $c_4 = 3$ and $c_q = 2\lfloor (q-1)/2 \rfloor$ for $q \ge 5$.

Lemma 2.1. *Let G be connected and strong C_4-free, and let P be a longest induced path starting at a vertex s, of order q. Let x be the other end of P, and suppose that the block B containing x is bipartite. Let Q be the subpath of P formed by the vertices of P that lie in B, and set $\ell = |Q|$. Then every neighbour of x lies in B, and x lies in no triangle. If $\ell = 2$, then $d(x) = 1$. If $\ell \ge 3$, then $d(x) \le \lfloor (\ell - 1)/2 \rfloor$. In every case $\ell \le q$, and $L(x) \le c_q$ whenever $q \ge 5$.*

Proof. A neighbour of x in a component of $G - x$ that does not contain s would extend P, so x is not a cut vertex. Every edge with one end in B and one end outside B is incident with a cut vertex of B. Therefore every neighbour of x lies in B. The block is bipartite, so x lies in no triangle.

Since the block-cut tree is a tree, P enters B only once. The intersection of P with B is a single subpath $Q = z_0, \dots, z_{\ell-1}$ with $z_{\ell-1} = x$. If s lies in B, then $Q = P$. If s lies outside B, the vertices of P that precede z_0 lie outside B, and z_0 is their only neighbour in B: any other edge from that prefix into B would be a second entry. A vertex of $B \setminus \{z_0\}$ is therefore adjacent to no vertex of the prefix. An extension of P at x through a vertex of B can fail only by a chord into Q.

Let y be a neighbour of x other than $z_{\ell-2}$. Then y lies in B, and y does not lie on Q, because Q is induced. The vertex y is not adjacent to the prefix. If y had no neighbour on Q, the path P followed by y would be induced. Thus y has a neighbour z_i on Q. This neighbour is not x. It is not $z_{\ell-2}$, for $yxz_{\ell-2}$ would be a triangle, and it is not $z_{\ell-3}$, for $z_{\ell-3}z_{\ell-2}xy$ would be a 4-cycle. Since B is bipartite, z_i has the colour of x, so $i \equiv \ell - 1 \pmod 2$ and $i \le \ell - 5$.

If $\ell = 2$, then Q is a single edge and there is no admissible index, so $d(x) = 1$. If $\ell \ge 3$, the number of admissible indices is $\lfloor(\ell-1)/2\rfloor - 1$. Two neighbours of x are nonadjacent and cannot share a blocker, for a common blocker would form a 4-cycle with them and with x. Hence $d(x) \le \lfloor(\ell-1)/2\rfloor$. In every case $\ell \le q$. For $q \ge 5$ one has $d(x) \le \lfloor(q-1)/2\rfloor$, and $L(x) = 2d(x) \le c_q$.

The same count applies inside B. Let z be the cut vertex of B lying on every path from a fixed root to B, or the root itself if the root lies in B, and let Q be a longest induced path of B starting at z, with other end x and with $\ell = |Q|$. A neighbour of x in B that had no neighbour on Q would extend Q inside B. If the root lies outside B, no vertex of $B \setminus \{z\}$ meets the component of the root at z, so that component contributes no further blocker. Thus $d_B(x) \le 1$ if $\ell = 2$, and $d_B(x) \le \lfloor(\ell-1)/2\rfloor$ if $\ell \ge 3$. □

Theorem 2.2. (1) *Every terminal instance in $\mathcal{B}$ satisfies $S(G) \le 2(D+q-2) - m + c_q R$.*

(2) *Every $F \in \mathcal{B}$ with a vertex s such that $p_s(F) \le q$ satisfies*

$$S(F) \le c_q(n-1) - (c_q - 2)D - m.$$

In particular ROOT(q) *holds in $\mathcal{B}$ for every $q \ge 3$. If every block is bipartite or a triangle, one may take $c_q = 2\lfloor(q-1)/2\rfloor$ for all $q \ge 3$.*

Proof. Two terminal facts. An endpoint x of a longest rooted induced path is not a cut vertex: a neighbour of x in another component of $G-x$ would extend the path. If z is at distance $q-1$ from s and $v_0 = s, \dots, v_{q-1} = z$ is a geodesic, any neighbour $y \ne v_{q-2}$ of z must meet an earlier path vertex (else the rooted path extends); meeting v_i with $i \le q-4$ shortens $d(s,z)$, and meeting v_{q-3} gives a C_4; so $y \sim v_{q-2}$, and two such y would give a C_4. Hence $L(z) = 2d(z) - 3t(z) \le 2$. Moreover, if $x \ne s, z$, $G - x$ is connected and z survives, then $q - 1 \le d_{G-x}(s,z) \le p_s(G-x) - 1 \le q - 1$, so the instance stays terminal with the same q.

Endpoint losses. Let x be the endpoint of a longest rooted path of order q. It lies in a single block. If that block is a triangle the loss $L(x)$ is 1, if an edge 2, and if a cycle of length at least 5 it is 4. If the block is bipartite, Lemma 2.1 gives $L(x) \le c_q$ for $q \ge 5$.

No nontrivial bipartite block when $q \le 4$. Let B be a nontrivial bipartite block, and let z be the cut vertex of B that lies on every path from s to B, or let $z = s$ if $s \in B$. Let Q be a longest induced path of B starting at z, and let x be its other end. Every vertex of a nontrivial block has degree at least two in the block, so $d_B(x) \ge 2$. The last paragraph of Lemma 2.1 gives $d_B(x) \le 1$ if $|Q| \le 4$ and $d_B(x) \le \lfloor(|Q|-1)/2\rfloor$ in any case with $|Q| \ge 3$. The inequality $d_B(x) \ge 2$ therefore forces $|Q| \ge 5$. An induced s–z path meets B only at z. Concatenating it with Q yields an induced path that starts at s. The order of this path is $|Q|$ if $s \in B$, and at least $|Q|+1$ if $s \notin B$. In either case $p_s(G) \ge 5$. A terminal instance with $q \le 4$ therefore has no nontrivial bipartite block. Every block is an edge, a triangle, or a cycle of length at least five, so the graph is a cactus.

Bases. For $q \le 4$ the preceding paragraph leaves only cacti. For a C_4-free cactus with c_3 triangle blocks and c_L longer cycle blocks, $S = 2(n-1) + 2c_L - c_3$ and $m \le c_3$. For $q = 3$ no longer cycle occurs, and (1) follows. For $q = 4$ every longer cycle is a C_5 through s. A chosen length-3 geodesic uses at most one edge of each such C_5: if it used two edges and then left the cycle, replacing that two-edge arc by the other three-edge arc would give an induced rooted path of order 5. So each C_5 contributes two vertices to R, $R \ge 2c_L$, and $[2(D+2) - m + 3R] - S = R - 2c_L + c_3 - m \ge 0$. For $q = 2$ the root is universal and $S = 2D - m$.

Induction on n. Let E be the set of endpoints of rooted induced paths of order q and z a farthest vertex, so $z \in E$. If some $x \in E \setminus \{z\}$ exists, delete it: $x \notin N[s]$, x is not a cut vertex, its loss is at most c_q, the instance remains terminal with the same q, D and m, and R drops by one. If $E = \{z\}$, delete z: the new rooted maximum is $q-1$, the eccentricity $q-2$, R is unchanged and the loss is at most 2; induction gives $S \le 2(D+q-3) - m + c_{q-1}R + 2 \le 2(D+q-2) - m + c_qR$. Deletion preserves membership in $\mathcal{B}$. This proves (1); the bipartite-or-triangle variant is identical.

(2) Attach a new path with $q-1$ edges at s. The result is terminal of index q, lies in $\mathcal{B}$, has order $n+q-1$, mass $S+2(q-1)$ and root degree $D+1$, and m is unchanged. Substituting into (1) gives (2). Since $c_q \le q-1$ and $n-1-D \ge 0$,

$$\begin{aligned} c_q(n-1) - (c_q-2)D - m &= c_q(n-1-D) + 2D - m \\ &\le (q-1)(n-1-D) + 2D - m \\ &= (q-1)(n-1) - (q-3)D - m, \end{aligned}$$

so ROOT(q) follows. □

A matching upper bound of shape $S \le (r-1)n - 2r(r-3)$, for graphs in $\mathcal{B}$ of radius $r \ge 5$ and path order $2r-1$, has been proposed from a structural description of radius-extremal graphs. That description is not proved in this paper, and the radius bound in $\mathcal{B}$ is not claimed.

Lemma 2.3. *Let G be strong C_4-free and triangle-free, and let $x_0x_1\cdots x_k$ be an induced path that cannot be extended at x_k. If $d(x_k) \ge 2$, then $k \ge d(x_k)+1$. If moreover G is bipartite, then $k \ge 2d(x_k)$.*

Proof. Let $y \ne x_{k-1}$ be a neighbour of x_k. It is not on the path, and, since the path cannot be extended, y has a neighbour x_i with $i \le k-1$. Here $i \ne k-1$ (a triangle) and $i \ne k-2$ (the 4-cycle $x_{k-2}x_{k-1}x_ky$), so $i \le k-3$. A vertex x_i is adjacent to at most one neighbour of x_k. Hence the $d(x_k)-1 \ge 1$ such neighbours need distinct blockers among $x_0, \dots, x_{k-3}$, and $k-2 \ge d(x_k)-1$. If G is bipartite, the blocker has the colour of x_k, so $i \equiv k \pmod 2$ and $i \le k-4$; there are $\lfloor k/2 \rfloor - 1$ such positions, and $\lfloor k/2 \rfloor \ge d(x_k)$. □

Proposition 2.4. *Every connected regular bipartite strong C_4-free graph satisfies* GEN_K *for every K and satisfies REAL-133. Every Moore graph of diameter* 2 *and girth* 5 *satisfies REAL-133.*

Proof. Let G be d-regular. If $d = 1$ then $G = K_2$, with $r = 1$, $\ell = 1$, $p = 2$ and $S = n$, and both statements hold. Let $d \ge 2$. As G is triangle-free, $\lambda \equiv d$ and $\ell = d$. Lemma 2.3 applied to a longest induced path gives $p \ge 2d+1$, and Theorem 2.4 gives $p \ge 2r-1$; so $p \ge r+d$ both when $r \le d+1$ and when $r \ge d+1$. If $p \le 2K$, then $2d+1 \le 2K$, so $S = dn < Kn$. A Moore graph of diameter 2 and girth 5 has degree $d \ge 2$, radius 2 and $\ell = d$. The degree is an integer, so the floor form and the real form ask for the same inequality. The first part of Lemma 2.3 gives $p \ge d+2 = r+\ell$. This is an alternate proof of the case already recorded by Chen [3] immediately after his Theorem 3.6. It is not a new extremal classification. □

Remark 2.5. A Moore graph of degree 57, if it exists, has $n = 57^2 + 1 = 3250$ and $S = 57n$. Proposition 2.4 gives REAL-133, which only needs $p \ge 59$, and this conclusion is the integer case of the fact cited above. The statement GEN_{57} holds for it with equality in the mass bound, but GEN_{56} asks for $p \ge 113$ because $S > 56n$. We do not know whether such a graph would have $p \le 112$. If it did, it would refute GEN_{56} while satisfying REAL-133. Its existence would therefore not by itself separate the two statements, but it shows that a proof of every GEN_K is a different assertion from REAL-133.

3. Negative results

The examples in this section are explicit graphs.

Example 3.1. Let $s \geq 3$, $t \geq 3$ and $m \geq 1$ be integers. Let $G_{s,t,m}$ be the cycle $C_{2s} = c_0 \cdots c_{2s-1}$ with a pendant path of t edges at every cycle vertex, together with m internally disjoint paths c_0xzyc_1. Then $n = 2s(t+1) + 3m$, $e = 2s(t+1) + 4m$ and $T = 0$, so $e - n - T = m \geq 1$. For $(s,t,m) = (3,3,1)$ one has $n = 27$, $e = 28$, $r = 6$ and $p = 11$. These graphs satisfy REAL-133, since $\ell = 2e/n < 8/3$. They also show that $e = n + T$ fails for radius-extremal strong C_4-free graphs of every radius $r \geq 6$: here $r(G_{s,t,m}) = s + t$ and $p(G_{s,t,m}) = 2(s+t) - 1$.

Two vertices have at most one common neighbour, so G is strong C_4-free and triangle-free. An added path of length four does not shorten distances among the older vertices, and c_0 remains within distance two of every vertex of such a path, so the eccentricity of c_0 is still $s + t$. A vertex w of an added path meets the tip of the arm at c_s only through c_0 or c_1, hence at distance at least $s + t$. Thus $r = s + t$. For the path order, let B be the block formed by C and the added paths. An induced path P meets B in a subpath Q and can leave B only at an end of Q that is a cycle vertex, through at most t arm vertices. The same case division as for a single ear shows $|P| \leq 2s + 2t - 1$, using $t \geq 3$ and $s \geq 3$, and the cycle minus one vertex together with the arms at its two ends attains this order.

The last sentence of Example 3.1 abbreviates a case division which we now record. If Q avoids both c_0 and c_1, then $|Q| \leq 2s-2$. If Q contains exactly one of them, then either Q lies on the cycle with $|Q| \leq 2s - 1$, or $|Q| \leq 2s + 2$ with at most one cycle end, or $|Q| \leq 7$ with no cycle end. If Q contains both, the same three shapes occur with $|Q| \leq 2s-1$, $|Q| \leq 2s+2$ and $|Q| \leq 6$. Adding at most t arm vertices at each cycle end of Q gives $|P| \leq \max\{2s-1+2t,\ 2s+2+t,\ 7\} = 2s+2t-1$.

Example 3.2. Forbidding induced paths on eleven vertices and requiring $L(v) \geq 6$ at every vertex does not forbid an induced star $K_{1,6}$. Two Petersen graphs sharing a vertex give a triangle-free strong C_4-free graph on $n = 19$ vertices with $S = 60$, minimum deletion loss 6, $p = 9$ and $\lambda = 6$ at the shared vertex. For $b \geq 2$, gluing b copies of the Petersen graph along a common edge gives a 2-connected graph B_b with $n = 8b + 2$, $S = 28b + 2$, minimum loss 6, $p = 9$ and $\lambda = 2b + 1$ at the two hubs. Thus singleton losses do not bound λ. These graphs are not counterexamples to GEN_5: the eight non-hub vertices of one Petersen block have deletion loss $28 < 5 \cdot 8 + 1$.

For an oriented edge (u, v) of a connected strong C_4-free graph G, let q be the maximum order of an induced path whose first two vertices are u, v in that order, and let E be the set of terminal vertices of the induced paths of that order. The endpoint inequality $\mathrm{EP}(q)$ asserts $D_G(E) \leq q|E|$ for every such oriented edge of maximum order q. For a root s with $p_s(G) = q$, if E is the set of terminal vertices of the induced paths of order q starting at s, the rooted inequality $\mathrm{EP}^+(q)$ asserts $D_G(E) \leq (q-1)|E|$.

Proposition 3.3. *For every integer $q \geq 4$, $\mathrm{EP}(q)$ holds for all connected strong C_4-free graphs if and only if $\mathrm{EP}^+(q-1)$ does.*

Proof. Both graphs below remain strong C_4-free, so $S = 2e - 3T$ and the deletion loss of a vertex set is determined by the edges and triangles that meet it.

First assume $\mathrm{EP}^+(q-1)$, and let (u, v) be an oriented edge of maximum induced-path order q, with terminal set E. Delete $W = N[u] \setminus \{v\}$, and let H be the component of v in $G - W$. An induced path of G that starts with uv has no further vertex in $N[u]$, because the path is induced, so the remainder is an induced path of H starting at v. Conversely, every induced path of H starting at v extends by u to an induced path of G, since no vertex of H other than v is adjacent to u. Thus $p_v(H) = q - 1$, and the terminal set in H is still E.

Each $x \in E$ has at most one neighbour in W: any such neighbour is adjacent to u, and two of them would be two common neighbours of u and x. Every neighbour of x outside H lies in W. Let m_W be the number of edges between E and W, so $m_W \leq |E|$, and let T_{extra} be the number of triangles that meet E and are not contained in H. Every such triangle uses one of those edges.

Comparing edges and triangles,

$$D_G(E) = D_H(E) + 2m_W - 3T_{\text{extra}} \le D_H(E) + 2|E|.$$

The rooted inequality in H is $D_H(E) \le (q-2)|E|$, because $\mathrm{EP}^+(q-1)$ contributes the factor $(q-1)-1$. Therefore $D_G(E) \le q|E|$.

Conversely, assume $\mathrm{EP}(q)$, and let (H, v) be a connected strong C_4-free graph with $p_v(H) = q-1$ and terminal set E. Form G by adding a vertex u adjacent to v and, for each $x \in E$, a new vertex a_x adjacent exactly to u and to x. Every terminal x fails to be adjacent to v: an induced path of order $q-1 \ge 3$ from v to x has no chord vx. The vertex a_x has degree two. Any cycle through a_x uses both edges ua_x and a_xx, and then continues from x to the other neighbour of u on the cycle. That other neighbour is v or some a_y. In the first subcase the x–v path in H has length at least two, so the cycle has length at least five. In the second the cycle has the form $u, a_x, x, \ldots, y, a_y$ and again has length at least five. A cycle through u that avoids every a_x would need two neighbours of u in H, but the only neighbour of u in H is v. Thus G is strong C_4-free. No new triangle appears: the two neighbours of a_x are nonadjacent, and the neighbours of u are pairwise nonadjacent.

An induced path that starts with uv contains no a_x. Otherwise the edge ua_x would be a chord, since a_x is not the successor of u on that path. The induced uv-paths are therefore the paths u followed by an induced path of H starting at v. Their maximum order is q, and their terminal set is E. Exactly $|E|$ new edges meet E, namely the edges a_xx, and no new triangle meets E. Hence $D_G(E) = D_H(E) + 2|E|$. The oriented inequality gives $D_G(E) \le q|E|$, so $D_H(E) \le (q-2)|E|$, which is $\mathrm{EP}^+(q-1)$. □

Example 3.4. EP(9) fails. Let G have vertices $0, \ldots, 14$ and adjacency lists

0: $4, 8, 11$	1: $5, 8, 10$	2: $6, 8, 9$	3: $7, 10$
4: $0, 9, 10$	5: $1, 9, 11$	6: $2, 10, 11$	7: $3, 11$
8: $0, 1, 2, 12$	9: $2, 4, 5$	10: $1, 3, 4, 6, 13$	11: $0, 5, 6, 7, 14$
12: $8, 13, 14$	13: $10, 12$	14: $11, 12$.	

The graph is triangle-free and strong C_4-free, with $n = 15$, $e = 23$ and $S = 46$. For the oriented edge $(12, 8)$ the maximum order is $q = 9$, attained only by

$$12, 8, 2, 9, 4, 10, 3, 7, 11 \qquad \text{and} \qquad 12, 8, 2, 9, 5, 11, 7, 3, 10.$$

Thus $E = \{10, 11\}$ and $D_G(E) = 20 > 18$. No induced path that starts with $12, 8$ has order greater than 9. The neighbours of 8 other than 12 are $0, 1, 2$. A continuation is legal only when the new vertex meets the current end and meets no earlier vertex of the path. Under that rule every maximal induced path is one of the following, and the only two of order 9 are the paths already displayed.

maximal induced path starting $12, 8$	order
$12, 8, 0, 4, 9, 5$	6
$12, 8, 0, 4, 10, 3, 7$	7
$12, 8, 0, 4, 10, 6$	6
$12, 8, 0, 11, 5, 9$	6
$12, 8, 0, 11, 6, 10, 3$	7
$12, 8, 0, 11, 7, 3, 10$	7
$12, 8, 1, 5, 9, 4$	6
$12, 8, 1, 5, 11, 6$	6
$12, 8, 1, 5, 11, 7, 3$	7
$12, 8, 1, 10, 3, 7, 11$	7
$12, 8, 1, 10, 4, 9$	6
$12, 8, 1, 10, 6, 11, 7$	7
$12, 8, 2, 6, 10, 3, 7$	7
$12, 8, 2, 6, 10, 4$	6
$12, 8, 2, 6, 11, 5$	6
$12, 8, 2, 6, 11, 7, 3$	7
$12, 8, 2, 9, 4, 10, 3, 7, 11$	9
$12, 8, 2, 9, 5, 11, 7, 3, 10$	9

The first of the two paths of order 9 ends at 11. The neighbours of 11 are $0, 5, 6, 7, 14$: the vertex 7 is the predecessor, 0 meets 8, 5 meets 9, 6 meets both 2 and 10, and 14 meets 12. The second path ends at 10. The neighbours of 10 are $1, 3, 4, 6, 13$: the vertex 3 is the predecessor, 1 meets 8, 4 meets 9, 6 meets both 2 and 11, and 13 meets 12. Neither path extends. The same computation gives a failure of $\mathrm{EP}^{+}(8)$ on the subgraph induced by $0, \dots, 11$, rooted at 8: $p_8 = 8$, $E = \{10, 11\}$ and $D(E) = 16 > 14$. The inequalities $\mathrm{EP}(8)$ and $\mathrm{EP}^{+}(6)$ are not settled here.

Example 3.5. Let $k = p - r$. Call an induced subgraph H of G an *eligible core* if it is proper, connected and dominating, and $p(H) \le p(G) - 2$. The budget CB asks for an eligible core with

$$S(G) - S(H) \le k\big(n - |H|\big) + |H|.$$

Together with REAL-133 for H and an induction on the order, the budget would imply REAL-133. The Petersen graph has $S = 30$, $p = 5$, $r = 2$ and $k = 3$, hence $\ell = 3$. An eligible core has $p(H) \le 3$. A connected triangle-free strong C_4-free P_4-free graph is a star, by Lemma 2.7 and the absence of triangles. In the Petersen graph a star with $k \le 3$ leaves has centre of degree 3. The vertices it dominates are the centre, its three neighbours, and at most two further neighbours of each leaf, at most $4 + 2k$ vertices in all. This is 10 only for $k = 3$, so the eligible cores are exactly the ten closed neighbourhoods $N[v]$. Each has $S(H) = 6$, and $S(G) - S(H) = 24 > 3 \cdot 6 + 4 = 22$.

The unrestricted budget therefore fails. The example has $\ell = 3$. It does not determine a cutoff such as $\ell > 5$, and this paper contains no census of eligible cores.

Example 3.6. The degree, triangle and distance data of the Heawood graph ($n = 14$, cubic, $T = 0$, $r = 3$, $S = 42$, and $Z = 28$ nonadjacent pairs with no common neighbour) satisfy the counting identities used above, including $\sum_v d(v)^2 = n(n-1) + 6T - 2Z$, together with the lower bounds $p \ge 2r - 1$ and $p \ge \min_v \lambda(v) + 2$, if p is replaced by the value 5. But $5 < r + \ell = 6$. This system of scalar constraints does not imply REAL-133. The Heawood graph itself has $p = 7$.

Example 3.7. Let F be the incidence graph of the projective plane $\mathrm{PG}(2, q)$, with $f = |F| = 2(q^2 + q + 1)$ and $S(F) = (q + 1)f$, and choose a root of F. Let $t = f - 1$ and $b \ge 1$. Take a 6-cycle $c_0 \cdots c_5$, attach a pendant path with t edges at each cycle vertex, and attach b copies of F at c_0 by identifying their roots with c_0. The result is strong C_4-free, because each new block meets the rest in a cut vertex, and

$$n = 6f + b(f - 1), \qquad S = 12f + b(q + 1)f, \qquad r = t + 3, \qquad p = 2t + 5 = 2r - 1.$$

Distances inside the flower are unchanged. Every vertex of an attached copy is within distance $f-1 \le t$ of c_0, so each cycle vertex has eccentricity $t+3$, while a vertex x of a copy has distance $d_F(x,c_0)+t+3$ from the tip of the arm at c_3. An induced path meets the cycle in at most five consecutive vertices, and each of its two ends outside the cycle contributes at most t vertices. Mass is additive at cut vertices by Lemma 2.3. For fixed q, as $b \to \infty$ one has $\ell \to (q+1)f/(f-1)$. Letting $q \to \infty$, with $r = f+2$, this limit grows like $\sqrt{r/2}$. For $q = 5$ and $b = 17$ one has $n = 1409$, $r = 64$, $p = 127$ and $S = 7068 > 5n$. These graphs satisfy RAD(r) for their own radius, and they show that ℓ is unbounded on the class $p = 2r-1$. Any theorem RAD(r) must use a bound that depends on r.

Appendix A. A conditional reduction for GEN_3

Theorem 5.4 is the triangle-free case of GEN_3. The three local statements named at the end of this appendix are not proved for graphs that contain triangles. With them, the hand arguments below would give GEN_3 in general. Without them, that case remains open. The cell $r = 3$, $p = 6$, $T \ge 2$, $n \ge 17$ and $3 < \ell \le 4 - 16/n$ remains open as well. By Corollary 5.11 one has $\ell \ne 4$, and graphs with at most one triangle are excluded by Theorems 5.4 and 5.6.

Theorem A.1. *Let G be a counterexample to GEN_3 of minimum order: G is connected and strong C_4-free, $p(G) \le 6$, $S(G) \ge 3n+1$, and n is minimum. Then G is 2-connected; $S(G) - S(G-U) \ge 3|U|+1$ for every nonempty proper $U \subset V(G)$; $2\lambda(v) - t(v) \ge 4$ for every v, so $\lambda(v) \ge 2$; and every vertex with $\lambda(v) = 2$ has degree 2 and nonadjacent neighbours, and no two vertices of degree 2 are adjacent.*

Proof. Since $S \le 2 < 3n$ for $n \le 2$, one has $n \ge 3$. Let v be a cut vertex. For each component C of $G-v$ let ρ_C be the largest order of an induced path in $G[C \cup \{v\}]$ with end v. Two such paths from different components form an induced path on $\rho_C + \rho_{C'} - 1 \le 6$ vertices, so some component C_1 has $\rho_{C_1} \le 3$. Put $G_1 = G[C_1 \cup \{v\}]$ and $G_2 = G - C_1$, with $n_i = |G_i|$, so $n = n_1 + n_2 - 1$ and $n_1 \ge 2$. Both graphs are connected, smaller than G, and P_7-free. Lemma 5.3 gives $S(G_1) \le 2(n_1-1)$, and minimality gives $S(G_2) \le 3n_2$. By Lemma 2.3, $S(G) \le 2(n_1-1) + 3n_2 = 3n - (n_1 - 1) < 3n$.

Every component H of $G-U$, for U nonempty and proper, is connected, smaller, and P_7-free, so $S(H) \le 3|H|$. Summing, $S(G-U) \le 3(n-|U|)$, and $S(G) \ge 3n+1$ gives the deletion bound. The case $U = \{v\}$ and Lemma 2.2 give $2\lambda(v) - t(v) \ge 4$. If $\lambda(v) = 2$, then $t(v) = 0$, so $d(v) = 2$ and the neighbours are nonadjacent. Every degree-2 vertex has $\lambda = 2$ and hence lies in no triangle. If $u \sim w$ both have degree 2, with other neighbours x and y, then $x \ne y$ because $t(u) = 0$. The set $\{u\}$ is a unit of x and $\{w\}$ is a unit of y, so Lemma 2.2 with $U = \{u, w\}$ gives loss $2+2+1+1 = 6 < 7$. □

The same mass bound for a disconnected graph is the sum of the bounds on its components. An isolated vertex has $S = 0$. Call a graph *admissible* if it is strong C_4-free and P_7-free, $L(v) \ge 4$ for every vertex, and no two adjacent vertices have degree 2. A minimum counterexample is admissible. In an admissible graph, $\lambda(v) = 2$ forces $t(v) = 0$ and $d(v) = 2$. Write B for the vertices of degree at least 3, the *branch vertices*, and Z for the degree-2 vertices, the *ears*. Put $D = G[B]$, $d_D(v) = |N(v) \cap B|$ and $q(v) = |N(v) \cap Z|$. All triangles lie in D.

Lemma A.2. *In an admissible graph, $q(v) \le 4$ for every vertex v.*

Proof. Let u_i be ears at v with other neighbours a_i. The a_i are distinct, nonadjacent to v, and of degree at least 3. Put $E_i = N(a_i) \setminus \{u_i\}$. Then $|E_i| \ge 2$ and $E_i \cap N[v] = \emptyset$.

If $a_i \not\sim a_j$, then $E_i = \{w, p\}$ and $E_j = \{w, q\}$ with $pq \in E$, $w \not\sim p$ and $w \not\sim q$. Indeed two common neighbours of a_i and a_j are forbidden, so $|E_i \cap E_j| \le 1$. Since $|E_i| \ge 2$, the private part $E_i \setminus E_j$ is nonempty, and likewise $E_j \setminus E_i$. If p lies in the first private part and q in the second with $p \not\sim q$, the path $pa_iu_ivu_ja_jq$ is induced. Thus the private parts are complete to each other.

If one of them contained two vertices, those two vertices and one vertex of the other private part would form a 4-cycle with the corresponding a-vertex. Both private parts therefore have size one, and $|E_i \cap E_j| = 1$. An edge from the common vertex to a private vertex is another 4-cycle. In particular a_i and a_j have degree 3 and lie in no triangle.

No three of the vertices a_i are pairwise nonadjacent. Otherwise the sets E_i meet pairwise in a single vertex. If the three intersections are the three pairs of a triple, the claim forces a triangle on that triple, and one a_i together with two vertices of the triple forms a 4-cycle. Otherwise $E_i = \{w, p_i\}$ with $p_1p_2p_3$ a triangle and w nonadjacent to each p_i, and $vu_1a_1wa_2p_2p_3$ is an induced P_7.

Five ears are impossible. The graph F they determine on $\{a_1, \dots, a_5\}$ has independence number at most two. If some vertex a of F is adjacent to the other four, those four induce a matching. The independence number forces that matching to be perfect, say $\{b_1b_2, c_1c_2\}$. Then $b_1 \not\sim c_1$, so the preceding paragraph says that b_1 lies in no triangle, but ab_1b_2 is a triangle. Otherwise every vertex of F has a non-neighbour, hence has degree 3 in G and lies in no triangle. One neighbour of each such vertex is its ear, so F has maximum degree at most two. It is triangle-free and has no 4-cycle, and its independence number is at most two, so F is a 5-cycle. Label that cycle $a_1a_2a_3a_4a_5a_1$, and write u_i for the ear joining v to a_i. The seven vertices $u_5, v, u_1, a_1, a_2, a_3, a_4$, in that order, induce a P_7: each u_i has neighbours v and a_i only, v is nonadjacent to every a_j, and the only edges of the cycle among a_1, a_2, a_3, a_4 are the three path edges. □

Proposition A.3. *Let G be admissible, with B, Z, D, d_D and q as above. Suppose that $d_D(v) \le 3$ for every v, that $d_D(v) = 3$ implies $q(v) = 0$, and that $d_D(v) = 2$ with the two neighbours in B nonadjacent implies $q(v) \le 2$. Then $S(G) \le 3n$.*

Proof. Ears have nonadjacent neighbours and Z is an independent set, so $S(G) = S(D) + 4|Z|$. Since $|Z| = \frac{1}{2}\sum_{v\in B} q(v)$ and $n = |B| + |Z|$,

$$S - 3n = \sum_{v\in B}\Big(d_D(v) - t_D(v) - 3 + \tfrac{1}{2}q(v)\Big).$$

Lemma A.2 gives $q(v) \le 4$. If $d_D(v) = 2$ and $t_D(v) = 0$, the two neighbours in B are nonadjacent, for an edge between them would be a triangle. The three suppositions and the ear bound make every summand nonpositive: the summand is at most -1 when $(d_D, t_D) = (0, 0)$, at most 0 when $(d_D, t_D) = (1, 0)$, at most 0 when $(d_D, t_D) = (2, 0)$, at most 0 when $(d_D, t_D) = (2, 1)$, and equal to $-t_D(v) \le 0$ when $d_D(v) = 3$. The suppositions exclude $d_D(v) \ge 4$. □

The three suppositions in Proposition A.3 are not proved in this paper. Theorem A.1 shows that a minimum counterexample to GEN_3 is admissible, so those suppositions would imply GEN_3. They do not, by themselves, make GEN_3 a theorem.

Acknowledgments

The authors are responsible for the text, including any errors. Language-model tools were used to prepare drafts.

References

[1] E. Camby and O. Schaudt, *A new characterization of P_k-free graphs*, Algorithmica **75** (2016), 205–217.

[2] A. Brandstädt and C. T. Hoàng, *On clique separators, nearly chordal graphs, and the Maximum Weight Stable Set Problem*, Theoret. Comput. Sci. **389** (2007), 295–306.

[3] H. Chen, *Partial results on the WOWII-133 Graffiti conjecture*, Zenodo, 2026, `https://doi.org/10.5281/zenodo.22054834` (version record 22054835).

[4] M. Chudnovsky and J. Stacho, *3-colorable subclasses of P_8-free graphs*, SIAM J. Discrete Math. **32** (2018), 1111–1138.

[5] R. Chen and B. Xu, *Nearly optimal coloring of some C_4-free graphs*, Discrete Appl. Math. (2026), https://doi.org/10.1016/j.dam.2026.07.010; arXiv:2409.06944.
[6] R. Chen, D. Wu and B. Xu, *Structure of some (P_7, C_4)-free graphs with application to colorings*, Discrete Appl. Math. **357** (2024), 14–23; arXiv:2302.11115.
[7] E. DeLaVina, *Written on the Wall II (Conjectures of Graffiti.pc)*, http://cms.dt.uh.edu/faculty/delavinae/research/wowII/.
[8] E. DeLaViña, C. E. Larson, R. Pepper and B. Waller, *A characterization of graphs where the independence number equals the radius*, Graphs Combin. **28** (2012), 315–332.
[9] S. Fajtlowicz, *A characterization of radius-critical graphs*, J. Graph Theory **12** (1988), 529–532.
[10] P. Erdős, M. Saks and V. T. Sós, *Maximum induced trees in graphs*, J. Combin. Theory Ser. B **41** (1986), 61–79.
[11] Formal Conjectures (Google DeepMind), file FormalConjectures/WrittenOnTheWallII/GraphConjecture133.lean at commit 5faef0c4dc3d226f8f9a9fc8f581b6221283eb6e, https://github.com/google-deepmind/formal-conjectures/blob/5faef0c4dc3d226f8f9a9fc8f581b6221283eb6e/FormalConjectures/WrittenOnTheWallII/GraphConjecture133.lean.
[12] Formal Conjectures, pull request #5727, merged as commit 5faef0c4dc3d226f8f9a9fc8f581b6221283eb6e, https://github.com/google-deepmind/formal-conjectures/pull/5727.
[13] T. Karthick and F. Maffray, *Square-free graphs with no six-vertex induced path*, SIAM J. Discrete Math. **33** (2019), 874–909; arXiv:1805.05007.
[14] D. Seinsche, *On a property of the class of n-colorable graphs*, J. Combin. Theory Ser. B **16** (1974), 191–193.