

Solutions for
Analysis I&II (Forth Edition)
by Terence Tao

Ilia Koloiarov

June 28, 2026

Contents

1	Introduction	2
2	Starting at the beginning: the natural numbers	3
2.1	The Peano axioms	4
2.2	Addition	6
2.3	Multiplication	13
3	Set Theory	19
3.1	Fundamentals	20
A	The basics of mathematical logic	21
A.1	Mathematical statements	22
A.2	Implication	23
A.3	The structure of proofs	24
B	The decimal system	25

Chapter 1

Introduction

No exercises in this chapter.

Chapter 2

Starting at the beginning: the natural numbers

2.1 The Peano axioms

Axiom 2.1. *0 is a natural number.*

Axiom 2.2. *If n is a natural number, then $n++$ is also a natural number.*

Axiom 2.3. *0 is not the successor of any natural number; i.e., we have $n++ \neq 0$ for every natural number n .*

Axiom 2.4. *Different natural numbers must have different successors; i.e., if n, m are natural numbers and $n \neq m$, then $n++ \neq m++$. Equivalently, if $n++ = m++$, then we must have $n = m$.*

Axiom 2.5. (Principle of mathematical induction). *Let $P(n)$ be any property pertaining to a natural number n . Suppose that $P(0)$ is true, and suppose that whenever $P(n)$ is true, $P(n++)$ is also true. Then $P(n)$ is true for every natural number n .*

Definition 2.1.3. *We define 1 to be the number $0++$, 2 to be the number $(0++)++$, etc.*

Proposition 2.1.4. *3 is a natural number.*

Proof. Given the Axiom 2.1, 0 is a natural number. $1 := 0++$ is a natural number by Axiom 2.2. Same as $2 := 1++$ and $3 := 2++$. $\square$

Proposition 2.1.6. *4 is not equal to 0.*

Proof. By Definition 2.1.3, $4 := 3++$, thus $3++ \stackrel{\neq}{=} 0$ what contradicts Axiom 2.3 $\square$

Proposition 2.1.8. *6 is not equal to 2*

Proof. Let's assume that $6 = 2$. By Definition 2.1.3, $6 = 5++$ and $2 = 1++$. That means, $5 = 1$ by Axiom 2.4. Repeating the procedure, we end up with $4 \stackrel{\neq}{=} 0$ what contradicts previously proven Proposition 2.1.6 $\square$

Lemma 2.1.8.a. *For any natural number n , $n \neq n++$.*

Proof. For the base case, $0 \neq 0++$ by Axiom 2.3. For the step case, we need to prove if $n \neq n++$, then $n++ \neq (n++)++$. Let's assume that $n++ = (n++)++$. Then by Axiom 2.4, $n \stackrel{\neq}{=} n++$, what contradicts the inductive hypothesis. Now we have closed the induction. $\square$

Proposition 2.1.16. *(Recursive definitions). Suppose for each natural number n , we have some function $f_n : \mathbb{N} \rightarrow \mathbb{N}$ from the natural numbers to the natural numbers. Let c be a natural number. Then we can assign a unique natural number to each natural number n , such that $a_0 = c$ and $a_{n++} = f_n(a_n)$ for each natural number n .*

Proof. First, none of a_n redefines a_0 by Axiom 2.3. Second, let's assume that the procedure gives the unique value for a_n . Then it also provides the unique value for a_{n++} by Axiom 2.4 what closes the induction. $\square$

2.2 Addition

Definition 2.2.1. (*Addition of natural numbers*). Let m be a natural number. To add zero to m , we define $0 + m := m$. Now suppose inductively that we have defined how to add n to m . Then we can add $n++$ to m by defining $(n++) + m := (n + m)++$.

Proposition 2.2.1.a. *The sum of two natural numbers is again a natural number.*

Proof. We induct on n . Using Definition 2.2.1, $0 + m := m$ is a natural number by the proposition's assumption. Let's assume that $n + m$ is a natural number. Then $(n++) + m := (n + m)++$ by Definition 2.2.1 and $(n + m)++$ is a natural number by Axiom 2.2. That closes the induction. $\square$

Lemma 2.2.2. *For any natural number n , $n + 0 = n$.*

Proof. Guess what, we use induction. Given that 0 is a natural number and Definition 2.2.1, $0 + 0 := 0$ what proves the base case. Let's assume that $n + 0 := n$, then $(n++) + 0 := (n + 0)++$ using Definition 2.2.1, and $(n + 0)++ = n++$ by the induction assumption what proves the step case. That closes the induction. $\square$

Lemma 2.2.3. *For any natural numbers n and m , $n + (m++) = (n + m)++$.*

Proof. Let's induct on n . To prove the base case, $0 + (m++) := m++ := (0 + m)++$ using Definition 2.2.1. For the step case, we need to prove $(n++) + (m++) = ((n++) + m)++$. Let's assume $n + (m++) = (n + m)++$, then

$$\begin{aligned} (n++) + (m++) &:= (n + (m++))++ && \text{Definition 2.2.1} \\ &= ((n + m)++)++ && \text{Inductive Hypothesis} \\ &= ((n++) + m)++ && \text{Definition 2.2.1} \end{aligned}$$

what closes the induction. $\square$

Corollary 2.2.3.a. *For any natural number n , $n++ = n + 1$.*

Proof. The base case is set up by Definition 2.1.3. For the step case, we need to prove $(n++)++ = (n++) + 1$, assuming $n++ = n + 1$:

$$\begin{aligned} (n++)++ &= (n + 1)++ && \text{Inductive Hypothesis} \\ &= (n++) + 1 && \text{Definition 2.2.1} \end{aligned}$$

$\square$

Proposition 2.2.4. (Addition is commutative). *For any natural numbers n and m , $n + m = m + n$.*

Proof. Let's (again) induct on n . For the base case we need to prove $0 + m = m + 0$. The left term equals to m by Definition 2.2.1. The right term equals also to m by Lemma 2.2.2. Thus, both terms are equal. For the step case, we need to prove $(n++) + m = m + (n++)$, assuming $n + m = m + n$:

$$\begin{aligned}
 (n++) + m &= (n + m)++ && \text{Definition 2.2.1} \\
 &= (m + n)++ && \text{Inductive Hypothesis} \\
 &= (m + (n++)) && \text{Lemma 2.2.3}
 \end{aligned}$$

what closes the induction. □

Proposition 2.2.5. (Addition is associative). *For any natural numbers a, b, c , we have $(a + b) + c = a + (b + c)$.*

Proof. Let's induct on a . For the base case, we need to prove

$$\begin{aligned}
 (0 + b) + c &= 0 + (b + c) \\
 \text{Definition 2.2.1} \quad & \quad b + c = b + c \quad \quad \text{Lemma 2.2.2}
 \end{aligned}$$

For the step case, we need to prove $((a++) + b) + c = (a++) + (b + c)$, assuming $(a + b) + c = a + (b + c)$:

$$\begin{aligned}
 ((a++) + b) + c &= ((a + b)++) + c && \text{Definition 2.2.1} \\
 &= ((a + b) + c)++ && \text{Definition 2.2.1} \\
 &= (a + (b + c))++ && \text{Inductive Hypothesis} \\
 &= (a++) + (b + c) && \text{Definition 2.2.1}
 \end{aligned}$$

what closes the induction. □

Proposition 2.2.6. (Cancellation Law). *Let a, b, c be natural numbers such $a + b = a + c$. Then we have $b = c$.*

Proof. Let's induct on a . The base case, we need to prove $0 + b = 0 + c$ what leads to $b = c$ by Definition 2.2.1. For the step case, we need to prove if $(a++) + b = (a++) + c$,

assuming $a + b = a + c$, then $b = c$.

$$(a++) + b = (a++) + c$$

$$(a + b)++ = (a + c)++$$

$$(a + b) = (a + c)$$

$$b = c$$

Definition 2.2.1

Axiom 2.4

Inductive Hypothesis

□

Definition 2.2.7. (Positive natural numbers). *A natural number n is said to be positive iff it is not equal to 0.*

Proposition 2.2.8. *If a is positive and b is a natural number, then $a + b$ is positive.*

Proof. Let's induct on b . For the base case, $a + 0 = a$ by Definition 2.2.1. The result is positive by the propositional assumption. For the step case, we need to prove $a + (b++)$ is positive, assuming $a + b$ is positive. $a + (b++) = (a + b)++$ by Lemma 2.2.3. Given Axiom 2.3, $(a + b)++$ is positive as well, what closes the induction. □

Corollary 2.2.9. *If a and b are natural numbers such that $a + b = 0$, then $a = 0$ and $b = 0$.*

Proof. Let's assume that $a \neq 0$ or $b \neq 0$. In either case, $a + b$ is a positive number by Proposition 2.2.8 what contradicts $a + b \stackrel{\text{def}}{=} 0$. □

Lemma 2.2.10. *Let a be a positive number. Then there exists exactly one natural number b such that $b++ = a$.*

Proof. Suppose for sake of contradiction, there is another natural number c such that $c++ = a$. Then $b++ = c++$, what makes $b = c$ by Axiom 2.4. □

Definition 2.2.11. (Ordering of the natural numbers). *Let n and m be natural numbers. We say that n is greater than or equal to m , and write $n \geq m$ or $m \leq n$, iff we have $n = m + a$ for some natural number a . We say that n is strictly greater than m , and write $n > m$ or $m < n$, iff $n \geq m$ and $n \neq m$.*

Lemma 2.2.11.a. *For any natural number n , $n++ > n$.*

Proof. For any natural number, we know that $n++ = n + 1$ from Corollary 2.2.3.a. Then using the fact that $n \neq n++$ from Lemma 2.1.8.a, we can conclude $n++ > n$ by Definition 2.2.11. □

Proposition 2.2.12. (Basic properties of order for natural numbers). *Let a, b, c be natural numbers. Then*

Proposition 2.2.12.a. (Order is reflexive) $a \geq a$.

Proof. Any natural number a can be represented $a = a + 0$ using Lemma 2.2.2 what proves by Definition 2.2.11 that $a \geq a$. $\square$

Proposition 2.2.12.b. (Order is transitive) *If $a \geq b$ and $b \geq c$, then $a \geq c$.*

Proof. Given Definition 2.2.11 and the propositional assumption, b can be represented as $b + \tilde{c}$ and a can be represented as $b + \tilde{b}$ where $\tilde{b}$ and $\tilde{c}$ are natural numbers. Then combining both representations, $a = b + \tilde{b} = (b + \tilde{c}) + \tilde{b} = b + (\tilde{c} + \tilde{b})$ using Proposition 2.2.5. $\tilde{c} + \tilde{b}$ is a natural number by Proposition 2.2.1.a what leads us to $a \geq c$ by Definition 2.2.11. $\square$

Proposition 2.2.12.c. (Order is antisymmetric) $a \geq b$ and $b \geq a$ iff $a = b$.

(Original) *if $a \geq b$ and $b \geq a$, then $a = b$.*

Proof. The forward pass: by the propositional assumption and Definition 2.2.11, $a = b + \tilde{b}$ and $b = a + \tilde{a}$. Combining both terms, $a = b + \tilde{a} = (a + \tilde{a}) + \tilde{a}$ what equals to $a + (\tilde{a} + \tilde{a})$ using Proposition 2.2.5. Recalling Cancellation Law from Proposition 2.2.6, $(\tilde{a} + \tilde{a})$ must be equal to 0 since $a = a + 0$ by Definition 2.2.1. By Corollary 2.2.9, $\tilde{a} = 0$. Plugging it back to $b = a + \tilde{a} = a + 0 = a$.

The reverse pass: if $a = b$, then $a \geq b$ and $b \geq a$. If $a = b$, we can use the reflexivity of order from Proposition 2.2.12.a to transform $a \geq a$ and $b \geq b$ to $a \geq b$ and $b \geq a$ respectively. That closes the proof in both directions. $\square$

Proposition 2.2.12.d. (Addition preserves order) $a \geq b$ if and only if $a + c \geq b + c$.

Proof. Let's start with forward case: if $a \geq b$, then $a + c \geq b + c$. If $a \geq b$, then we can represent $a = b + \tilde{b}$ by Definition 2.2.11. Plugging the expression into $a + c$ results in $(b + \tilde{b}) + c$. Using Proposition 2.2.5 and Proposition 2.2.4, we can transform the expression into $(b + c) + \tilde{b}$ what proves $a + c \geq b + c$ by Definition 2.2.11.

Now backward: if $a + c \geq b + c$, then $a + c = b + c + \tilde{b}$ by Definition 2.2.11. Then using Cancellation Law from Proposition 2.2.6, $a = b + \tilde{b}$ which is $a \geq b$ by Definition 2.2.11. $\square$

Proposition 2.2.12.e. $a < b$ if and only if $b = a + (d++)$ for any natural number d .

Proof. For the forward case we need to prove that if $a < b$, then $b = a + d$ and $d \neq 0$. If $a < b$, then $b = a + d$ and $b \neq a$ by Definition 2.2.11. Let's assume that d is not a

positive natural number (i.e. $d = 0$). Then $a + d = a + 0 = a \stackrel{\neq}{=} b$ by the antecedent $b \neq a$.

For the backward pass: $b = a + (d++) = (a++) + d$ by Lemma 2.2.3 and Definition 2.2.1. If $a = b$, then $b \neq (a++)$ by Axiom 2.4. If $b \neq (a++)$ and $b = (a++) + d$, then $b > (a++)$ by Definition 2.2.11. If $b > (a++)$, then $b > a$ by Lemma 2.2.11.a and Proposition 2.2.12.b. If $a \neq b$ and $b = a + (d++)$, then $b > a$ by Definition 2.2.11. $\square$

Proposition 2.2.12.f. $a < b$ if and only if $a++ \leq b$.

Proof. For forward case, we need to prove if $a++ \leq b$, then $a < b$. Combining Definition 2.2.11 and associativity from Proposition 2.2.5, we can infer $b = (a++) + d = a + (d++)$ where $d++$ is a positive natural number by Axiom 2.3 and Definition 2.2.7. Thus, we can conclude that $b > a$ by Proposition 2.2.12.e.

For the backward: if $a++ \leq b$, then $b = (a++) + d = a + (d++)$ by Proposition 2.2.4. Since $d++ \neq 0$ by Axiom 2.3, then $d++$ is a positive number. If $d++$ is a positive number, then $a < b$ by Proposition 2.2.12.e. $\square$

Lemma 2.2.12.g. $0 \leq b$ for all natural numbers.

Proof. b can be represented as $b = 0 + b$ by Definition 2.2.1, what makes it $0 \leq b$ by Definition 2.2.11. $\square$

Proposition 2.2.13. (Trichotomy of order for natural numbers). *Let a and b be natural numbers. Then exactly one of the following statements is true: $a < b, a = b, a > b$.*

Proof. Let's start that **no more than one** statement holds at a time. If $a = b$, then neither $a < b$ nor $a > b$ holds by Definition 2.2.11. If $a > b$ and $a < b$, then by Definition 2.2.11, $a \geq b, a \leq b$ and $a \neq b$. If $a \geq b$ and $a \leq b$, then $a \stackrel{\neq}{=} b$ using Proposition 2.2.12.c, what contradicts the antecedent. Thus, no more than one statement holds at a time.

Now let's prove that there is **at least one** statement holds inducting on a . For the base case, $0 \leq b$ by Lemma 2.2.12.g. For the step case, suppose that one of the trichotomy statements holds for a . We need to prove whether any holds for $a++$. If $a < b$, then $a++ \leq b$ by Proposition 2.2.12.f. If $a = b$, then $a++ > b$ by Lemma 2.2.11.a. If $a > b$, then $a = b + (d++)$ by Proposition 2.2.12.e. Then we can take the successor $a++ = (b + (d++))++$ by Axiom 2.4, what equals to $b + (d++)++$ by Definition 2.2.1. $(d++)++$ is a positive number by Definition 2.2.7 and Axiom 2.3. Thus, $a++ > b$ still holds by Proposition 2.2.12.e. That covers all possible cases and closes the induction. $\square$

Proposition 2.2.14. (Strong principle of induction). *Let m_0 be a natural number, and let $P(m)$ be a property pertaining to an arbitrary natural number m . Suppose that for each $n \geq m_0$, we have the following implication: if $P(m)$ is true for all natural numbers $m_0 \leq m < n$, then $P(n)$ is also true (in particular, this means that $P(m_0)$ is true, since in this case the hypothesis is vacuous). Then we can conclude that $P(n)$ is true for all natural numbers $n \geq m_0$.*

Proof. We'll induct on n . For the base case m_0 : using Definition 2.2.11, $P(m)$ is true for $m_0 \leq m \leq m_0$ and $m \neq m_0$. If $m_0 \leq m$ and $m_0 \geq m$, then $m_0 \stackrel{!}{=} m$ by Proposition 2.2.12.c, what contradicts the fact that $m \neq m_0$. That means that there are no such m that satisfy $m_0 \leq m < m_0$. Thus, $P(m)$ is vacuously true for $m_0 \leq m < m_0$.

For the step case, suppose $P(m)$ for $m_0 \leq m < n \implies P(n)$, we need to prove $P(m)$ for $m_0 \leq m < n++ \implies P(n++)$. Since $P(m)$ holds for $m_0 \leq m < n$ and for $m = n$ (due to the given implication on $P(n)$), $P(m)$ holds for $m_0 \leq m \leq n$. The latter term is equivalent to $m_0 \leq m < n++$ by Proposition 2.2.12.f. **That means $P(m)$ is true for $m_0 \leq m < n++$, and it means $P(n++)$ is true. That closes the induction.** $\square$

Proposition 2.2.15. (Principle of backwards induction). *Let n be a natural number, and let $P(m)$ be a property pertaining to the natural numbers such that whenever $P(m++)$ is true, then $P(m)$ is true. Suppose that $P(n)$ is true. Prove that $P(m)$ is true for all natural numbers $m \leq n$.*

Proof. Let's induct on n . If $n = 0$, then $m \leq 0$. Combining with Lemma 2.2.12.g, we can conclude that $m = 0$ by Proposition 2.2.12.c. Since $P(n)$ and $n = m$ are true, we proved $P(m)$ for $m \leq n$.

For the step case, suppose the following implication holds: $P(n) \implies P(m)$ is true for $m \leq n$. Now we need to prove $P(n++) \implies P(m)$ is true for $m \leq n++$. Let's assume that $P(n++)$ is true, then $P(n)$ is also true by the "backward" property. If $P(n)$ is true, then $P(m)$ is true for $m \leq n$. Since $P(m)$ is true for $m \leq n$ and for $m = n++$ (by our assumption), then $P(m)$ is true for $m \leq n++$. That proves the implication and closes the induction. $\square$

Proposition 2.2.16. (Principle of induction starting from the base case n). *Let n be a natural number, and let $P(m)$ be a property pertaining to the natural numbers such that whenever $P(m)$ is true, $P(m++)$ is true. Show that if $P(n)$ is true, then $P(m)$ is true for all $m \geq n$.*

Proof. Since $m \geq n$, then we can represent $m = n + k$. So, let's induct on k . For the base case $k = 0$, $m = n$. Since $P(n)$ is true and $m = n$, then $P(m)$ is also true.

For the step case, we assume if $P(n)$ is true, then $P(n + k)$ is true for all $n + k \geq n$. Now we need to prove that the same holds for $P(n + (k++))$. Given the fact that $m = n + k$, $n + (k++) = (n + k)++$ by Definition 2.2.1, and the property assumption, if $P(m)$ is true, then $P(m++)$ is also true, we can conclude that $P(n + (k++))$ is true. That closes the induction. $\square$

2.3 Multiplication

Definition 2.3.1. (Multiplication of natural numbers). *Let m be a natural number. To multiply zero to m , we define $0 \times m := 0$. Now suppose inductively that we have defined how to multiply n to m . Then we can multiply $n++$ to m by defining $(n++) \times m := (n \times m) + m$.*

Proposition 2.3.1.a. *Product of two natural numbers is a natural number.*

Proof. Let's induct on n . The base case would be $n := 0$. By definition 2.3.1, $0 \times m := 0$ which is a natural number. Now suppose $n \times m$ is a natural number. Then $(n++) \times m := (n \times m) + m$ by Definition 2.3.1. Since $(n \times m)$ is a natural number by the inductive hypothesis and m is a natural number, the sum of two natural numbers is also a natural number by Proposition 2.2.1.a. That closes the induction. $\square$

Lemma 2.3.1.b. $1 \times n = n$ for all natural numbers.

Proof.

$$\begin{aligned}
 1 \times n &= (0++)n && \text{By Definition 2.1.3} \\
 &= (0 \times n) + n && \text{By Definition 2.3.1} \\
 &= 0 + n && \text{By Definition 2.3.1} \\
 &= n && \text{By Definition 2.2.1}
 \end{aligned}$$

$\square$

Lemma 2.3.2.a. *For any natural number n , $n \times 0 = 0$.*

Proof. Let's induct on n . For the base case $n := 0$, $n \times 0 = 0 \times 0 := 0$ by Definition 2.3.1. Now for the step case, let's assume that $n \times 0 = 0$. Then

$$\begin{aligned}
 (n++) \times 0 &:= (n \times 0) + 0 && \text{Definition 2.3.1} \\
 &= 0 + 0 && \text{Inductive Hypothesis} \\
 &= 0 && \text{Definition 2.2.1}
 \end{aligned}$$

what closes the induction. $\square$

Lemma 2.3.2.b. *For any natural numbers n, m , $m \times (n++) = (m \times n) + m$.*

Proof. Let's induct on m . For the base case,

$$\begin{aligned} m \times (n++) &= 0 \times (n++) \\ &= 0 \\ &= (0 \times n) + 0 \end{aligned}$$

For the step case, we assume that $m \times (n++) = (m \times n) + m$. Now we need to prove that $(m++) \times (n++) = ((m++) \times n) + (m++)$.

$$\begin{aligned} (m++) \times (n++) &= (m \times (n++)) + (n++) && \text{Definition 2.3.1} \\ &= ((m \times n) + m) + (n++) && \text{Inductive Hypothesis} \\ &= (m \times n) + (m + (n++)) && \text{Proposition 2.2.5} \\ &= (m \times n) + ((n++) + m) && \text{Proposition 2.2.4} \\ &= (m \times n) + n + (m++) && \text{Proposition 2.2.3} \\ &= ((m \times n) + n) + (m++) && \text{Proposition 2.2.5} \\ &= ((m++) \times n) + (m++) && \text{Definition 2.3.1} \end{aligned}$$

that closes the induction. □

Lemma 2.3.2.c. (Multiplication is commutative). *Let n, m be natural numbers. Then $n \times m = m \times n$.*

Proof. Let's induct on n . For the base case $n := 0$,

$$\begin{aligned} n \times m &= 0 \times m \\ &:= 0 && \text{Definition 2.3.1} \\ &= m \times 0 && \text{Lemma 2.3.2.a} \end{aligned}$$

For the step case, let's assume that $n \times m = m \times n$. Then we need to prove that $(n++) \times m = m \times (n++)$

$$\begin{aligned} (n++) \times m &:= (n \times m) + m && \text{Definition 2.3.1} \\ &= (m \times n) + m && \text{Inductive Hypothesis} \\ &= m \times (n++) && \text{Lemma 2.3.2.b} \end{aligned}$$

what closes the induction. □

Lemma 2.3.3. (Natural numbers have no zero divisors). *Let n, m be natural numbers. Then $n \times m = 0$ if and only if at least one of n, m is equal to zero. In particular, if n and m are both positive, then nm is also positive.*

Proof. We start with the left to right case. If $n \times m = 0$, then we need to prove that either n or m is zero. Let's assume than neither n nor m is zero (i.e. we can use $n++$ and $m++$ instead). Then $(n++) \times (m++) = (n \times (m++)) + m++$ (or $(m++) + (n++) = (m \times (n++)) + (n++)$) by Definition 2.3.1. We can conclude that in both cases, adding a natural number with a positive natural number will result in positive number by Proposition 2.2.8. That contradicts to $(n++) \times (m++) \stackrel{!}{=} 0$. Thus, either n or m must be zero.

The right to left case: if either n or m is zero, then $n \times m = 0 \times m := 0$ by Definition 2.3.1 or $n \times m = n \times 0 = 0 \times n = 0$ by Lemma 2.3.2.a. What closes both directions.

If both, n and m , are positive natural numbers (i.e. $n++$ and $m++$), then $(n++) \times (m++)$ is also positive as shown above. $\square$

Proposition 2.3.4. (Distributive law). *For any natural numbers a, b, c , we have $a(b + c) = ab + ac$ and $(b + c)a = ba + ca$.*

Proof. The case $a(b + c) = ab + ac$. Let's induct on a . The base case,

$$0(b + c) = 0b + 0c$$

$$0 = 0 + 0$$

Definition 2.3.1

$$0 = 0$$

Definition 2.2.1

The step case, we assume that $a(b + c) = ab + ac$ to prove $a++(b + c) = (a++)b + (a++)c$

$$a++(b + c) = a(b + c) + (b + c)$$

Definition 2.3.1

$$= ab + ac + b + c$$

Inductive Hypothesis

$$= ab + b + ac + c$$

Propositions 2.2.4 and 2.2.5

$$= (a++)b + (a++)c$$

Definition 2.3.1

what closes the induction for the former case. The case $(b + c)a = ba + ca$. Again, inducting on a . For the base case:

$$(b + c)0 = b0 + c0$$

$$0 = 0 + 0$$

Lemma 2.3.2.a

$$0 = 0$$

Lemma 2.2.2

For the step case, assuming that $(b + c)a = ba + ca$, we need to prove $(b + c)(a++) =$

$$b(a++) + c(a++)$$

$$\begin{aligned}
(b+c)(a++) &= (b+c)a + (b+c) && \text{Lemma 2.3.2.b} \\
&= ba + ca + b + c && \text{Inductive Hypothesis} \\
&= ba + b + ca + c && \text{Proposition 2.2.4 and 2.2.5} \\
&= b(a++) + c(a++) && \text{Lemma 2.3.2.b}
\end{aligned}$$

what closes the induction for the latter case. $\square$

Proposition 2.3.5. (Multiplication is associative). *For any natural numbers a, b, c we have $(a \times b) \times c = a \times (b \times c)$.*

Proof. Let's induct on a . The base case:

$$\begin{aligned}
(0 \times b) \times c &= 0 \times (b \times c) \\
0 \times c &= 0 && \text{By Definition 2.3.1 and Proposition 2.3.1.a} \\
0 &= 0 && \text{By Definition 2.3.1}
\end{aligned}$$

For the step case, we need to prove that $((a++) \times b) \times c = (a++) \times (b \times c)$, given $(a \times b) \times c = a \times (b \times c)$:

$$\begin{aligned}
((a++) \times b) \times c &= ((a \times b) + b) \times c && \text{By Definition 2.3.1} \\
&= (a \times b) \times c + (b \times c) && \text{By Proposition 2.3.4} \\
&= a \times (b \times c) + (b \times c) && \text{By the Inductive Hypothesis} \\
&= (a++) \times (b \times c) && \text{By Definition 2.3.1}
\end{aligned}$$

what closes the induction. $\square$

Proposition 2.3.6. (Multiplication preserves the order). *If a, b are natural numbers such that $a < b$, and c is positive, then $ac < bc$.*

Proof. Since c is positive, we can denote it as $\bar{c}++$. Let's induct on $\bar{c}$. The base case:

$$\begin{aligned}
a &< b \\
0 + a &< 0 + b && \text{By Definition 2.2.1 and Proposition 2.2.12.d} \\
(a \times 0) + a &< (b \times 0) + b && \text{By Definition 2.3.1} \\
a(0++) &< b(0++) && \text{By Definition 2.3.1 and Lemma 2.3.2.c} \\
a(\bar{c}++) &< b(\bar{c}++)
\end{aligned}$$

For the step case, given $a(\bar{c}++) < b(\bar{c}++)$, we need to prove that $a((\bar{c}++)++) < b((\bar{c}++)++)$:

$$\begin{array}{ll}
a(\bar{c}++) < b(\bar{c}++) & \text{By the Inductive Hypothesis} \\
a(\bar{c}++) + \bar{c}++ < b(\bar{c}++) + \bar{c}++ & \text{By Proposition 2.2.12.d} \\
a((\bar{c}++)++) < b((\bar{c}++)++) & \text{By Lemma 2.3.2.c and Definition 2.3.1}
\end{array}$$

□

Corollary 2.3.7. (Cancellation law). *Let a, b, c be natural numbers such that $ac = bc$ and $c \neq 0$. Then $a = b$.*

Proof. Let's apply the contrapositive property of implication: if $a \neq b$ or, in other words, $a > b$ exclusively or $a < b$, then $ac \neq bc$ or $c = 0$. If $c \neq 0$ and $a > b$, then $ac \leq bc$ by Proposition 2.3.6, which means $ac \neq bc$ by Proposition 2.2.13. Since Proposition 2.2.13 requires $c = 0$, we need to consider $c = 0$. If $c = 0$, then the consequent is trivially satisfied. □

Proposition 2.3.8. (Euclidean algorithm). *Let n be a natural number, and let q be a positive number. Then there exist natural numbers m, r such that $0 \leq r < q$ and $n = mq + r$.*

Proof. Now, we fix q and induct on n . For the base case $n = 0$, we have $0 = mq + r$. To satisfy the statement, both terms of RHS must be equal to zero. Indeed, there exist $m = 0$ and $r = 0$ that satisfy the statement and do not violate the condition $0 \leq r < q$. For the step case, we assume that the implication holds for n , and we need to prove that it holds for $n++$.

$$\begin{array}{ll}
n++ = (mq + r)++ & \text{Inductive Hypothesis} \\
= mq + (r++) & \text{By Proposition 2.2.4 and Definition 2.2.1}
\end{array}$$

if $r++ < q$, then $0 \leq r < r++ < q$ by Lemma 2.2.11.a and Proposition 2.2.12.b. If $r++ = q$, then $mq + (r++) = mq + q = (m++)q + 0 = \bar{m}q + \bar{r}$ by Definition 2.3.1. The case $r++ > q$ is not relevant due to Proposition 2.2.12.f. □

Definition 2.3.9. (Exponentiation for natural numbers). *Let m be a natural number. To raise m to the power 0, we define $m^0 := 1$. Now suppose recursively that m^n has been defined for some natural number n , then we define $m^{n++} := m^n \times m$.*

Proposition 2.3.10. $(a + b)^2 = (a^2 + 2ab + b^2)$ for all natural numbers a, b .

Proof.

$$\begin{aligned}(a+b)^2 &= (a+b)^{1++} \\ &= (a+b) \times (a+b) && \text{By Definition 2.3.9} \\ &= (a+b)a + (a+b)b && \text{By Proposition 2.3.4} \\ &= aa + ab + ab + bb && \text{By Proposition 2.3.4} \\ &= a^2 + (1 \times ab) + ab + b^2 && \text{By Definition 2.3.9 and Proposition 2.3.1.b} \\ &= a^2 + (1++)ab + b^2 && \text{By Definition 2.3.1} \\ &= a^2 + 2ab + b^2 && \text{By Definition 2.1.3}\end{aligned}$$

□

Chapter 3

Set Theory

3.1 Fundamentals

Axiom 3.1. (Sets are objects.) *If A is a set, then A is also an object. In particular, given two sets A and B , it is meaningful to ask whether A is also an element of B .*

Axiom 3.2. (Equality of sets). *Two sets A and B are equal, $A = B$, iff every element of A is an element of B and vice versa. To put it another way, $A = B$ if and only if every element x of A belongs also to B , and every element y of B belongs also to A .*

Axiom 3.3. (Empty set). *There exists a set $\emptyset$ or $\{\}$, known as the empty set, which contains no elements, i.e. for every object x we have $x \notin \emptyset$.*

Proposition 3.1.1. *There is exactly one empty set $\emptyset$.*

Proof. Let us assume that $\emptyset'$ is also an empty set. Based on Axiom 3.3, since no elements are in both sets, then all elements (in particular, the lack of them), that are in $\emptyset$, are also in $\emptyset'$ and vice versa. That makes $\emptyset' = \emptyset$. $\square$

Lemma 3.1.5. (Single choice). *Let A be a non-empty set. Then there exists an object x such that $x \in A$.*

Proof. Let us prove by contradiction. If A is a non-empty set, then no x exists such that $x \in A$. Or in other words, for every object x we have $x \notin A$. But this is the empty set by Axiom 3.3. $\square$

Axiom 3.4. (Singleton sets and pair sets). *If a is an object, then there exists a set $\{a\}$ whose only element is a , i.e., for every object y , we have $y \in \{a\}$ if and only if $y = a$; we refer to $\{a\}$ as the singleton set whose element is a . Furthermore, if a and b are objects, then there exists a set $\{a, b\}$ whose only elements are a and b ; i.e., for every object y , we have $y \in \{a, b\}$ if and only if $y = a$ or $y = b$; we refer to this set as the pair set formed by a and b .*

Proposition 3.1.6. *There is only one singleton set for each object a .*

Proof. Let's assume that there's another singleton set a' for a different from a which is a singleton set of a . Since only $a \in a'$ by Axiom 3.4, then for all $x \in a$ it is also $x \in a'$. What makes $a = a'$ by Axiom 3.2 what contradicts our initial assumption. $\square$

Proposition 3.1.7. *Given two objects a and b , there is only one pair set formed by a and b .*

Appendix A

The basics of mathematical logic

A.1 Mathematical statements

Exercise A.1.1. *What is the negation of the statement “either X is true or Y is true, but not both”?*

Solution. X and Y are false or X and Y are true. $\square$

Exercise A.1.2. *What is the negation of the statement “ X is true if and only if Y is true”?*

Solution. X is false iff Y is false $\square$

Exercise A.1.3. *Suppose that you have shown that whenever X is true, then Y is true, and whenever X is false, then Y is false. Have you demonstrated that X and Y are logically equivalent? Explain.*

Solution. A well-defined statement can be either true or false only (all possible values). Now, we consider the assumptions: (1) if X is false, then Y is false (Y is never true in that case), and (2) if X is true, then Y is true (Y is never false here). Thus, both statements coincide in all possible values what makes them “equivalent”. $\square$

Exercise A.1.4. *Suppose that you have shown that whenever X is true, then Y is true, and whenever Y is false, then X is false. Have you demonstrated that $X \Leftrightarrow Y$? Explain.*

Solution. No. If X is true, then Y is true: we exclude only the case X is true and Y is false. If Y is false, then X is false: we exclude the case X is true and Y is false. The both cases represent the same statement. As the result, the case X is False and Y is True is missing. Thus, X and Y are not being proven to be logically equivalent. $\square$

Exercise A.1.5. *Suppose that you have shown that X is true iff Y is true, and you know that Y is true iff Z is true. Is this enough to show that X, Y, Z are logically equivalent? Explain.*

Solution. Yes. Let’s construct a logical table of all possible cases:

X	Y	Z	Possible?	If not, why?
False	False	False	Yes	
False	False	True	No	$Y \Leftrightarrow Z$
False	True	False	No	$X \Leftrightarrow Y$ or $Y \Leftrightarrow X$
False	True	True	No	$X \Leftrightarrow Y$
True	False	False	No	$X \Leftrightarrow Y$
True	False	True	No	$X \Leftrightarrow Y$ or $Y \Leftrightarrow X$
True	True	False	No	$Y \Leftrightarrow Z$
True	True	True	Yes	

What results X, Y, Z to be logically equivalent by definition. $\square$

Exercise A.1.6. Suppose that you have shown that whenever X is true, then Y is true; that if Y is true, then Z is true; and if Z is true, then X is true. Is this enough to show that X, Y, Z are logically equivalent? Explain.

Solution. Yes. Let's construct a logical table of all possible cases:

X	Y	Z	Possible?	If not, why?
False	False	False	Yes	
False	False	True	No	$Y \implies Z$
False	True	False	No	$X \implies Y$
False	True	True	No	$Z \implies X$
True	False	False	No	$X \implies Y$
True	False	True	No	$X \implies Y$
True	True	False	No	$Y \implies Z$
True	True	True	Yes	

What results X, Y, Z to be logically equivalent by definition. $\square$

A.2 Implication

The main ideas in the chapter are the following:

- Vacuous hypothesis is a hypothesis where an antecedent is false. Since false implies false/true is always true, it does not provide any new information.

- We can also draw the following conclusions:

$$\begin{array}{ll}
 X \implies Y \iff \neg Y \implies \neg X & \text{Contrapositive} \\
 X \implies Y \iff Y \implies X & \text{Converse} \\
 X \implies Y \iff \neg X \implies \neg Y & \text{Inverse}
 \end{array}$$

- Although, vacuous hypothesis can be viewed as “useless”, it can still be used for a proof by contradiction using Contrapositive property, for instance. For that case, we need to have a consequent of the original implication to be known to be false.

A.3 The structure of proofs

Appendix B

The decimal system