

The pretty pretty pretty Priddy good complex

Mark E. Walker

January 23, 2026

Abstract

These are notes for the RTG learning seminar on Koszul algebras. Nothing here is original to me. I thank Eloísa Grifo and Adam LaClair for comments that changed these notes, usually for the better. *Dicatus melioribus mathematicis.*

My goals in this talk are to introduce quadratic algebras, Koszul algebras, quadratic duality and the Priddy complex, with an emphasis on examples.

See here for an explanation of the title.

1 Quadratic Algebras

Throughout k is a field and V is a finite dimensional k -vector space with basis x_1, \dots, x_n . The tensor algebra is the graded k -algebra

$$T(V) = \bigoplus_{j \geq 0} T^j(V) = \bigoplus_{j \geq 0} \overbrace{V \otimes_k \cdots \otimes_k V}^j = k\langle x_1, \dots, x_n \rangle,$$

the polynomial ring in n *non-commuting* variables. By a k -algebra we will always mean a quotient of $T(V)$ by a two-sided, homogeneous ideal $I = I_2 \oplus I_3 \oplus \cdots$:

$$A = T(V)/I.$$

So, A is an \mathbb{N} -graded k -algebra, with $A_0 = k$ and $A_1 = V$, that is generated as a k -algebra by $A_1 = V$. Note that we do *not* assume A is commutative.

Remark 1. Even though people in this room are mostly interested in commutative algebras, the quadratic dual of a commutative quadratic algebra is nearly always non-commutative.

We say A is *quadratic* if I is generated (as a two sided ideal) by its degree two part I_2 ; that is, if there is some vector subspace

$$W \subseteq V \otimes_k V = k\langle x_1, \dots, x_n \rangle_2$$

(which will be I_2) such that

$$A = T(V)/\langle W \rangle.$$

Here $\langle W \rangle$ is the smallest *two sided* ideal containing W . So the degree three part of $\langle W \rangle$ is

$$\langle W \rangle_3 = V \otimes_k W + W \otimes_k V \subseteq T^3(V),$$

the degree four part is

$$\langle W \rangle_4 = V \otimes_k V \otimes_k W + V \otimes_k W \otimes_k V + W \otimes_k V \otimes_k V \subseteq T^4(V),$$

and in general the degree d part is

$$\langle W \rangle_d = \sum_{i=0}^{d-2} T^i(V) \otimes_k W \otimes_k T^{d-i-2}(V) \subseteq T^d(V) = k\langle x_1, \dots, x_n \rangle_d.$$

The following examples will be referred to throughout these notes.

Example (Symmetric algebra). Let

$$W = k \cdot \{v \otimes v' - v' \otimes v \mid v, v' \in V\} = k \cdot \{x_i x_j - x_j x_i\} \subseteq V \otimes_k V.$$

We may identify W with $\Lambda^2(V)$; in general, $\Lambda^j(V)$ may be viewed as a subspace of $T^j(V)$ via the anti-symmetrization map; see (3.1) below. Then

$$T(V)/\langle W \rangle = S(V) = k[x_1, \dots, x_n],$$

the symmetric algebra on V , aka the usual polynomial ring in n commuting variables.

Example (Exterior algebra). Let

$$W = k\{v \otimes v' + v' \otimes v, v \otimes v \mid v, v' \in V\} \subseteq V \otimes_k V.$$

In other words, $W = \Gamma^2(V) = T^2(V)^{\Sigma_2}$, the subspace of $T^2(V)$ invariant under the evident action of Σ_2 . Since

$$v \otimes v' + v' \otimes v = (v + v') \otimes (v + v') - v \otimes v - v' \otimes v',$$

the first set of generators is actually redundant. Then

$$T(V)/\langle W \rangle = \Lambda(V) = \Lambda(x_1, \dots, x_n)$$

the exterior algebra on V .

Example (W is Zero). Taking things to an extreme: If $W = 0$ then $A = T(V) = k\langle x_1, \dots, x_n \rangle$ is the tensor algebra.

Example (W is Everything). Taking things to the opposite extreme: If $W = V \otimes_k V$ then $A = T(V)/\langle W \rangle = k \oplus V$ with (nearly) trivial multiplication: $v \cdot v' = 0$ for $v, v' \in V$ (and in general $(a, v) \cdot (a', v') = (aa', av' + a'v)$).

Example (Toy example). The commutative algebra $A = k[x, y]/(x^2, y^2)$ is quadratic with $W = k \cdot \{x^2, xy - yx, y^2\} \subseteq k\langle x, y \rangle$.

Example (Toy example dual). Say $W = k \cdot \{xy + yx\} \subseteq k\langle x, y \rangle$. Then $A = k\langle x, y \rangle/\langle W \rangle$ is the k -span of $\{x^i y^j\}$ but x and y anti-commute (and do not square to 0).

2 Koszul algebras

Let $A = \frac{T(V)}{I} = \frac{k\langle x_1, \dots, x_n \rangle}{I}$ be a k -algebra. We regard k as an (left) A -module by identifying it with A/\mathfrak{m}_A with $\mathfrak{m}_A := A_{\geq 1}$.

Definition 2. A k -algebra $A = T(V)/I$ is *Koszul* if the minimal graded free resolution of k (as a left A -module) is linear, in the sense that it has the form

$$\begin{array}{ccccccccccc} \cdots & \longrightarrow & W_j \otimes_k A(-j) & \longrightarrow & W_{j-1} \otimes_k A(-j+1) & \longrightarrow & \cdots & \longrightarrow & W_2 \otimes_k A(-2) & \longrightarrow & W_1 \otimes_k A(-1) & \longrightarrow & A & \longrightarrow & 0 \\ & & \parallel & & \parallel & & & & \parallel & & \parallel & & \parallel & & \\ \cdots & \longrightarrow & A(-j)^{b_j} & \longrightarrow & A(-j+1)^{b_{j-1}} & \longrightarrow & \cdots & \longrightarrow & A(-2)^{b_2} & \longrightarrow & A(-1)^{b_1} & \longrightarrow & A & \longrightarrow & 0 \end{array} \quad (2.1)$$

for some vector spaces W_1, W_2, \dots which are regarded to be of degree 0.

Another way of defining this concept: A is Koszul iff each differential in the minimal graded free resolution of k is represented by a matrix of linear forms (i.e., elements of A_1). Yet another way: A is Koszul if and only if $\text{Tor}_j^A(k, k)_i = 0$ unless $i = j$.

Proposition 3. *If A is Koszul then A is quadratic. In fact, so long as the minimal free resolution of k starts as $\cdots \rightarrow A(-2)^{b_2} \rightarrow A(-1)^{b_1} \rightarrow A$ then A is quadratic.*

Exercise 4. Prove the proposition. *Tips:* For $A = T(V)/I$, mess around with long exact sequences to prove that there are isomorphisms of graded k -vector spaces.

$$\text{Tor}_2^A(k, k) \cong \text{Ker}(\mathfrak{m}_A \otimes_k \mathfrak{m}_A \xrightarrow{\cdot} \mathfrak{m}_A) \cong I/\mathfrak{m}_{T(V)}I.$$

and hence $\dim_k \text{Tor}_2^A(k, k)_j$ is the number of minimal generators of I of degree j .

Remark 5. The converse of Proposition 3 is false. (Justification: We would not be running this seminar if the converse were true.)

3 Examples and non-examples of Koszul algebras

Here are some examples of W 's that give Koszul algebras:

Example (Symmetric algebra). The symmetric algebra $S(V) = k[x_1, \dots, x_n]$ is Koszul; see Example 1. We prove this by exhibiting a linear resolution.

In the beginning, God created the Koszul complex, and it was good.

The minimal free resolution of k is the Koszul¹ complex on x_1, \dots, x_n , which has the form

$$0 \rightarrow A(-n) \rightarrow A(-n+1)^n \rightarrow \cdots \rightarrow A(-j)^{\binom{n}{j}} \rightarrow \cdots \rightarrow A(-2)^{\binom{n}{2}} \rightarrow A(-1)^n \rightarrow A.$$

This is a linear resolution of k and so A is Koszul.

It will be helpful to exhibit this resolution in a basis-free way:

$$\begin{array}{cccccccccccccccc} 0 & \longrightarrow & \Lambda^n(V) \otimes_k S(V)(-n) & \longrightarrow & \cdots & \longrightarrow & \Lambda^j(V) \otimes_k S(V)(-j) & \longrightarrow & \cdots & \longrightarrow & \Lambda^2(V) \otimes_k S(V)(-2) & \longrightarrow & V \otimes_k S(V)(-1) & \longrightarrow & S(V) & \longrightarrow & 0. \\ & & \parallel & & & & \parallel & & & & \parallel & & \parallel & & \parallel & & \parallel & \\ 0 & \longrightarrow & \Lambda^n(S(V)(-1)) & \longrightarrow & \cdots & \longrightarrow & \Lambda^j(S(V)(-1))^n & \longrightarrow & \cdots & \longrightarrow & \Lambda^2(S(V)(-1))^n & \longrightarrow & S(V)(-1)^n & \longrightarrow & S(V) & \longrightarrow & 0. \end{array}$$

One way of describing the differential uses that we can realize $\Lambda^j(V)$ as a subspace of $T^j(V)$ by the “anti-symmetrization” map

$$v_1 \wedge \cdots \wedge v_j \mapsto \sum_{\sigma} \text{sign}(\sigma) v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(j)}. \quad (3.1)$$

Then the differential on the Koszul complex is the restriction of the “multiplication” map

$$T^j(V) \otimes_k S(V) \rightarrow T^{j-1}(V) \otimes_k S(V), \quad v_1 \otimes \cdots \otimes v_j \otimes f \mapsto v_1 \otimes \cdots \otimes v_{j-1} \otimes v_j \cdot f. \quad (3.2)$$

Exercise 6. Verify all the unproven claims made in the previous example.

Example (Exterior algebra). The exterior algebra $\Lambda(V)$ (see Example 1) is Koszul.

Let $\Gamma^j(V)$ be the “symmetric tensors”, that is, the subspace of $T^j(V)$ invariant under the evident action of the symmetric group Σ_j . E.g., $\Gamma^2(V)$ is the k -span of $\{v \otimes v' + v' \otimes v\} \cup \{v \otimes v\}$ (the first set is actually redundant). Notice that these are precisely the defining relations of the exterior algebra.

¹Not a coincidence that the same word appears!

Then the minimal free resolution of k is the infinite linear resolution

$$\cdots \rightarrow \Gamma^2(V) \otimes_k \Lambda(V)(-2) \rightarrow V \otimes_k \Lambda(V)(-1) \rightarrow \Lambda(V) \rightarrow k.$$

As before, the differential is the restriction to $\Gamma^j(V) \otimes_k \Lambda(V) \rightarrow \Gamma^{j-1}(V) \otimes_k \Lambda(V)$ of the multiplication map 3.2 in the previous example.

Example (W is zero). If $W = 0$ then $A = T(V)$. It is Koszul since

$$0 \rightarrow V \otimes_k T(V)(-1) \xrightarrow{\cdot} T(V) \rightarrow k$$

is exact. (Note that the projective dimension of k over $T(V)$ is finite — in fact, equal to 1 — to $T(V)$ is like a regular ring.)

Example (W is everything). $W = T^2(V)$. Then $A = k \oplus V$ with (nearly) trivial multiplication. This is Koszul since the minimal resolution of k is

$$\cdots \rightarrow V \otimes_k V \otimes_k V \otimes_k A \xrightarrow{v_1 \otimes v_2 \otimes v_3 \otimes a \mapsto v_1 \otimes v_2 \otimes v_3 a} V \otimes_k V \otimes_k A \xrightarrow{v_1 \otimes v_2 \otimes a \mapsto v_1 \otimes v_2 a} V \otimes_k A \xrightarrow{\cdot} A \rightarrow k \rightarrow 0.$$

Remark 7. Do you notice that the previous four examples occur in “dual pairs”?

That is, the “totalization” of the minimal resolution of k over $S(V)$ is $\Lambda(V) \otimes_k S(V)$, whereas the totalization of the minimal resolution of k over $\Lambda(V)$ is $\Gamma(V) \otimes_k \Lambda(V)$, and we have

$$(\Gamma(V) \otimes_k \Lambda(V))^* \cong \Lambda(V^*) \otimes S(V^*).$$

Likewise, the totalization of the minimal resolution of k over $k \oplus V$ is $T(V) \otimes_k (k \oplus V)$ whereas the minimal resolution of k over $T(V)$ is $(V \oplus k) \otimes T(V)$, which are similarly dual.

This is no fluke!

Example (Toy example). $A = k[x, y]/(x^2, y^2)$ is Koszul. This will be justified later.

Example (Toy example dual). $k\langle x, y \rangle / \langle xy + yx \rangle$ is Koszul. This will be justified later.

Some more examples:

Example 1. (McCullough-Peeva) Let $A = k[x, y, z]/(x^2, y^2, xy + xz, xy + yz)$; that is, take W to be the k -span of $xy - yx, xz - zx, yz - zy, x^2, y^2, xy + xz, xy + yz$. Then A is Koszul. This can be justified by the fact that its “Koszul dual” A^1 admits a quadratic Gröbner basis.

Example 2. (McCullough-Peeva) Here is a non-example:

$$A = k[x_1, x_2, x_3, x_4]/(x_1^2, x_2^2, x_3^2, x_4^2, x_1x_2 + x_1x_3 + x_1x_4),$$

so that W the k -span of

$$\{x_i x_j - x_j x_i\} \cup \{x_1^2, x_2^2, x_3^2, x_4^2, x_1x_2 + x_1x_3 + x_1x_4\}.$$

This can be shown to not be Koszul by using that if A were Koszul then we would have $P_A(t) \cdot H_A(-t) = 1$; this will be proven by a tentative Eloísa next week. But

$$1/H_A(-t) = \frac{1}{1 + 4t + 5t^2 + t^3} = 1 + 4t + 11t^2 + 25t^3 + 49t^4 + 82t^5 + 108t^6 + 71t^7 - 174t^8 + \cdots$$

cannot be the Poincaré series of k .

Example 3. (Conca) Here is another non-example: Let R be the commutative quadratic algebra

$$R = k[x, y, z, t]/(x^2, y^2, z^2, t^2, xy + zt).$$

(So W is the k -span of all the commutators $xy - yx, xz - zx, \dots$ along with $x^2, y^2, z^2, t^2, xy + zt$.) Then $b_{3,4}^R(k) = 5$ and hence R is not Koszul.

Exercise 8. Compute the quadratic dual of Conca’s example (once you learn what that is).

4 Quadratic duals

Recall that we write V^* for $\text{Hom}_k(V, k)$ and that x_1, \dots, x_n is a basis of V . By the *dual basis* of V^* we mean the basis x_1^*, \dots, x_n^* of V^* with $x_i^*(x_j) = \delta_{i,j}$. (Roughly, $x_i^* = \partial/\partial x_i$.)

The inclusion $W \subseteq V \otimes_k V$, induces a surjection by taking k -linear duals

$$V^* \otimes_k V^* \cong (V \otimes_k V)^* \twoheadrightarrow W^*$$

and we write the kernel of this surjection as $W^\perp \subseteq V^* \otimes_k V^*$. The *quadratic dual* of the quadratic algebra $A = T(V)/\langle W \rangle$ is the quadratic algebra

$$A^! = T(V^*)/(W^\perp).$$

Remark 9. We have identified $V^* \otimes_k V^*$ with $(V \otimes_k V)^*$, but there are multiple ways of doing so. Let's agree to use the following one: $\gamma_1 \otimes \gamma_2 \in V^* \otimes_k V^*$ corresponds to the map $V \otimes_k V \rightarrow k$ given by $v_1 \otimes v_2 \mapsto \gamma_1(v_2)\gamma_2(v_1)$. If we think of this as a pairing $V^* \otimes_k V^* \otimes_k V \otimes_k V \rightarrow k$, the convention we use is to combine adjacent terms: $\gamma_1 \otimes \gamma_2 \otimes v_1 \otimes v_2 \mapsto \gamma_2(v_1)\gamma_1 \otimes v_2 \mapsto \gamma_1(v_2)\gamma_2(v_1)$.

Example (Symmetric algebra). If $W = \Lambda^2(V)$, the k -span of $\{v_1 \otimes v_2 - v_2 \otimes v_1\}$, so that $A = S(V)$, then W^\perp is the k -span of $\{\gamma \otimes \gamma \mid \gamma \in V^*\}$. (Recall that this span includes $\gamma_1 \otimes \gamma_2 + \gamma_2 \otimes \gamma_1$.) Thus

$$S(V)^! = \Lambda(V^*).$$

Example (Exterior algebra). If W is the k -span of $\{v \otimes v \mid v \in V\}$, so that $A = \Lambda(V)$, then W^\perp is the k -span of $\{\gamma_1 \otimes \gamma_2 - \gamma_2 \otimes \gamma_1 \mid \gamma_1, \gamma_2 \in V^*\}$. Thus

$$\Lambda(V)^! = S(V^*).$$

This is an example of quadratic duality:

Proposition 10. For any quadratic algebra A , we have $(A^!)^! \cong A$. We call A and $A^!$ quadratic duals.

Exercise 11. Prove the previous Proposition.

Example (Toy example). $k[x, y]/(x^2, y^2)$ and $k\langle x^*, y^* \rangle / \langle x^*y^* + y^*x^* \rangle$ are quadratic duals.

Example (Extreme examples). The tensor algebra $T(V)$ and the trivial algebra $k \oplus V^*$ are quadratic duals:

$$T(V)^! = (k \oplus V^*) \text{ and } (k \otimes V)^! = T(V^*).$$

Here is another example:

Example 4. Let $A = k[x, y, z]/(x^2, y^2, xy + xz, xy + yz)$; that is, take

$$W = \text{the } k\text{-span of } \{xy - yx, xz - zx, yz - zy, x^2, y^2, xy + xz, xy + yz\}.$$

Then

$$A^! = k\langle x^*, y^*, z^* \rangle / (z^*z^*, x^*y^* + y^*x^* - x^*z^* - z^*x^* - y^*z^* - z^*y^*)$$

Exercise 12. Verify the previous example.

Exercise 13. Prove that if $\text{char}(k) \neq 2$ and $n \geq 2$, then for a quadratic algebra A , A and $A^!$ cannot both be commutative.

What is the story when $\text{char}(k) = 2$?

5 The pretty pretty pretty Priddy good complex

We come to the Priddy² complex of a quadratic algebra $A = T(V)/\langle W \rangle$. You should think of this as a generalization of the Koszul complex and indeed some authors refer to the Priddy complex as the “(generalized) Koszul complex”.

Let us note that the degree d component of $A^!$ is a quotient of $T^d(V^*)$ and so its dual $(A_d^!)^*$ can be identified with a subspace of $T^d(V)$. For instance $(A_1^!)^* = V$, $(A_2^!)^* = W$, and

$$(A_3^!)^* = W \otimes_k V \cap V \otimes_k W \subseteq T^3(V).$$

In general,

Lemma 14. *For any quadratic algebra $A = T(V)/\langle W \rangle$, we have*

$$(A_d^!)^* = \bigcap_i T^i(V) \otimes W \otimes T^{d-i-2}(V)$$

for each d , as subspaces of $T^d(V)$.

For example

$$(A_4^!)^* = (W \otimes_k V \otimes_k V) \cap (V \otimes_k W \otimes_k V) \cap (V \otimes_k V \otimes_k W) \subseteq T^4(V).$$

Definition 15. For a quadratic algebra $A = T(V)/\langle W \rangle$, the *Priddy complex* of A (also known as the *generalized Koszul complex*) is the complex

$$\cdots \rightarrow (A_j^!)^* \otimes_k A(-j) \rightarrow (A_{j-1}^!)^* \otimes_k A(-j+1) \rightarrow \cdots \rightarrow (A_3^!)^* \otimes_k A(-3) \rightarrow W \otimes_k A(-2) \rightarrow V \otimes_k A(-1) \rightarrow A \rightarrow 0.$$

where the differential is given by the restriction of the multiplication map

$$T^j(V) \otimes_k A(-j) \rightarrow T^{j-1}(V) \otimes_k A(-j+1), \quad v_1 \otimes \cdots \otimes v_j \otimes a \mapsto v_1 \otimes \cdots \otimes v_{j-1} \otimes v_j \cdot a.$$

Exercise 16. Verify that this really is a complex (i.e., that the differential squares to 0).

Example (Toy example). Recall that if $A = k\langle x, y \rangle / \langle xy + yx \rangle$ then $A^! = k[x^*, y^*] / (x^*x^*, y^*y^*) = k\langle x^*, y^* \rangle / \langle x^*x^*, y^*x^* - x^*y^*, y^*y^* \rangle$. So $A^!$ has k -basis $1, x^*, y^*, y^*x^* + x^*y^*$ and $(A^!)^*$ has k -basis $1, x, y, xy + yx$. So the priddy complex for A is

$$0 \rightarrow (xy + yx) \otimes A(-2) \rightarrow x \otimes A(-1) \oplus y \otimes A(-1) \rightarrow A \rightarrow 0 \quad (5.1)$$

with differential $(xy + yx) \otimes a \mapsto x \otimes ya + y \otimes xa$ etc. Using the evident bases, this is the complex

$$0 \rightarrow A(-2) \xrightarrow{\begin{bmatrix} y \\ x \end{bmatrix}} A(-1)^2 \xrightarrow{\begin{bmatrix} x & y \end{bmatrix}} A \rightarrow 0.$$

Exercise 17. Verify the details of the previous example.

Theorem 18. *A quadratic algebra A is Koszul if and only if the Priddy complex is a resolution of k .*

Example. The Priddy complex is a resolution of k for the symmetric algebra, the exterior algebra, the tensor algebra and the trivial algebra.

Example (Toy example). One³ may verify directly that the Priddy complex for $A = k\langle x, y \rangle / \langle xy + yx \rangle$ shown in (5.1) above is a resolution of k . So, A is Koszul.

²I knew Stuart Priddy back in the day

³i.e., you, dear reader

Proof of Theorem. Since the Priddy complex is clearly a linear complex, one direction is immediate: If the Priddy complex resolution k , then A is Koszul.

Assume A is Koszul. This means we have an exact sequence of graded A -modules of the form

$$\cdots \longrightarrow W_j \otimes_k A(-j) \longrightarrow \cdots \longrightarrow W_2 \otimes_k A(-2) \longrightarrow W_1 \otimes_k A(-1) \longrightarrow A \longrightarrow k \longrightarrow 0. \quad (5.2)$$

for some finite dimensional vector spaces W_j , $j \geq 1$, each viewed as being concentrated in degree zero. For any d , the degree d part of $W_j \otimes_k A(-j)$ is $W_j \otimes_k A_{d-j}$ and the degree d part of (5.2) is the exact sequences of finite dimensional k -vector spaces

$$0 \rightarrow W_d \rightarrow W_{d-1} \otimes_k A_1 \rightarrow W_{d-2} \otimes_k A_2 \rightarrow \cdots \rightarrow W_1 \otimes_k A_{d-1} \rightarrow A_d \rightarrow 0.$$

We can use this to recursively identify each W_d and what the maps are. In detail:

For $d = 1$ we get the exact sequence

$$0 \rightarrow W_1 \rightarrow A_1 \rightarrow 0,$$

and since $I_1 = 0$ we have $A_1 = V$ and thus (up to isomorphism) we deduce that

- $W_1 = V$ and
- the differential $W_1 \otimes_k A(-1) \rightarrow A$ is given by multiplication: $v \otimes a \mapsto v \cdot a$.

This confirms what we already knew: the minimal resolution of k starts as $V \otimes_k A(-1) \xrightarrow{\cdot} A$ or in other words as $A(-1)^n \xrightarrow{(x_1, \dots, x_n)} A$.

For $d = 2$ we get the exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & W_2 & \longrightarrow & W_1 \otimes_k A_1 & \longrightarrow & A_2 \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & W_2 & \longrightarrow & V \otimes V & \longrightarrow & \frac{V \otimes_k V}{W} \longrightarrow 0 \end{array}$$

and thus (up to isomorphism) we have

- $W_2 = W \subseteq T^2(V)$ and
- the differential $W_2 \otimes_k A(-2) \rightarrow W_1 \otimes_k A(-1)$ is multiplication, in the sense that it is the restriction of the map on $T^2(V) \otimes_k A(-2) \rightarrow V \otimes_k A(-1)$ given by $v_1 \otimes v_2 \otimes a \mapsto v_1 \otimes v_2 a$.

For $d = 3$ we get

$$\begin{array}{cccccccc} 0 & \longrightarrow & W_3 & \longrightarrow & W_2 \otimes_k A_1 & \longrightarrow & W_1 \otimes_k A_2 & \longrightarrow & A_3 \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & W_3 & \longrightarrow & W \otimes V & \longrightarrow & \frac{V \otimes_k V \otimes_k V}{V \otimes_k W} & \longrightarrow & \frac{V \otimes_k V \otimes_k V}{W \otimes_k V + V \otimes_k W} \longrightarrow 0 \end{array}$$

and thus (up to isomorphism) we have

- $W_3 = (W \otimes_k V \cap V \otimes_k W) = (A_3^!)^* \subseteq T^3(V)$ and
- the differential $W_3 \otimes_k A(-3) \rightarrow W_2 \otimes_k A(-2)$ is multiplication: $v_1 \otimes v_2 \otimes v_3 \otimes a \mapsto v_1 \otimes v_2 \otimes v_3 a$.

For $d = 4$ we get

$$\begin{array}{ccccccccccc}
0 & \longrightarrow & W_4 & \longrightarrow & W_3 \otimes_k A_1 & \longrightarrow & W_2 \otimes_k A_2 & \longrightarrow & W_1 \otimes_k A_3 & \longrightarrow & A_4 \longrightarrow 0 \\
& & \parallel & & \parallel & & \parallel & & \parallel & & \parallel \\
0 & \longrightarrow & W_4 & \longrightarrow & (W \otimes_k V \cap V \otimes_k W) \otimes V & \longrightarrow & W \otimes \frac{V \otimes_k V}{W} & \longrightarrow & \frac{V \otimes_k V \otimes_k V \otimes_k V}{V \otimes_k W \otimes_k V + V \otimes_k V \otimes_k W} & \longrightarrow & A_4 \longrightarrow 0
\end{array}$$

and thus (up to isomorphism) we have

- $W_4 = (W \otimes_k V \otimes_k V \cap V \otimes_k W \otimes_k V \cap V \otimes_k V \otimes_k W) = (A_4^!)^* \subseteq T^4(V)$ and
- the differential $W_4 \otimes_k A(-4) \rightarrow W_3 \otimes_k A(-3)$ is multiplication.

By the Freshman's Principle of Mathematical Induction, we have proven that (5.2) is isomorphic to the Priddy complex. \square

Corollary 19. *A quadratic algebra A is Koszul if and only if its quadratic dual $A^!$ is Koszul.*

Proof. This holds since the k -dual of the Priddy complex for A is the Priddy complex for $A^!$:

$$((A^!)^* \otimes_k A)^* \cong A^! \otimes_k A^* \cong A^* \otimes_k A^! \cong ((A^!)^!)^* \otimes_k A^!.$$

\square

Example (Toy example). $k[x, y]/(x^2, y^2)$ is Koszul since, as we saw above, its quadratic dual is.

Exercise 20. Write down the first few terms of the Priddy complex for $k[x, y]/(x^2, y^2)$.

Corollary 21. *If A is Koszul,*

$$\mathrm{Ext}_A^*(k, k) \cong A^!.$$

Proof. Since the Priddy complex $(A^!)^* \otimes_k A$ resolves k , we have

$$\mathrm{Ext}_A^*(k, k) = H^* \mathrm{Hom}_A((A^!)^* \otimes_k A, k) = H^* \mathrm{Hom}_k((A^!)^*, k) = A^!$$

as graded k -vector spaces. To complete the proof one should verify the multiplication rules match up; I leave that as an exercise. \square

Corollary 22. *If A is Koszul,*

$$A \cong \mathrm{Ext}_{\mathrm{Ext}_A^*(k, k)}^*(k, k).$$

Remark 23. The previous Corollary admits a vast generalization: If R is any complete local ring, or any reasonable graded ring, we have

$$R \sim \mathbb{R} \mathrm{Hom}_{\mathbb{R} \mathrm{Hom}_R(k, k)}(k, k).$$

If A is Koszul, then the dga $\mathbb{R} \mathrm{Hom}_A(k, k)$ is formal: it is quasi-isomorphic, as a dga, to its homology.

Remark 24. In general, $\mathrm{Ext}_A^*(k, k)$ is the k -linear dual of $\mathrm{Tor}_*^A(k, k)$. If A is commutative, $\mathrm{Tor}_*^A(k, k)$ is a graded commutative ring. So, when A is commutative both $\mathrm{Ext}_A^*(k, k)$ and its k -linear dual are algebras; in fact, $\mathrm{Ext}_A^*(k, k)$ is what is known as a Hopf algebra in this case.

So, if A is Koszul and commutative, $A^!$ is actually a Hopf algebra.

Exercise 25. Assume A is Koszul. Prove k has finite projective dimension over A if and only if $A^!$ is artinian.

Remark 26. If $pd_A(k) < \infty$ then every finitely generated graded A -module has finite projective dimension. This holds since $\mathrm{Tor}_*^A(M, k)$ can be computed by either resolving M or k . So if A is a Koszul algebra such that $A^!$ is artinian, then A is like a regular ring.

Exercise 27. Show that if A is commutative and Koszul and A^\dagger is artinian, then A is a polynomial ring.

Remark 28. For those who know about derived categories, if A is Koszul then $\text{Thick}_k(A) \cong \text{Perf}(A^\dagger)$ and $\text{Thick}_k(A^\dagger) \cong \text{Perf}(A)$. When $A = S(V)$ and $A^\dagger = \Lambda(V)$, this recovers the famed Bernstein-Gelfand-Gelfand correspondence: in this case $\text{Thick}_k(\Lambda(V)) = D^b(\Lambda(V))$ and $\text{Perf}(S(V)) = D^b(S(V))$.