

Math 342W/642/742W

Recitation – Day #11 (3.18.25)

Problem #1

Consider the following R code from the class demos and labs. The numbers on the right are line numbers that will be referred to later. They are not part of the code.

```
1 y = MASS::Cars93$Price
2 var(y)
3 [1] 93.30458
4 X = as.matrix(cbind(1, MASS::Cars93[,c(7,8,12,13,14,15,17,18)]))
5 n = nrow(X)
6 n
7 [1] 93
8 Xt = t(X)
9 XtX = Xt %*% X
10 XtXinv = solve(XtX)
11 XtXinvXt = XtXinv %*% Xt
12 b = XtXinvXt %*% y
13 H = X %*% XtXinvXt
14 I_minus_H = diag(n) - H
15 yhat = H %*% y
16 e = I_minus_H %*% y
17 var(e)
18      [,1]
19 [1,] 32.70302
```

(This problem is adapted from Problem #2 of Professor Kapelner's Spring 2024 Midterm I.)

- What is returned by R when evaluating `length(b)`?
- What is returned by R when evaluating `ncol(XtX)`?
- What is returned by R when evaluating `ncol(H)`?
- What is returned by R when evaluating `Matrix::rankMatrix(I_minus_H)`?
- Compute SST, SSE, SSR, and R^2 to the nearest two decimal places.
- What is the number of random predictors that can be added before the code throws an error and halts?

Problem #2

- (a) Assume that $\mathbf{X} \in \mathbb{R}^{n \times (p+1)}$ with $p > 1$ composed of random realizations from iid standard normal random variables, full rank. Let \mathbf{Q}, \mathbf{R} be the matrix results of the QR-decomposition procedure run on \mathbf{X} . Let $\mathbf{y} \in \mathbb{R}^n$ which represents a vector of measurements of a phenomenon of interest.

Prove that

$$\frac{\|\mathbf{Q}\mathbf{Q}^T\mathbf{y}\| + \|(I - \mathbf{Q}\mathbf{Q}^T)\mathbf{y}\|}{\|\mathbf{y}\|} \geq 1$$

-
- (b) Claim: $\exists c \neq 0$ such that $\mathbf{X}_{\cdot 3} = c\mathbf{Q}_{\cdot 3}$.
Show whether the claim is true or not.

(This problem is adapted from Problem #4 of Professor Kapelner's Spring 2024 Midterm I.)