

# FINITE ELEMENTS

NEAL COLEMAN<sup>†</sup>

ABSTRACT. Study the

## INTRODUCTION

### TRIANGLE

Let  $p, q, r \in \mathbb{R}^2$  be the vertices of a triangle  $T \subset \mathbb{R}^2$  with area  $A_T$ . Let  $\theta_p$  be the interior angle at vertex  $p$ , and likewise  $\theta_q, \theta_r$ . Let  $P$  be the length of the side opposite  $p$ , likewise with  $Q$  and  $R$ . (We may sometimes abuse notation and denote by  $P, Q, R$  the sides themselves.)

Area is computed by

$$\begin{aligned} A_T &= \frac{1}{2} \det \begin{bmatrix} p - r & q - r \end{bmatrix} \\ &= \frac{1}{2} \det \begin{bmatrix} p - q & r - q \end{bmatrix} \\ &= \frac{1}{2} \det \begin{bmatrix} q - p & r - p \end{bmatrix} \end{aligned}$$

### BARYCENTRIC COORDINATES

Let  $s = \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$ . Then we can write  $s = \alpha_p p + \alpha_q q + \alpha_r r$ , with  $\alpha_p + \alpha_q + \alpha_r = 1$ . Define

$$B = \left[ \begin{array}{c|c|c} p & q & r \\ \hline 1 & 1 & 1 \end{array} \right]$$

and

$$\alpha = \begin{bmatrix} \alpha_p \\ \alpha_q \\ \alpha_r \end{bmatrix}$$

so that  $s = B\alpha$ .

---

2000 *Mathematics Subject Classification.* Primary 51R10; Secondary 51R12.

<sup>†</sup> :)

Observe that

$$\begin{aligned}
\det B &= \det \begin{bmatrix} q & r \end{bmatrix} - \det \begin{bmatrix} p & r \end{bmatrix} + \det \begin{bmatrix} p & q \end{bmatrix} \\
&= (q_x r_y - q_y r_x) - (p_x r_y - p_y r_x) + (p_x q_y - p_y q_x) \\
&= p_x q_y - p_x r_y - r_x q_y - p_y q_x + p_y r_x + r_y q_x - r_x r_y + r_x r_y \\
&= (p_x - r_x)(q_y - r_y) - (p_y - r_y)(q_x - r_x) \\
&= 2A_T
\end{aligned}$$

Using cofactor expansion we calculate

$$B^{-1} = \frac{1}{2A_T} \begin{bmatrix} q_y - r_y & -(q_x - r_x) & \det \begin{bmatrix} q & r \end{bmatrix} \\ -(p_y - r_y) & p_x - r_x & -\det \begin{bmatrix} p & r \end{bmatrix} \\ p_y - q_y & -(p_x - q_x) & \det \begin{bmatrix} p & q \end{bmatrix} \end{bmatrix}$$

#### BUILDING BLOCKS OF ELEMENTS

Define  $v_p : T \rightarrow \mathbb{R}$  to be the restriction of the unique affine-linear function with  $v_p(p) = 1$ ,  $v_p(q) = 0$ ,  $v_p(r) = 0$ . Define  $v_q$  and  $v_r$  similarly.

In fact  $v_p(s)$  is the  $p$  coordinate in the barycentric expansion of  $s = (x, y)$ . (This is because linear functions are uniquely determined by their values.) If  $e_i$  is the  $i^{\text{th}}$  standard basis element of  $\mathbb{R}^3$  we have

$$v_p(s) = e_1^T B^{-1} s = \frac{1}{2A_T} \left( (q_y - r_y)x - (q_x - r_x)y + \det \begin{bmatrix} q & r \end{bmatrix} \right)$$

and its gradient is constant and equal to

$$\nabla v_p = \frac{1}{2A_T} \begin{bmatrix} q_y - r_y \\ -(q_x - r_x) \end{bmatrix}$$

**Integrals.** We now compute  $L^2$  norm of  $v_p$  and the  $L^2$  inner product of  $v_p, v_q$ .

To compute the squared norm  $\|v_p\|^2$  of  $v_p$ :

$$\|v_p\|^2 = \int_T (v_p)^2 = \int_0^1 t^2 |v_p^{-1}(t)| dt$$

Observing that  $|v_p^{-1}(t)| = (1-t)P$  we have

$$\begin{aligned}
\int_T (v_p)^2 &= \int_0^1 t^2 (1-t)P dt \\
&= P \left( \frac{1}{3} - \frac{1}{4} \right) \\
&= \frac{1}{12}P
\end{aligned}$$

Likewise  $\|v_q\|^2 = \frac{1}{12}Q$  and  $\|v_r\|^2 = \frac{1}{12}R$ .

To compute the inner product  $\langle v_p, v_q \rangle_2$  of  $v_p$  and  $v_q$  we foliate  $T$  by lines parallel to  $R$ , which are level sets of  $v_r$ .

$$\langle v_p, v_q \rangle = \int_0^1 \int_{v_r^{-1}(t)} v_p v_q dt$$

We use a linear parametrization of  $v_r^{-1}(t)$  with independent variable  $s$ , traversing from  $P$  when  $s = 0$  to  $Q$  when  $s = 1$ .

In abuse of notation re-use the symbols  $v_p, v_q$  for their restrictions to  $v_r^{-1}(t)$ . Then  $v_p(0) = 0$  and  $v_p(1) = (1-t)$ , while  $v_q(0) = 1-t$  and  $v_q(1) = 0$ . So  $v_p(s) = s(1-t)$  and  $v_q(s) = (1-s)(1-t)$ .

Thus

$$\int_{v_r^{-1}(t)} v_p v_q = \int_0^1 s(1-s)(1-t)^2 R ds = \frac{1}{6} R (1-t)^2$$

and so

$$\begin{aligned} \langle v_p, v_q \rangle_2 &= \int_0^1 \frac{1}{6} R (1-t)^2 dt \\ &= \frac{1}{18} R \end{aligned}$$

The  $L^2$  norm of  $\nabla v_p$  is nothing more than

$$\|\nabla v_p\|^2 = \frac{(q_y - r_y)^2 + (q_x - r_x)^2}{4A_T^2} = \frac{P^2}{4A_T^2}$$

The inner product of  $\nabla v_p$  and  $\nabla v_q$  is

$$\langle \nabla v_p, \nabla v_q \rangle = -\frac{1}{4A_T^2} \left( (q_y - r_y)(p_y - r_y) + (q_x - r_x)(p_x - r_x) \right) = -\frac{\langle q - r, p - r \rangle}{4A_T^2}$$

This can also be interpreted as proportional to the cosine of the interior angle at  $r$ :

$$\langle \nabla v_p, \nabla v_q \rangle = -\frac{1}{4A_T^2} PQ \cos \theta_r$$

## TRIANGULATIONS AND ELEMENTS

Let  $\Omega \subset \mathbb{R}^2$  be a precompact domain with piecewise linear boundary.

## COMPUTATIONS WITH TRIANGLES

Triangles admit a self-similar triangulation.

## PROOF OF WEYL'S LAW

## REFERENCES

- |         |  |
|---------|--|
| [Kai91] | Vadim A. Kaimanovich, <i>Poisson boundaries of random walks on discrete solvable groups</i> , Probability measures on groups, X (Oberwolfach, 1990), Plenum, New York, 1991, pp. 205–238. MR MR1178986 (94m:60014) |
| [Pra75] | J.-J. Prat, <i>Étude asymptotique et convergence angulaire du mouvement brownien sur une variété à courbure négative</i> , C. R. Acad. Sci. Paris Sér. A-B <b>280</b> (1975), no. 22, A1539–A1542.                 |

anovich:91

Prat75

Sullivan83
------------

[Sul83] D. Sullivan, *The Dirichlet problem at infinity for a negatively curved manifold*, no. 18, 723–732.

CHICAGOLAND

*Email address:* coleman.neal@gmail.com