

# Volatile Organic Compound Detection Using Insect Odorant-Receptor Functionalised Field-Effect Transistors

by

Eddyn Oswald Perkins Treacher

A thesis submitted in fulfilment of the  
requirements of the degree of  
Doctor of Philosophy in Physics  
School of Physical and Chemical Sciences  
Te Herenga Waka - Victoria University of Wellington

Apr 2023





# Acknowledgements

Thanks for all the fish.



# Abstract

This is a thesis skeleton written with quarto. Make a copy of this thesis repo and start to write!

Make a new paragraph by leaving a blank line.



# Table of contents

<b>Acknowledgements</b>	<b>1</b>
<b>Abstract</b>	<b>3</b>
<b>1 Introduction</b>	<b>7</b>
<b>2 Carbon Nanotube and Graphene Field-Effect Transistors</b>	<b>9</b>
2.1 Device Functionalisation . . . . .	9
2.2 Insect Odorant Receptors . . . . .	9
<b>3 Carbon Nanotube and Graphene Field-Effect Transistors as Biosensor Platforms</b>	<b>11</b>
<b>4 Fabrication</b>	<b>13</b>
<b>5 Functionalisation of Carbon Nanotubes and Graphene with Odorant Receptors</b>	<b>15</b>
5.1 Linker molecules . . . . .	15
5.1.1 1-Pyrenebutanoic acid N-hydroxysuccinimide ester (PBASE) . . .	15
<b>6 Results</b>	<b>19</b>
<b>7 Results</b>	<b>21</b>
<b>8 Summary</b>	<b>23</b>
<b>References</b>	<b>25</b>





# 1 Introduction

This is a book created from markdown and executable code.

See for additional discussion of literate programming.

[1] 2



## **2 Carbon Nanotube and Graphene Field-Effect Transistors**

### **2.1 Device Functionalisation**

### **2.2 Insect Odorant Receptors**



### **3 Carbon Nanotube and Graphene Field-Effect Transistors as Biosensor Platforms**



## 4 Fabrication

Stuff I did to get the results.





## 5 Functionalisation of Carbon Nanotubes and Graphene with Odorant Receptors

### 5.1 Linker molecules

#### 5.1.1 1-Pyrenebutanoic acid N-hydroxysuccinimide ester (PBASE)




Figure 5.1: Two conformations of PBASE molecule with geometry optimised via *ab initio* calculation (computed using Gaussian software [1]). The difference between computed Hartree-Fock energies is 1.0 kJ/mol, small enough that the existence of both conformations is physically possible.

1-Pyrenebutanoic acid N-hydroxysuccinimide ester (variously known commercially and in the literature as 1-Pyrenebutyric acid N-hydroxysuccinimide ester, PBASE, PBSE, PASE, Pyr-NHS, PyBASE, PANHS) is an aromatic molecule commonly used for tethering biomolecules to the carbon rings of graphene and carbon nanotubes. The use of this bifunctional molecule for noncovalent functionalisation of proteins onto a single-walled carbon nanotube was first reported in 2001 by Chen *et al.* [2]. Two methods for protein functionalisation were successfully used, with the only differences being the solvent used to dissolve the PBASE powder (DMF, methanol) and the final concentration of the resulting solutions (6 mM, 1 mM respectively). The lower concentration may have been used for PBASE in methanol as PBASE appears to dissolve poorly in methanol at

higher concentrations. Subsequent publications appear to have largely either chosen or adapted one of these two methods, as demonstrated by the frequency of the use of 6 mM PBASE in DMF and 1 mM PBASE in methanol in Table 5.1. Cella *et al.*, Campos *et al.*, Zheng *et al.* and Ohno *et al.* directly cite Chen *et al.* when discussing functionalisation with PBASE [3]–[6].

However, despite these various methodologies appearing to possess a common ancestor, there is a large degree of variation in

We purchased PBASE from two suppliers, Sigma-Aldrich and Setareh Biotech. Sigma recommends DMF and methanol as suitable solvents for dissolving PBASE alongside chloroform and DMSO. Setareh Biotech indicates methanol can be used for dissolving PBASE. The two suppliers have conflicting information for suitable storage of PBASE, with Sigma recommending room temperature storage while Setareh Biotech recommends storage of  $-5$  to  $-30^{\circ}\text{C}$  and protection from light and moisture.

### 5.1 Linker molecules

Table 5.1: Comparison of PBASE functionalisation processes used for immobilisation of proteins and aptamers onto liquid-gated CNTFET and graphene FET sensors

Solvent	Channel	Conc. (mM)	Incubation type	Time (hr)	Rinse steps	References
DMF	CNTs	5	Immersed	1	PBS	Maehashi <i>et al.</i> [7]
		6	Immersed	1	DMF, PBS	García-Aljaro <i>et al.</i> [8]
		6	Immersed	1	DMF	Chen <i>et al.</i> [2]
		6	Immersed	1	DMF	Cella <i>et al.</i> [3]
		6	Immersed	1	DMF	Das <i>et al.</i> [9]
	Graphene	-	-	2	DMF	Kwong Hong Tsang <i>et al.</i> [10]
		-	-	20	-	Wiedman <i>et al.</i> [11]
		0.2	Immersed	20	DMF, IPA, DI water	Gao <i>et al.</i> [12]
		1	100 $\mu$ L droplet	6	DMF, IPA, DI water	Nekrasov <i>et al.</i> [13]
		5	Immersed	1	DMF, DI water	Hwang <i>et al.</i> [14]
		6	6 $\mu$ L droplet	2	DMF, DI water	Nur Nasufiya <i>et al.</i> [15]
		10	10 $\mu$ L droplet	2	DMF, DI water	Campos <i>et al.</i> [4]
		10	Immersed	2	DMF, PBS	Kuscu <i>et al.</i> [16]
		10	Immersed	1	DMF	Xu <i>et al.</i> [17]
		10	Immersed	12	DMF, ethanol, DI water	Khan <i>et al.</i> [18]
2-Methoxyethanol	Graphene	1	Immersed	1	DI water	Ono <i>et al.</i> [19]
Methanol	CNTs	1	Immersed	1	Methanol, DI water	Zheng <i>et al.</i> [5]
		1	Immersed	2	Methanol	Kim <i>et al.</i> [20]
	Graphene	5	Immersed	2	-	Sethi <i>et al.</i> [21]
		5	Immersed	1	Methanol, PBS	Ohno <i>et al.</i> [6]
DMSO	CNTs	10	-	1	DI water	Lopez <i>et al.</i> [22]
		10	Immersed	1	PBS	Strack <i>et al.</i> [23]



## 6 Results

What I found out.

See for more detailed results



## 7 Results

What I found out.

See for more detailed results





## 8 Summary

In summary, this book has no content whatsoever.

[1] 2



## References



# Bibliography

- [1] J. A. M. J. Frisch and G. W. Trucks and H. B. Schlegel and G. E. Scuseria and M. A. Robb and J. R. Cheeseman and G. Scalmani and V. Barone and G. A. Petersson and H. Nakatsuji and X. Li and M. Caricato and A. V. Marenich and J. Bloino and B. G. Janesko and R. G. J. E. Peralta, F. Ogliaro, et al. *Gaussian~16 Revision C.01*. 2016.
- [2] R. J. Chen, Y. Zhang, D. Wang, et al. “Noncovalent sidewall functionalization of single-walled carbon nanotubes for protein immobilization [11]”. In: *Journal of the American Chemical Society* 123.16 (2001), pp. 3838–3839. ISSN: 00027863. DOI: 10.1021/ja010172b. URL: <http://pubs.acs.org>.
- [3] Lakshmi N. Cella, Pablo Sanchez, Wenwan Zhong, et al. “Nano aptasensor for Protective Antigen Toxin of Anthrax”. In: *Analytical Chemistry* 82.5 (Mar. 2010), pp. 2042–2047. ISSN: 00032700. DOI: 10.1021/ac902791q. URL: <https://pubs.acs.org/doi/full/10.1021/ac902791q>.
- [4] Rui Campos, Jérôme Borme, Joana Rafaela Guerreiro, et al. “Attomolar label-free detection of dna hybridization with electrolyte-gated graphene field-effect transistors”. In: *ACS Sensors* 4.2 (Feb. 2019), pp. 286–293. ISSN: 23793694. DOI: 10.1021/acssensors.8b00344. URL: <https://pubs.acs.org/doi/full/10.1021/acssensors.8b00344>.
- [5] Han Yue Zheng, Omar A. Alsager, Bicheng Zhu, et al. “Electrostatic gating in carbon nanotube aptasensors”. In: *Nanoscale* 8.28 (July 2016), pp. 13659–13668. ISSN: 20403372. DOI: 10.1039/c5nr08117c. URL: <https://pubs.rsc.org/en/content/articlehtml/2016/nr/c5nr08117c> <https://pubs.rsc.org/en/content/articlelanding/2016/nr/c5nr08117c>.
- [6] Yasuhide Ohno, Kenzo Maehashi, and Kazuhiko Matsumoto. “Label-free biosensors based on aptamer-modified graphene field-effect transistors”. In: *Journal of the American Chemical Society* 132.51 (Dec. 2010), pp. 18012–18013. ISSN: 00027863. DOI: 10.1021/ja108127r. URL: <https://pubs.acs.org/doi/full/10.1021/ja108127r>.
- [7] Kenzo Maehashi, Taiji Katsura, Kagan Kerman, et al. “Label-free protein biosensor based on aptamer-modified carbon nanotube field-effect transistors”. In: *Analytical Chemistry* 79.2 (Jan. 2007), pp. 782–787. ISSN: 00032700. DOI: 10.1021/ac060830g. URL: <https://pubs.acs.org/doi/full/10.1021/ac060830g>.

- [8] Cristina García-Aljaro, Lakshmi N. Cella, Dhamanand J. Shirale, et al. “Carbon nanotubes-based chemiresistive biosensors for detection of microorganisms”. In: *Biosensors and Bioelectronics* 26.4 (Dec. 2010), pp. 1437–1441. ISSN: 09565663. DOI: 10.1016/j.bios.2010.07.077.
- [9] Basanta K. Das, Chaker Tlili, Sushmee Badhulika, et al. “Single-walled carbon nanotubes chemiresistor aptasensors for small molecules: Picomolar level detection of adenosine triphosphate”. In: *Chemical Communications* 47.13 (Mar. 2011), pp. 3793–3795. ISSN: 1364548X. DOI: 10.1039/c0cc04733c. URL: <https://pubs.rsc.org/en/content/articlehtml/2011/cc/c0cc04733c> <https://pubs.rsc.org/en/content/articlelanding/2011/cc/c0cc04733c>.
- [10] Deana Kwong Hong Tsang, Tyler J. Lieberthal, Clare Watts, et al. “Chemically Functionalised Graphene FET Biosensor for the Label-free Sensing of Exosomes”. In: *Scientific Reports* 9.1 (Sept. 2019), pp. 1–10. ISSN: 20452322. DOI: 10.1038/s41598-019-50412-9. URL: <https://www.nature.com/articles/s41598-019-50412-9>.
- [11] Gregory R. Wiedman, Yanan Zhao, Arkady Mustaev, et al. “An Aptamer-Based Biosensor for the Azole Class of Antifungal Drugs”. In: *mSphere* 2.4 (Aug. 2017). ISSN: 23795042. DOI: 10.1128/msphere.00274-17. URL: [/pmc/articles/PMC5566834/](https://pmc/articles/PMC5566834/) <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5566834/?report=abstract> <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5566834/>.
- [12] Zhaoli Gao, Han Xia, Jonathan Zauberman, et al. “Detection of Sub-fM DNA with Target Recycling and Self-Assembly Amplification on Graphene Field-Effect Biosensors”. In: *Nano Letters* 18.6 (June 2018), pp. 3509–3515. ISSN: 15306992. DOI: 10.1021/acs.nanolett.8b00572. URL: <https://pubs.acs.org/doi/full/10.1021/acs.nanolett.8b00572>.
- [13] Nikita Nekrasov, Natalya Yakunina, Averyan V. Pushkarev, et al. “Spectral-phase interferometry detection of ochratoxin a via aptamer-functionalized graphene coated glass”. In: *Nanomaterials* 11.1 (Jan. 2021), pp. 1–10. ISSN: 20794991. DOI: 10.3390/nano11010226. URL: <https://www.mdpi.com/2079-4991/11/1/226/htm> <https://www.mdpi.com/2079-4991/11/1/226>.
- [14] Michael T. Hwang, B. Landon Preston, Lee Joon, et al. “Highly specific SNP detection using 2D graphene electronics and DNA strand displacement”. In: *Proceedings of the National Academy of Sciences of the United States of America* 113.26 (June 2016), pp. 7088–7093. ISSN: 10916490. DOI: 10.1073/pnas.1603753113. URL: <https://www.pnas.org/doi/abs/10.1073/pnas.1603753113>.
- [15] Mohd Maidin Nur Nasyifa, A. Rahim Ruslinda, Nur Hamidah Abdul Halim, et al. “Immuno-probed graphene nanoplatelets on electrolyte-gated field-effect transistor for stable cortisol quantification in serum”. In: *Journal of the Taiwan Institute of Chemical Engineers* 117 (Dec. 2020), pp. 10–18. ISSN: 18761070. DOI: 10.1016/j.jtice.2020.12.008.

- [16] Murat Kuscü, Hamideh Ramezani, Ergin Dinc, et al. “Graphene-based Nanoscale Molecular Communication Receiver: Fabrication and Microfluidic Analysis”. In: (June 2020). arXiv: 2006.15470. URL: <https://arxiv.org/abs/2006.15470v2>.
- [17] Shicai Xu, Jian Zhan, Baoyuan Man, et al. “Real-time reliable determination of binding kinetics of DNA hybridization using a multi-channel graphene biosensor”. In: *Nature Communications* 8.1 (Mar. 2017), pp. 1–10. ISSN: 20411723. DOI: 10.1038/ncomms14902. URL: <https://www.nature.com/articles/ncomms14902>.
- [18] Niazul I. Khan, Mohammad Mousazadehkasin, Sujoy Ghosh, et al. “An integrated microfluidic platform for selective and real-time detection of thrombin biomarkers using a graphene FET”. In: *Analyst* 145.13 (June 2020), pp. 4494–4503. ISSN: 13645528. DOI: 10.1039/d0an00251h. URL: <https://pubs.rsc.org/en/content/articlehtml/2020/an/d0an00251h> <https://pubs.rsc.org/en/content/articlelanding/2020/an/d0an00251h>.
- [19] T Ono, K Kamada, R Hayashi, et al. “Lab-on-a-graphene-FET detection of key molecular events underpinning influenza 2 virus infection and effect of antiviral drugs 3 Running title: Graphene-FET detects reactions in an influenza infection MAIN TEXT”. In: *bioRxiv* (Mar. 2020), p. 2020.03.18.996884. DOI: 10.1101/2020.03.18.996884. URL: <https://doi.org/10.1101/2020.03.18.996884>.
- [20] Jun Pyo Kim, Byung Yang Lee, Joohyung Lee, et al. “Enhancement of sensitivity and specificity by surface modification of carbon nanotubes in diagnosis of prostate cancer based on carbon nanotube field effect transistors”. In: *Biosensors and Bioelectronics* 24.11 (July 2009), pp. 3372–3378. ISSN: 09565663. DOI: 10.1016/j.bios.2009.04.048. URL: <https://pubmed.ncbi.nlm.nih.gov/19481922/>.
- [21] Jagriti Sethi, Michiel Van Bulck, Ahmed Suhail, et al. “A label-free biosensor based on graphene and reduced graphene oxide dual-layer for electrochemical determination of beta-amyloid biomarkers”. In: *Microchimica Acta* 187.5 (May 2020), pp. 1–10. ISSN: 14365073. DOI: 10.1007/s00604-020-04267-x. URL: <https://link.springer.com/article/10.1007/s00604-020-04267-x>.
- [22] Ryan J. Lopez, Sofia Babanova, Kateryna Artyushkova, et al. “Surface modifications for enhanced enzyme immobilization and improved electron transfer of PQQ-dependent glucose dehydrogenase anodes”. In: *Bioelectrochemistry* 105 (Oct. 2015), pp. 78–87. ISSN: 1878562X. DOI: 10.1016/j.bioelechem.2015.05.010. URL: <https://pubmed.ncbi.nlm.nih.gov/26011132/>.
- [23] Guinevere Strack, Robert Nichols, Plamen Atanassov, et al. “Modification of carbon nanotube electrodes with 1-pyrenebutanoic acid, succinimidyl ester for enhanced bioelectrocatalysis”. In: *Methods in Molecular Biology* 1051 (2013), pp. 217–228. ISSN: 10643745. DOI: 10.1007/978-1-62703-550-7\_14. URL: <https://pubmed.ncbi.nlm.nih.gov/23934807/>.