

Weird stuff from not so weird stuff*

NEEV SHAH ¹

¹*Steward Observatory, The University of Arizona*

ABSTRACT

This example manuscript is intended to serve as a tutorial and template for authors to use when writing their own AAS Journal articles. The manuscript includes a history of AAS_TE_X and documents the new features in the previous versions as well as the new features in version 7. This manuscript includes many figure and table examples to illustrate these new features. Information on features not explicitly mentioned in the article can be viewed in the manuscript comments or more extensive online documentation. Authors are welcome to replace the text, tables, figures, and bibliography with their own and submit the resulting manuscript to the AAS Journals peer review system. The first lesson in the tutorial is to remind authors that the AAS Journals, the Astrophysical Journal (ApJ), the Astrophysical Journal Letters (ApJL), the Astronomical Journal (AJ), and the Planetary Science Journal (PSJ) all have a 250 word limit for the abstract. The limit is 150 for RNAAS manuscripts. If you exceed this length the Editorial office will ask you to shorten it. This abstract has 189 words.

Keywords: Galaxies (573) — Cosmology (343) — High Energy astrophysics (739) — Interstellar medium (847) — Stellar astronomy (1583) — Solar physics (1476)

1. INTRODUCTION

1. Background

2. Introduce XRBs, runaways, HMXBs - 4U1700-37 - well studied, many observational constraints - not well known if NS or not

3. Introduce GWs - GW190814 - unknown origin, conventionally difficult to form

4. Lower mass gap - GW190814, new NSBH GW, some EM observations in the mass gap as well. Unknown if real or not

5. Connection between XRBs and GWs - XRBs perhaps intermediate path.

6. Leverage XRB constraints to chart past history, connect it to GW190814 at low Z

7. What exactly are we trying to show? Forming asymmetric mass ratio systems starting from symmetric, followed by conservative mass transfer. Kick can lead to variety of outcomes, including XRBs, TZOs, GWs. GW190814 not a weirdo to be ignored

8. Why is it important

9. Describe paper outline

X-ray binaries (XRBs) and compact object mergers are some of the most exotic high energy phenomena in the universe. The presence of a compact object (or even two) in such systems implies that their progenitors must have consisted of at least one massive star that has ended its life in a spectacular explosion or implosion, leaving behind a neutron star or a black hole remnant. The formation of X-ray binaries and gravitational wave sources is a complex and poorly constrained process, arising from our uncertainties in the evolution of massive stars and their winds, coupled with the uncertainties in binary evolution such as mass transfer, common envelope evolution, and the effects of tides. The observations of XRBs in the galaxy and the local universe and the detection of GW events have provided us with a wealth of information that can be used to constrain their formation pathways, and thereby better understand the evolution of massive stars in binaries. The pathway of forming an XRB or compact binary starting from a pair of massive stars is complex, with numerous uncertainties at each step. To add to that, such bound systems are rare outcomes of binary evolution, with most binaries getting disrupted at the first supernova, or a premature merger before the formation of a compact object.

The usual pathway to forming an XRB involves a phase of mass transfer before the end of the life of the donor star. This flips the mass ratio, and if the system remains bound after the first supernova (depending on the natal kick and the mass lost in the explosion), we are left with a compact object orbiting a star. If the system is tight enough, the companion star will eventually grow large enough such that mass is transferred to the compact object, either through stellar winds or RLOF. This mass accretion on the compact object generates X-rays, and such a system is referred to as an X-ray binary. Eventually, another phase of mass transfer may occur, now from the companion star (originally the less massive star) to the compact object. This phase can either be stable or become unstable, leading to the formation of a common envelope. If the binary survives this phase of mass transfer and the companion star is massive enough to leave behind a compact object remnant, we may be left behind with a tight binary consisting of two compact objects which may eventually merge due to gravitational wave emission. Thus, high mass XRBs (HMXBs) are a key link in the evolutionary pathway of forming gravitational wave sources from massive binaries. HMXBs can be accurately studied in the local universe, with their orbital properties constrained via various observational probes.

One of the first X-ray sources to be discovered in the 1970s is the HMXB 4U1700-37. It also consists of potentially the most massive companion star, HD 153919, an O6.5 Iaf+ star with an estimated mass surpassing $40M_{\odot}$. The system is in a rather short orbit of just 3.41 days, and is also a runaway HMXB with a large spatial velocity. With the help of Gaia kinematics, this system can be traced back to have originated from the young open cluster NGC 6231. It is likely that around 2Myr ago, the originally more massive star in the binary exploded in a supernova which kicked it out of the cluster, while keeping the system bound so that it is visible as a HMXB today. Given the young age of the system at the time of explosion, the progenitor star that formed the compact object must have been quite massive. However, the compact object is estimated to have a mass of just $2.5M_{\odot}$, which is straddles the supposed mass gap between the heaviest neutron stars and the lightest black holes. This makes the past evolutionary history of the binary particularly interesting, and its well characterized orbital properties and kinematic speed can be used in constraining the progenitor properties of the binary.

Meanwhile, on 14th August 2019, the LIGO and Virgo detectors detected a gravitational wave event, GW190814, which consisted of a merger between a $23M_{\odot}$ black hole and a $2.5M_{\odot}$ compact object. The

nature of the less massive compact object is uncertain. The mass ratio of the two objects is particularly interesting, as it is the most asymmetric mass ratio observed in a compact binary merger. The formation of such an asymmetric system has been difficult to explain through standard formation channels. Given the similarity in masses between the HMXB and GW190814, (van der meijs+21) proposed that they may share a common formation pathway. In this work, we explore this connection in further detail with the help of detailed binary evolution model computed with MESA. We also utilize monte carlo simulations of the effects of supernova explosions in binaries, which helps us constrain the pre-supernova properties of the HMXB. This in turn informs our MESA binary models, which we evolve to the point of the first core-collapse in the system (defined as till carbon depletion). We then go on to evolve a similar progenitor binary at a lower metallicity to explore the scenarios in which we can form a GW190814-like system.

In Section (Methods), we describe our setup for computing the evolution of a binary system with MESA. We also describe our monte carlo method, wherein we use the observational constraints of a post-supernova binary to infer its properties pre-supernova. In Section (XRB), we demonstrate our monte carlo method by applying it to the HMXB, and constrain its pre-supernova properties. Motivated by those results, we use MESA to construct an evolutionary pathway for this system starting from a massive binary. We end the section with a discussion on the future evolution of the XRB, and its connection to TZO's. In Section (GW190814), we construct similar MESA models at lower metallicity to explore the formation of a GW190814-like system. We discuss the conditions in which such a system can form, and its similarities and differences with the HMXB described in the previous section. We also make toy estimates for the rates of such events, and what makes them rare. In Section (Conclusion), we summarize our results and discuss the implications of our work.

2. METHODS

2.1. MESA

We simulate the evolution of massive binaries using MESA (version xxx). Our inlists and choice of input parameters are public (zenodo/github) and we describe some of the important ones below.

1. MESA (star) - basic stellar physics assumptions - MLT, overshooting, other mixing, rotation, winds, supereddington treatment
2. MESA (binary) - Two stars evolved self consistently, along with orbit equations. Treatment of

mass transfer (Kolb & Ritter), AM transport, loss from system, tides, rotationally enhanced winds. Rotation only included in accretor, turned off in donor - justify why?

3. Monte Carlo - Outcome of supernova in a binary
 - Assumptions: Pre-SN circular orbits, instantaneous mass loss, Blaauw kick, Natal kick. Mass loss and kick vector gives post-SN orbital parameters, systemic velocity and eccentricity.

We use the Ledoux criterion to model convection, and assume a mixing length parameter of 2.0. We include semiconvection ($\alpha_{sc} = 1$), and thermohaline mixing ($\alpha_{th} = 1$). To account for convective boundary overshooting above the core, we use the exponential model as implemented in MESA, and we do not account for convective boundary overshooting for off-center convective layers. To aid in regions at the eddington limit where convection is also inefficient, we utilize the local implicit enhancement of the convective flux in superadiabatic regions introduced in MESA 15140 (`use superad reduction = .true.`).

For modeling rotation, we use the "shellular" approximation where the angular velocities ω are fixed on isobaric surfaces (which assumes strong anisotropic horizontal turbulence). We also assume rigid rotation at the beginning of our runs. We include several rotational mixing processes, and also enhanced wind mass loss due to rotation to keep the star to rotate subcritically. We turn off rotation in the donor to avoid numerical issues, and because the spin of the compact object is largely unconstrained for our XRB. Meanwhile, the companion is a visible star with a $v \sin(i)$ measurement, and hence we do not ignore rotation in the accretor star. We do not believe that the rotation of the donor star has a significant effect in the interpretation of our results.

We include stellar winds by combining several different mass loss prescriptions. For stars with a surface effective temperature greater than 11000K, we utilize the Bjorklund prescription, while for cooler stars (lesser than 10000K), we use the Decin winds, interpolating in temperature ranges in between. For the hot stars with a surface hydrogen mass fraction less than 0.4, we switch to the optically thick WR winds from Nugis and Lamers.

We treat mass transfer in a binary using an implicit scheme as described in Kolb and Ritter. We also include the effects of tides, which can have an important effect on the evolution of the binary given the short periods that we work with.

2.2. Monte Carlo

Here, we describe our method of inferring the pre-supernova parameters of a binary given its observational constraints post explosion. We follow (el badry+24, brandt,pods+95) in our setup.

To start with, we assume that the binary has a circular orbit before explosion. This is justified for our scenario, because we model short period binaries that undergo mass transfer, and hence tides are expected to have circularized the orbit. Once a star dies in a binary, its orbit is affected due to the following

- instantaneous mass loss in a supernova explosion (Δm) which we assume is lost from the binary.
- the speed and direction of a potential natal kick to the supernova remnant ($v_{kick}, \{\theta, \phi\}$)

A combination of these affects can lead to diverse outcomes post explosion. A binary could get disrupted if sufficient mass is lost, or if the natal kick is strong enough. If a binary survives, it may have an induced eccentricity, and the center of mass may also gain a systemic velocity. These affect the size and orbital period of the binary as well.




Figure 1. This is a caption for your figure.

3. RESULTS

3.1. Monte Carlo Results

1. Describe what we are doing in this section - Infer pre-SN parameters of XRB from observational constraints
2. Define relevant variables, initial population, assumptions (circular orbits) outcomes
3. Emphasize that population is not observationally motivated, so probabilities are not real. Meant to explore parameter space
4. Simulate outcomes of SN kicks - put observational constraints (define them)
5. Show corner plot - explain what it implies - Pre-SN period and mass constraints
6. Explain lower left corner gap in Pre-SN period - emphasize why kick is likely (based on MT period widening, Renzo+19), and kick needed to shrink
7. Explain black shaded area - unlikely to fit without RLOF, theoretically not expected due to orbit widening, wind mass loss over kinematic age gives minimum period 2d. In black is shown distribution for post-SN period ≥ 2 d

In this section, we describe the outcomes of our monte carlo simulations that simulate the effects of a supernova in a binary. Note that these are different from population synthesis simulations, as the initial binary population that we use is not observationally motivated, but is meant to explore the parameter space of interest relevant to the properties of the XRB.

We start with generating 10^7 binaries, with the following properties

1. The mass of the companion star is uniformly sampled between $40 - 60M_{\odot}$. This roughly spans the measured mass of the visible star today, and we do not expect it to change significantly during the supernova and until today. In our simulations, we assume this mass to be fixed before and after the SN.
2. For the progenitor of the $2.5M_{\odot}$ compact object, we sample the mass lost in the supernova explosion (Δm) uniformly between $0 - 20M_{\odot}$. We fix the post-SN mass of the compact object to be $2.5M_{\odot}$. The mass of the pre-SN star is then given by $2.5M_{\odot} + \Delta m$.
3. We sample the pre-SN period of the binary uniformly between $0 - 30$ days.

4. We sample the kick distribution from a maxwellian distribution with a scale of 265km/s and isotropic in direction.
5. We assume the pre-SN orbits to be circular.

Now that we have defined our pre-SN population, we simulate the outcomes of a SN explosion on the properties of the binary. Depending on the amount of mass lost in the explosion and the strength and direction of the natal kick to the compact object, the orbit of the binary can change or it may even become unbound. It may also induce some eccentricity in the orbit, and give the center of mass of the binary a systemic velocity. By sampling different combinations of mass loss and kick vectors, we can infer which binaries get unbound, which ones are bound, and what are the new orbits of the bound systems. We then compare them to the observational constraints that we have for the XRB, and select for systems that satisfy those conditions.

The observational constraints that we impose are as follows

1. The orbital period post-SN is between $2 - 3.41$ days. We place a lower limit of 2 days as orbital periods smaller than this imply that the current visible star would have filled its roche lobe post-SN which does not seem plausible given its current evolutionary state. As the current orbital period is measured to be around 3.41 days, we allow for post-SN periods smaller than this because wind mass loss post-SN could have widened the system to its current orbital period as well.
2. The systemic velocity inherited during the SN is $50-80\text{km/s}$ (the measured systemic velocity with respect to the open cluster is 65km/s). Here, we are making the assumption that most of the systemic velocity of the XRB with respect to its birth cluster today can be attributed to its post-SN systemic velocity. We are also implicitly assuming that this velocity has not significantly changed during its runaway phase for the past couple of Myr. However, the details of this would depend on accurately modeling the effects of the cluster potential, which we do not model in this work.
3. We assume that the induced eccentricity in the binary is less than one, meaning that we only select for systems that remain bound post-SN (there are disagreements in literature on whether the binary is close to circular or has a modest eccentricity of 0.2). Since the binary could have had a higher eccentricity post-SN (but less than one) and has undergone tidal circularization since then during the

runaway phase, we do not impose any constraints based on its measured eccentricity (controversial) today.

Fig. 1 shows the distribution of binary parameters that satisfy the observational constraints of the XRB. We emphasize that the probability distributions here are not to be taken literally, as the probability distribution of the pre-SN population was not physically motivated. We find that the pre-SN mass of the progenitor of the compact object must have been between $4 - 12M_{\odot}$. We also find that the pre-SN orbital period was likely smaller than 9 days. Although we do not put any constraints on the pre-SN orbital period, it is likely that it could not have been very small ($< 1 - 2$ days) as that would imply that the companion star would fill its roche lobe before the SN. We now point out some interesting regions in the corner plot that provide clues on the XRB's past. We find that large pre-SN periods (> 4 days) and small or no kicks are incompatible, as it would not be possible to shrink the binary orbit to its post-SN period in such cases. This is important because mass transfer in binaries tends to widen the orbital period in most scenarios. Given that this XRB likely went through a phase of mass transfer in the past (from the progenitor of the CO onto the current visible star), its post-SN period was likely large enough that a natal kick is necessarily needed to shrink the orbit post-SN to its current state.

Another excluded region in this plot is the lower left corner in the pre-SN period vs kick panel. This shows that small enough pre-SN periods and low kicks are incompatible with the imposed observational constraints. The reasons for this are a bit subtle, and we explain it for the case of no kicks, which can then be easily generalized to the presence of kicks. We have imposed a constraint that the post-SN period is larger than 2 days, and the systemic velocity is smaller than 80km/s. For a small enough pre-SN period, it so happens that to widen the orbit to satisfy the imposed constraints, the systemic velocity required exceeds the observational constraints. However, kicks allow some room for scatter in such situations, and thus the minimum pre-SN period decreases as we increase the strength of the kicks, which creates the diagonal boundary in the lower left corner in the pre-SN period vs kick panel.

We thus find that there is a very narrow range of pre-SN periods ($1.5 - 2.5$ days), where the observational constraints can be satisfied without a kick (quantified as kicks lesser than 20km/s). Such small pre-SN periods after an initial phase of stable MT are already quite unlikely given the theoretical expectations of widening of the orbit during MT. This strengthens the case for the requirement for a natal kick at the birth of the $2.5M_{\odot}$

compact object, irrespective of whether it is a NS or a low mass BH.

3.2. MESA-Model-XRB

1. Monte carlo constraints give us insight into past - conservative mass transfer (pre-SN mass), short period (pre-SN period)
2. Progenitor system - 40+30, 3d. Case A MT
3. Describe evolution, and also HR diagram. Focus on key phases - Fast+Slow Case A, Case AB, WR winds during helium burning, explosion, Accretor growth till TAMS
4. Compare with observed properties of the star
5. what do we get wrong (describe here or later in discussion)
6. Future of XRB - Binding energy curves, alpha common envelope, likely failure, TZO/transient
7. Perhaps discuss here i) why do we need a natal kick, why not a common envelope, why not Case B (or place this somewhere else if more appropriate)

3.3. Why do we need a natal kick?

Compact object that are formed in a successful supernova explosion may receive a natal kick at birth due to asymmetries in the explosion or in neutrino emission. Observationally, the natal kicks received by neutron stars is well described by a maxwellian distribution with a mean of 265km/s. However, for black holes, the story is a bit more complicated with evidence pointing towards weak or no kicks in some cases or strong kicks in others. For our X-ray binary of interest, we know that this is a runaway system from Gaia kinematics. The runaway nature does not necessarily imply that the compact object received a natal kick, as some mass lost in a spherically symmetric can also impart a systemic velocity to the binary and also induce eccentricity in the system. This is referred to as the "Blauuw kick" and the new binary orbit (if the system did not get disrupted) always has a larger orbital period (and therefore the orbital separation) because of the mass lost in the explosion. However, a decrease in the orbital period may be possible given a natal kick in a favourable direction. Given that the XRB currently is in a very tight orbit of just 3.41 days, a Blauuw kick implies that the pre-supernova orbit must have been even tighter. This may be inconsistent with most scenarios of mass transfer through RLOF, which in general leads to a widening of the orbit. Therefore, we assume that the pre-supernova

orbital period was larger than the current one, and that the shrinking of the orbit is due to a natal kick received by the compact object. Not just that, for the orbit to shrink, we find that the natal kick must be directed away from the orbital velocity of the star, as seen in (insert fig)

4. WHY NOT A COMMON ENVELOPE?

Given the short period of the XRB, it may seem appealing to invoke a common envelope formation scenario for this system. In this section, we discuss why it is difficult to form this X-ray binary through a common envelope instead of stable RLOF. To start with, the visible star in the system (and the originally less massive one) has a high mass of 40-60 M_{sun} . Since this star likely did not gain any mass during a supposed common envelope phase, the original donor that formed the compact object must have been more massive than 40-60 M_{sun} . In a common envelope phase, the donor would lose its envelope, and leave behind a core that would be far too massive to successfully explode and form a 2.5 M_{sun} compact object (however see Burrows et al etc papers). To add to that, our monte carlo experiments imply that the pre-SN donor mass is less than 15 M_{sun} , which is far too small of a core for a star that is more massive than 40-60 M_{sun} . Also, common envelope phases usually are thought to occur for asymmetric binaries, and this may imply that the donor should have been much more massive than the current visible star. All of these constraints may imply that forming this X-ray binary through a common envelope phase is unlikely, and that the likely scenario may involve stable mass transfer through RLOF.

5. WHY NOT CASE B MASS TRANSFER?

Previous studies of this system have suggested formation pathways wherein mass transfer occurs after the donor star has left the main sequence (Case B). However, this is unlikely because of the short orbital period of the binary today. Renzo 2019 (insert citation) showed that for Case B mass transfer, the orbital period of the binary widens significantly in most cases, except for scenarios where mass transfer is non conservative ($\beta < 0.6$) and there are extreme losses in angular momentum ($\gamma_{\text{RLOF}} \geq 2$). To add to that, as Case B occurs after the donor has left the MS, the initial orbital period of the binary must have been > 10 days, and this would have most likely increased post-MT. As we find in our monte carlo results, even with a favourable kick in the right direction, it is likely extremely rare to get the period post-SN to less than 4 days starting from a > 10 day period pre-SN.

6. XRB

HD153919 is a well studied runaway high mass X-ray binary. It consists of a bright O-type star with a compact companion that straddles the mass gap between the heaviest neutron stars and lightest black holes at 2.5 M_{sun} . The extreme mass ratio (20 to 1), short orbital period and the runaway nature of the binary have been particularly challenging to simultaneously explain through standard formation channels. Using Hipparcos, and more recently with Gaia DR3, the binary can be confidently traced back to have originated from the young open cluster NGC6231. By measuring the systemic velocity with respect to the cluster, we know that the binary must have been kicked out 2.2Myr ago. We also know that the cluster is very young with quite a few massive stars still present, with age estimates ranging from 4-10Myr. It is likely that a supernova explosion in the binary which formed the 2.5 M_{sun} compact object gave the system a systemic velocity that ejected it out of the cluster. Given the young age of the cluster at the time of explosion, the progenitor star that exploded must have originally been quite massive as well. Since this XRB is well studied, we have strong constraints on the binary's orbit, the individual masses and on the systemic velocity which presumably is dominated by the kick it received due to the supernova explosion. Given these constraints, we perform monte-carlo simulations to estimate what range of pre-supernova orbital configurations can explain the current observational constraints of the binary. We model the supernova explosion as an instantaneous mass loss event, and together with conservation of momentum, we can recompute the induced eccentricity in the binary. An eccentricity greater than one implies that the explosion disrupts the binary, while for eccentricities less than one we can recompute the new orbital period and separation, along with the systemic velocity induced in the binary due to the explosion. We also account for the effect of natal kicks that the compact object can receive during a supernova and its effect on the fate of the binary. To summarise, the observational constraints that we impose are

6.1. Results – XRB

In this section, we describe the evolution of our supposed progenitor system of the HMXB 4U-1700-27. Motivated by the results of our monte carlo simulations, we start with a binary of initial masses 40 M_{\odot} and 33.5 M_{\odot} in a tight orbit with a period of just 3.5d. As a result, the donor overflows its roche lobe just after 3.48Myr, during the Main Sequence (MS) itself. This leads to a phase of mass transfer from the donor to the accretor, commonly referred to as Case A. This phase actu-

ally consists of an initial thermal timescale mass transfer (0.07Myr), known as fast Case A, where the donor loses a majority of the mass lost during RLOF. As the donor regains thermal equilibrium, it slowly grows in size on a nuclear timescale, which leads to a steady period of mass transfer referred to as slow Case A. This phase ends around 4.9Myr, roughly when the donor has finished its supply of hydrogen in its core, and it starts contracting on a thermal timescale, ending the first phase of RLOF. At this stage, the donor has lost around $15M_{\odot}$, with $11M_{\odot}$ lost during the thermal timescale fast Case A itself. It now has a mass of $24.7M_{\odot}$, and having lost much of its envelope, the surface now exposes the products of CNO-burning. It has a helium rich ($Y = 0.59$) and hot surface ($T_{\text{eff}} \approx 37000K$), and a thin envelope of $\approx 6.8M_{\odot}$. Note that the mass of the helium core at this point is almost $18M_{\odot}$, which is potentially far too large to lead to a successful explosion. On the other hand, the accretor (still on the MS) has grown from its initial $33.5M_{\odot}$ to $\approx 45.5M_{\odot}$. Due to the increase in mass, it is now overluminous with a $\log_{10}(L/L_{\odot}) \approx 5.7$ and a surface temperature of $37000K$.

Case AB

As the donor star still has a substantial amount of hydrogen left in its envelope, it ignites hydrogen burning in a shell, which drives a rapid increase in its radius on a thermal timescale. As the binary only expanded from an initial period of $3.5d$ to $4.6d$, the donor star once again overflows its roche lobe during the hydrogen shell burning phase. As this occurs after the initial Case A mass transfer, this period is referred to as Case AB mass transfer (with the B referencing mass transfer occurring after the MS but before helium depletion). This also occurs on a thermal timescale, where the donor loses an additional $4M_{\odot}$, bringing its mass down to $19.4M_{\odot}$. Due to rotational spin down by tides, the accretor is able to accrete most of this mass, and it grows to almost $49.2M_{\odot}$. This phase ends just after 10000yr (when it happens is messy). Since the donor is now even more stripped of hydrogen (surface $X \approx 0.2$), it starts contracting and has a higher surface temperature. It soon ignites helium in its center, and initiates the core helium burning phase, which lasts for $\approx 0.44\text{Myr}$. During this phase, it has a hot helium rich surface, and would resemble a Wolf-Rayet (WR) star with a strong optically thick wind. These mass loss rates are of the order of $10^{-5} - 10^{-6}M_{\odot}/\text{yr}$, and the donor star loses an additional $11M_{\odot}$ during its WR phase until helium depletion. At this point we terminate our MESA models, and the originally more massive star now has a mass of $13.3M_{\odot}$, with a CO core mass of $11.3M_{\odot}$ (central C/O ≈ 0.37). There is no hydrogen left on the sur-

face which is now mostly consisting of helium, carbon and some oxygen. Meanwhile, the originally less massive star (and the current visible/donor in the XRB) now has a mass of $48.2M_{\odot}$. The orbital period of the binary has now increased to $8.27d$, which aligns with the expectations from our monte carlo simulations.

Our monte-carlo simulations imply that the progenitor star of the compact object must have been 7-14Msun prior to explosion. This is a strong indicator for past binary interaction, as the progenitor must have been much more massive early on given the young age of the cluster (atleast $\gtrsim 30\text{Msun}$). We also know that the companion is quite massive today, and this could be due to accretion of mass from the donor. These constraints hint towards an episode of stable and conservative mass transfer in the past, and our goal is to chart an evolutionary pathway from a pair of progenitor stars to the current observed X-ray binary. Motivated by the constraints from our monte-carlo sims, we model the evolution of a binary consisting of 2 massive star of 40 and 30Msun in a short orbit of 4 days. At such short periods, tides efficiently circularize and synchronize the orbit. Due to the small separation between the stars, the more massive star overflows its roche lobe during its Main Sequence itself, starting a phase of mass transfer that is commonly referred to as Case A mass transfer.




Figure 2. This is a caption for your figure.

6.2. What do we get wrong?

Although our MESA models and monte carlo simulations provide an evolutionary pathway for the XRB, there are a few limitations that we have not addressed yet (TBD). While the accretor model reasonably matches the properties of the visible star HD153919, such as its mass, surface temperature and luminosity, we underestimate its remaining MS lifetime. The XRB has a kinematic age of 2.2Myr, with the visible star still on the MS, while our accretor model only has 0.7Myr left on the MS after the death of the donor star, and only 0.5Myr to reach the current radius of HD153919. There could be a couple of potential explanations for addressing this discrepancy.

1. The measured systemic velocity is lower than what it was in the past. This could happen if the XRB slows down to its current systemic velocity as it escapes the cluster potential. A higher systemic velocity in the past would imply a smaller kinematic age.
2. A larger initial donor mass (currently assumed to be $40M_{\odot}$) would imply a shorter lifetime for the progenitor of the compact object, thereby extending time spent in the post-RLOF phase for our accretor model. This is complicated by the fact that a larger donor mass would also imply a slightly larger pre-SN mass that may be harder to explode and form a $2.5M_{\odot}$ compact object.
3. A smaller initial accretor mass (currently assumed to be $33.5M_{\odot}$) would imply a smaller mass for the star post-RLOF, thereby extending its MS lifetime. This could be in tension with the measured luminosity of the star, as we find that (initially) lower mass accretors are less luminous at present day than what is inferred from observations.
4. The effects of rejuvenation in the accretor are larger than currently modeled in our binaries. Rejuvenation increases the core mass, and the extra supply of fuel can extend the MS lifetime. Thus, more rejuvenation could help address the discrepancy between the MS lifetimes in our models and the XRB's kinematic age.

It could also be that the discrepancy could be addressed by a combination of the above factors, or something else entirely. Previous studies have also seen a 2x discrepancy in the ages as measured from single star evolutionary grids and kinematic ages.

Secondly, our donor models at carbon depletion have a total mass of $13.3M_{\odot}$ and a Carbon-Oxygen core of

$11.3M_{\odot}$. Population synthesis models would predict such stars to implode and form direct collapse black holes, and not successfully explode to leave behind a $2.5M_{\odot}$ compact object. However, the explodability of massive stars is still an active area of research, with 3D simulations pointing towards a more nuanced picture than simple prescriptions just based on the core mass.

7. GW190814

GW190814 was an exceptional event that consisted of a merger between a 23Msun black hole and 2.5Msun compact object. The GW data alone was not enough to characterize the black hole or neutron star nature of the less massive compact object, and the formation of such a system of such an asymmetric mass ratio has been puzzling for standard formation channels.

7.1. MESA model - GW190814

1. Based on "success" of XRB model, propose similar solution for GW190814
2. What will be different here? - Why is XRB a TZO - kick strength and direction
3. Why switch to lower masses - weaker winds
4. No observational constraints other than masses - more freedom
5. Describe fiducial GW model at Z/10 - key phases, HR diagram, final masses, donor explosion.
6. Describe possible futures
7. Evolve accretor as single star - next interaction depends on post-SN orbit. At smaller radii / lack of convective envelope / before RSG phase, binding energy high, merger likely. At larger radii, core envelope distinction, lower binding energy, CE ejection likely.
8. End state - large BH (implosion) + NS/low mass BH (after successful explosion + strong kick) - asymmetric mass ratio GW event

A key difference for stars at lower metallicity is that the wind mass loss becomes significantly weaker. From our previous MESA models at solar metallicity, we saw that the mass loss during the "WR phase" (core-helium burning) is extremely important in getting the donor star's mass down to what we expect from our monte carlo simulations. Utilizing the same stellar masses and computing their binary evolution at lower metallicity would leave us with pre-SN donor stars with sufficiently

high masses that would likely fail to explode. This motivates us to lower the initial masses of our stars while trying to model the progenitor of a GW190814-like system at lower metallicity. For our fiducial GW190814-like model, we start with a binary of $30M_{\odot} + 25M_{\odot}$ in a 3 day orbit. We show the evolution of this binary on the HR diagram in Fig. 3. Qualitatively, it follows a similar pathway as our fiducial XRB model. As stars at lower metallicity are more compact, mass transfer begins slightly later in the evolution. After 5.1Myr, the donor star overflows its roche lobe during the MS, initiating Case A mass transfer. This phase lasts around 1Myr, after which the donor has reached TAMS and starts contracting, detaching itself from its RL. At this stage, the donor star has a mass of $19M_{\odot}$, while the accretor has grown to about $36M_{\odot}$. Due to the short orbital period, tidal effects keep the accretor below critical rotation, and mass transfer remains close to fully conservative. The donor soon re-expands once it starts hydrogen shell burning, and starts a second phase of Case AB mass transfer, similar to our fiducial XRB model. At the end of this phase, we are left with a binary of $16.5M_{\odot} + 38.3M_{\odot}$ in a tight 3.85d orbit. Around the same time, the donor contracts and ignites helium burning in its core, and the star has a mass of $14.4M_{\odot}$ at helium depletion, with a carbon-oxygen core mass of $12.3M_{\odot}$, and a C/O ratio of 0.36, which is roughly similar to what we got for our fiducial XRB. On the other hand, the accretor, which is now the more massive star in this system has a mass of $38.5M_{\odot}$. Assuming that the donor star successfully explodes and leaves behind a NS or a low mass BH similar to $2.5M_{\odot}$, and the strength and direction of the natal kick, we can have a variety of outcomes. Large enough kicks can potentially disrupt the binary itself, while low kicks would just slightly widen the orbit. The now more massive star will continue growing as it burns hydrogen in its core, and begin a phase of reverse MT if it overflows its RL. The outcome of a phase of reverse RLOF onto a compact object sensitively depends on the evolutionary state of the star at that point. Due to the highly asymmetric mass ratio, MT is likely going to be unstable and a CE phase is likely. If the star interacts while it is still on or just past its MS, its envelope binding energies are likely going to be sufficiently high that it would meet the same fate as our fiducial XRB and form a transient/TZO. However, if the natal kick to the first born CO is large enough but not leading to disruption, the binary can be pushed into a wide orbit where the reverse MT starts much later in the evolution of the star. This is crucial for the successful ejection of a CE, as donor stars with large and loosely bound convective envelopes are easier to eject.

The star will likely develop a loosely bound convective envelope once it has expanded sufficiently and become cool enough to be a RSG. Interaction during the RSG phase significantly increases the possibility of a successful CE ejection, and we would be left behind with a tight binary, consisting of a helium core of about $20M_{\odot}$ and a $2.5M_{\odot}$ CO. Such a massive helium core is expected to implode and form a $20M_{\odot}$ BH without a natal kick. We are now left with a DCO system consisting of a $20M_{\odot}$ BH and a $2.5M_{\odot}$ CO, which would then merge after a GW driven inspiral. Such a system would appear to be similar in compact object masses to GW190814.




Figure 3. This is a caption for your figure.

8. FUTURE EVOLUTION

Now that we have been able to chart a reasonable past evolutionary history of the XRB using our MESA binary models, we evolve the system further to see what the future of this system could look like? The current mass of the compact object is $2.5M_{\odot}$ while the companion is $40\text{--}60M_{\odot}$ that is yet to fill its roche lobe. As the star evolves further, it will grow in size and soon overflow its roche lobe. However, given the highly asymmetric nature of the system, the subsequent mass transfer is likely to soon become unstable and the compact object will plunge into the companion star's envelope, starting a phase of common envelope evolution. This




Figure 4. This is a caption for your figure.




Figure 5. This is a caption for your figure.

we can compute the binding energy of the envelope as well. This requires us to assume a particular definition of core, and everything above it would be considered the envelope. If a common envelope ejection is successful, what remains of the donor star is its core, and we assume that it may have some sort of response that can expand its size by up to twice that of what the core size originally was at the star of the common envelope phase. If we assume a perfect use of orbital energy to unbind the envelope, i.e $\alpha = 1$, we can compute what would be the final separation between the compact object and the remnant core. If the remnant core fits inside the roche lobe of its new orbit, it implies that common envelope ejection was successful and we are left with a tight but detached binary. However, if the remnant core is too large for its final orbit, it means that there was a merger between the compact object and remnant core, and common envelope ejection was not successful.

In Fig. 4, we show the envelope binding energy of our fiducial accretor model at its current radius of about $25R_{\odot}$. We define it as

$$BE(m) = \int_m^M dm' \left(-\frac{Gm'}{r(m')} + \alpha_{th} u(m') \right) \quad (1)$$

where α_{th} is the parameter controlling the fraction of internal energy. Since there is no one definition for what is considered as the core, and what is the stellar envelope (especially for a MS star), we calculate the binding energy as a function of core mass. The binding energy at mass coordinate m is the energy required to lift up and eject all the stellar material above that mass coordinate. For a MS star, we consider two physically motivated definitions for the location of the core. One is the mass of the convective core, and the other is mass of the convective core *and* the overlying overshooting region. These two limits span the shaded area in Fig. 4. The red dashed and blue solid lines denote the binding energy curves with and without the contribution of the internal energy. Addition of internal energy provides an extra energy source which makes it easier to unbind the envelope (lower binding energy). We find that for our stellar model at its current evolutionary state, the binding energies lie between $[2.5 - 5.7] \times 10^{50}$ erg, depending on the assumed core mass definition and whether we include the internal energy or not.

In an ensuing phase of MT, if it turns unstable, the compact object plunges into the stellar envelope and the system enters a common envelope phase. The success (successful envelope ejection leaving behind a tight binary) or failure (failed ejection and premature merger) is a complex hydrodynamical process that can only be approximated in 1D stellar models. This is often done

phase is highly complex and multidimensional with various physical processes occurring at different time and length scales. Importantly, the success or failure of the common envelope phase is an active area of research with no concrete predictions. However, various techniques have been introduced to simulate (or skip over) this phase, enabling predictions to be made albeit at the cost of accuracy. One of the most commonly used approaches is the energy balance formalism or $\alpha\lambda$ mechanism. This parameterized method quantifies what fraction (α) of the orbital energy is required to unbind the donor star's envelope, which is parametrized by λ . To simulate the outcome of our XRB, we evolve it till it overflows its roche lobe and the mass transfer becomes unstable (quantified by $\dot{M} > 0.1 M_{\odot}/\text{yr}$). Using the orbital separation, we know how much orbital energy is available, and using the interior structure of the star,

with the $\alpha\lambda$ prescription, where λ quantifies the envelope binding energy, and α quantifies how much of the orbital energy can be used to unbind the envelope.

We calculate the α_{CE} as

$$\alpha_{\text{CE}}(m) = \frac{\text{BE}(m)}{-\frac{\text{GM}_1\text{M}_2}{2a_{\text{pre-CE}}} + \frac{\text{GM}_2}{2r(m)}} \quad (2)$$

A α_{CE} greater than one implies that just the orbital energy is not sufficient to eject the envelope, and additional sources of energy are required. As can be seen in the shaded region in Fig. 5, we find that successful CE ejection requires values of α_{CE} between 13–17 if we exclude the internal energy and 7.5–9.5 if we include it. These numbers are quite high, and usually imply that the common envelope phase would be unsuccessful, as the core of the star and the compact object would prematurely merge before ejecting the common envelope. Such a merger could potentially give rise to an interesting transient or even the formation of a Thorne-Zytkow object (TZO). However, due to the low likelihood of successful CE ejection, the HMXB 4U1700-37/HD153919 is unlikely to form a double compact object system, and therefore unlikely to be the progenitor of a gravitational wave source.

9. DISCUSSION

1. Very asymmetric HMXB in our own galaxy - well studied observationally, lots of constraints. Helps in charting past history
2. sibling/analogous to GW190814 - insights from galactic binaries to weird low Z GW events

3. common origin - conservative mass transfer + q 1 massive binaries
4. different endings - strength and direction of natal kick - XRB shrinks (seen as XRB, future TZO), GW widens (interacts during RSG, CE ejection, GW driven merger)
5. pathway to forming asymmetric binaries starting from symmetric ones
6. Perhaps Case A is needed, as in Case B helium core is developed and may be unable to explode
7. need successful explosion for both XRB/GW to form NS/low mass BH, and for GW need a strong natal kick as well

10. CONCLUSION

ACKNOWLEDGMENTS

AUTHOR CONTRIBUTIONS

Facilities: HST(STIS), Swift(XRT and UVOT), AAVSO, CTIO:1.3m, CTIO:1.5m, CXO

Software: astropy (Astropy Collaboration et al. 2013, 2018, 2022), Cloudy (G. J. Ferland et al. 2013), Source Extractor (E. Bertin & S. Arnouts 1996)

APPENDIX

REFERENCES

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, ApJ, 935, 167, doi: [10.3847/1538-4357/ac7c74](https://doi.org/10.3847/1538-4357/ac7c74)
- Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393, doi: [10.1051/aas:1996164](https://doi.org/10.1051/aas:1996164)
- Ferland, G. J., Porter, R. L., van Hoof, P. A. M., et al. 2013, RMxAA, 49, 137. <https://arxiv.org/abs/1302.4485>