

Travelling waves in a mathematical model for oncolytic virotherapy

Negar Mohammadnejad, Thomas Hillen *

Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, AB, T6G 2R3, Canada

Received 5 January 2026; revised 24 April 2026; accepted 8 June 2026

Abstract

Oncolytic virotherapy (OVT) is a promising cancer treatment strategy in which engineered viruses selectively infect and destroy tumor cells. Motivated by the biological mechanisms underlying viral spread and tumor invasion into the tissue, we analyze a non-cooperative reaction–diffusion model capturing the invasion of tumor tissue by oncolytic viruses. Using carefully constructed upper and lower solutions together with Schauder’s fixed point theorem, we establish the existence of positive travelling-wave solutions. In particular, we identify a minimal wave speed value $\bar{c}$ such that positive travelling waves exist for all $c \geq \bar{c}$. Moreover, we find a lower value $c_m \leq \bar{c}$ and we show that virus invasion fronts do not exist for wave speeds less than c_m . This analysis highlights parameter regions where the existence of travelling waves remains ambiguous, suggesting new mathematical questions about the propagation of viral treatments through tumor environments.

© 2026 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

Keywords: Oncolytic virotherapy; Reaction-diffusion equations; Travelling wave solutions; Positive travelling wave solutions

1. Introduction

In 1991, the first genetically engineered oncolytic virus (OV)—a thymidine kinase–deficient herpes simplex virus type 1 (HSV-1) mutant—was developed and used to treat malignant glioma in nude mice [20]. Oncolytic virotherapy (OVT) has since emerged as a promising cancer treat-

* Corresponding author.

E-mail addresses: negar3@ualberta.ca (N. Mohammadnejad), thillen@ualberta.ca (T. Hillen).

ment strategy in which oncolytic viruses are introduced into a patient to selectively infect and eliminate tumor cells [18].

Over the last two decades, mathematical models for oncolytic viruses were developed and analysed [38,13,33,6,4,31,12,23,2,1]. Besides clinically relevant conclusions on the administration of oncolytic virotherapy, it turned out that the basic oncolytic virus model has very interesting, and non-trivial, mathematical properties. The model is often formulated as a system of reaction-diffusion equations and it can lead to oscillations, spatial pattern formation, invasion fronts, spiral waves and spatio-temporal chaotic behaviour [26,6,3]. The analysis of these patterns has mostly focussed on numerical simulations. In the present work, we rigorously establish the existence of travelling wave solutions for the basic oncolytic virus reaction–diffusion model, which describes the spatial interactions between uninfected cancer cells C , infected cancer cells I , and free virions V on an unbounded spatial domain.

We notice that the three-component reaction-diffusion system is non-cooperative, hence the standard theory for monotone or cooperative systems (e.g., [8,17,19,36,39,35]) cannot be applied. Instead, we construct appropriate upper and lower solutions and apply Schauder's fixed point theorem to show existence of travelling waves with speeds in an interval $[\bar{c}, \infty)$, and non-existence in an interval $[0, c_m)$ with $c_m \leq \bar{c}$.

Before presenting our analysis, we begin with a brief overview of existing oncolytic virotherapy (OVT) models (Section 1.1), followed by a short introduction to travelling wave methods (Section 1.2). In the sections that follow, we introduce the oncolytic virotherapy model studied here in Section 1.3 and summarize the main results in Section 1.4. We use Section 2 to state the necessary conditions that allow us to construct the upper and lower solutions in preparation for the main existence result. The proof of the main existence theorem is completed in Section 3. In Section 4 we use the negative one-sided Laplace transform to show non-existence for small speeds. Finally, in Section 5 we present numerical simulations of the full PDE system and compare them with our analytical findings.

1.1. Oncolytic virotherapy modelling

Early theoretical work on oncolytic virotherapy (OVT) was shaped by the models of Wodarz and Komarova [37,38,13], who established a general ODE framework capturing key features of tumor growth and viral spread. Their analyses revealed that OVT dynamics fall into fast- and slow-spread regimes, and that treatment outcomes depend sensitively on assumptions about infection kinetics. These studies provided a foundation for later spatial and PDE-based models of tumor–virus interactions. Building on these ideas, Tian [33] introduced a three-variable ordinary differential system tracking healthy cancer cells, infected cells, and free virus, showing how viral burst size can determine therapeutic success. More detailed spatial models soon followed. Pooladvand et al. [26] developed a three-dimensional, radially symmetric PDE model for adenovirus therapy in solid tumors, allowing susceptible cells, infected cells, and free virus to diffuse. Their analysis showed that increasing viral infectivity alone does not guarantee tumor elimination—consistent with experimental observations that virotherapy rarely succeeds as a standalone treatment. Their work helped motivate further spatial modelling, including the diffusion-based system of [2], which examined temporal and spatial infection patterns in the absence of immune effects. Additional computational studies have explored why OVT can fail. Bhatt et al. [4] combined immersed-boundary and Voronoi-based simulations to identify key limiting mechanisms: rapid loss of infected cells (leading to premature viral clearance), emergence of virus-resistant tumor cells, and insufficient viral spread rates. Morselli

et al. [23] used a stochastic agent-based model to examine how spatial constraints and microenvironmental factors shape viral propagation, comparing random versus pressure-driven viral movement and reproducing infection patterns seen in earlier biological and theoretical studies. Although these agent-based and high-dimensional spatial models provide rich biological insight, their computational cost often limits extensive analytical investigation. For this reason, reaction–diffusion models remain valuable for uncovering the spatial structure of viral spread and identifying conditions favouring tumor control or eradication. Baabdulla et al. [2] extend the framework of Pooladvand et al. [26]. Pooladvand's original model—formulated in 3-D radial symmetry—displayed transitions to coexistence states and oscillatory dynamics through a Hopf bifurcation, matching oscillations previously reported in experimental systems. Baabdulla et al. [2] removed the radial symmetry and analyzed the full 2-D and 3-D diffusion problem. They computed invasion speeds numerically, and compared spatial infection profiles. They found complex spatiotemporal behaviours that are often seen in excitable media [14,27]. Their analysis highlighted how spatial heterogeneity shapes viral spread and helps clarify the circumstances under which OVT can suppress—or nearly eradicate—tumor mass.

In the absence of oncolytic viruses (OVs), tumors typically develop an immunosuppressive microenvironment that suppresses the host immune response. The introduction of OVs, however, induces pro-inflammatory signalling within the tumor and can substantially alter tumor–immune dynamics. Motivated by the work of Al-Tuwairqi et al. [1], a reaction–diffusion model was proposed in [21] to capture OVT–immune system interactions. This model extends that of Baabdulla et al. [2] by incorporating a fourth variable Y representing the immune response. Their analysis demonstrated the existence of stacked travelling waves and, through the computation of associated wave speeds, indicated that OVT as a monotherapy is unlikely to achieve complete tumor eradication. This was further explored in [22], where strategies to enhance the efficacy of OVT were investigated, with particular emphasis on the importance of combination treatments involving immunotherapies. These studies highlight that explicitly modelling the immune response is essential for understanding the full therapeutic potential of OVT.

Eftimie et al. [6] developed an ODE model incorporating both lymphoid and peripheral immune compartments to describe the interactions among uninfected and infected tumor cells, memory and effector immune cells, and two viral types (Ad and VSV). Their model successfully reproduced experimental tumor growth and immune-response patterns, identifying conditions under which sustained tumor elimination is possible. Importantly, they showed that viral persistence alone cannot clear the tumor without a coordinated anti-tumor immune response.

Storey et al. [31] further distinguished the roles of innate and adaptive immunity in OVT, noting that the innate response may act too rapidly and clear the virus before sufficient tumor infection occurs. To overcome this limitation, combination therapies pairing OVT with PD-1/PD-L1 checkpoint inhibition have been explored. Such approaches mitigate T-cell exhaustion, enhance immune-mediated tumor clearance, and reduce the viral infectivity threshold required for therapeutic success. Their findings also underscore the importance of treatment scheduling: administering a second viral dose too early can redirect the immune response toward viral, rather than tumor, elimination, thereby reducing overall therapeutic efficacy.

1.2. Travelling waves

Modelling with partial differential equations (PDEs)—and reaction–diffusion equations in particular—has had a profound influence on the mathematical study of biological populations,

with significant applications in cell biology [28,15], ecology [29,16], and epidemiology [40,35]. Many works (e.g., [24,5,34,30]) have investigated travelling waves and spatial dynamics for n -dimensional ($n \geq 1$) reaction–diffusion systems of the form

$$\frac{\partial u_i}{\partial t} = D_i \Delta u_i + f_i(u_1, \dots, u_n), \quad t \geq 0, \quad x \in \mathbb{R}, \quad (1)$$

for $1 \leq i \leq n$. Here $u_i(x, t)$ denotes the density of species of type $i = 1, \dots, n$, D_i denotes their diffusion coefficient, and $f_i(u_1, \dots, u_n)$ describes their birth, death, and other interactions. Such models give rise to travelling wave solutions that represent invasion fronts. The most prominent model arises for one species $n = 1$ and is known as the Fisher–KPP model [24,5]. It is a classical and widely studied prototype for biological invasions and describes the spatiotemporal dynamics of a population whose individuals migrate via linear diffusion and proliferate according to logistic-type growth.

In cancer biology, reaction–diffusion frameworks have been especially useful in describing tumor invasion, particularly when the spread of malignant cells is coupled to degradation of surrounding tissue [7].

The system (1) is called *cooperative* if each $f_i(u)$ is non-decreasing in all components of u , with the possible exception of the i th one. Cooperative systems have been extensively studied, and a general travelling wave theory is available in this setting [8,36,17].

For *non-cooperative* systems—characterized by kinetic Jacobians that include negative off-diagonal entries—this general theory does not apply. The model studied in this work (system (3)) is non-cooperative and the above mentioned theory is not available. Instead we follow the framework of [32] to construct upper and lower solutions that connect correctly to the left boundary and remain uniformly bounded throughout the domain. This framework then enables the application of Schauder’s fixed-point theorem, yielding the existence of travelling–wave solution for speeds in an interval $[\bar{c}, \infty)$.

The non-monotonicity of our system introduces significant technical challenges and restricts us to parameter regimes where existence can be rigorously established. When determining the minimal wave speed, we encounter parameter regions in which the existence—or nonexistence—of travelling waves cannot be conclusively determined. Such indeterminate parameter zones are common in travelling wave analysis and motivate future work aimed at resolving these “hazy” regimes where classical arguments do not decisively settle the question of the wave’s existence.

Reaction–diffusion systems can exhibit a phenomenon known as *stacked waves* [9], which refers to the occurrence of invasion fronts connecting different steady states and travelling at different speeds. This is also the case for our oncolytic virus model [21]. In the absence of virus and infected cells, the cancer population alone satisfies a Fisher–KPP equation

$$C_t = D \Delta C + C(1 - C),$$

which admits invasion fronts with speed $c \geq 2\sqrt{D}$ [24]. Once the virus is introduced, an additional invasion front emerges, corresponding to the propagation of the virus into the cancer population, which in general travels at a different speed. It is this virus invasion wave that is the primary focus of our analysis.

1.3. The main oncolytic virus system

In this work, we will study the A. Baabdulla [2] reaction diffusion model on $\mathbb{R}^n$. The proposed model from [2] is given by

$$\begin{aligned}\frac{dC}{dt} &= D\Delta C + rC \left(1 - \frac{C+I}{L}\right) - \beta CV, \\ \frac{dI}{dt} &= D\Delta I + \beta CV - \eta I, \\ \frac{dV}{dt} &= D_V\Delta V + \eta bI - \omega V.\end{aligned}\tag{2}$$

In this formulation, C denotes the population of uninfected cancer cells, I the population of infected cancer cells, and V the concentration of free oncolytic virus.

We consider system (2) together with the initial conditions

$$C(x, 0) = C_0 > 0, \quad I(x, 0) = I_0 = 0, \quad V(x, 0) = V_0 > 0,$$

The boundary behavior at $|x| \rightarrow \infty$ will be specified later when we introduce travelling wave coordinates.

In the system of equations (2), the diffusion coefficients of the susceptible and infected tumor cell populations are denoted by D , while the free virus diffuses with coefficient D_V . The operator Δ denotes the Laplacian, i.e. the sum of all second-order spatial derivatives.

In the first equation, the second term represents the logistic proliferation of the uninfected cancer cells in the absence of viral therapy, while the third term, $-\beta CV$, models the loss of susceptible tumor cells due to infection by free virus. Free virions interact with uninfected cancer cells and transfer them into the infected class. Consequently, the same interaction appears with opposite sign in the second equation as the term $+\beta CV$, which describes the influx of cells into the infected population. Thus, the second equation accounts for the accumulation of infected tumor cells through infection, while the term $-\eta I$ represents their removal due to lysis.

In the third equation, the term ηbI incorporates the burst size b , describing the number of viral particles released upon lysis of infected cells, and the final term, $-\omega V$, corresponds to the natural clearance rate of free virus.

To simplify our analysis, we apply the non-dimensionalization method by considering

$$\hat{C} = \frac{C}{L}, \quad \hat{I} = \frac{I}{L}, \quad \hat{V} = \frac{\beta V}{r}, \quad \hat{t} = rt, \quad \hat{x} = x\sqrt{\frac{r}{D_V}}, \quad \text{and} \quad \hat{D} = \frac{D}{D_V}.$$

After dropping the hats, we get the non-dimensional system:

$$\begin{aligned}\frac{dC}{dt} &= D\Delta C + C(1 - C - I) - CV, \\ \frac{dI}{dt} &= D\Delta I + CV - aI, \\ \frac{dV}{dt} &= \Delta V + \theta I - \gamma V,\end{aligned}\tag{3}$$




Fig. 1. Bifurcation diagram of the model (4) with bifurcation parameter θ . The equilibrium E_1 loses stability at $\theta \approx 12.7$, while E_2 becomes biologically relevant and remains stable up to the Hopf bifurcation at $\theta_H \approx 220$. Beyond this point, the system exhibits oscillatory dynamics. The parameter values used here are $a = 0.96$ and $\gamma = \frac{40}{3}$. The circles mark the θ values selected in Section 5 for numerical simulations.

where

$$a = \frac{\eta}{r}, \quad \theta = \frac{\eta b \beta L}{r^2}, \quad \gamma = \frac{\omega}{r}.$$

We initially examine the aforementioned model in the spatially homogeneous scenario, where all spatial dependencies are disregarded. Consequently, the system governing $C(t)$, $I(t)$, $V(t)$ simplifies to:

$$\begin{aligned} \frac{dC}{dt} &= C(1 - C - I) - CV, \\ \frac{dI}{dt} &= CV - aI, \\ \frac{dV}{dt} &= \theta I - \gamma V. \end{aligned} \tag{4}$$

We obtain the following equilibrium points for system (4)

$$E_0 = (0, 0, 0),$$

$$E_1 = (1, 0, 0),$$

$$E_2 = (C^*, I^*, V^*) = \left(\frac{a\gamma}{\theta}, \frac{\gamma(\theta - a\gamma)}{\theta(\gamma + \theta)}, \frac{\theta - a\gamma}{\gamma + \theta} \right).$$

The analysis of the equilibrium points E_0 , E_1 , and E_2 was carried out in detail in [2]. We show a bifurcation diagram in Fig. 1 with θ as bifurcation parameter. For low values of θ , the cancer only state E_1 is stable, and by increasing θ , it undergoes a transcritical bifurcation at $\theta_t = \gamma a$, and the coexistence state, E_2 , becomes biologically relevant and stable. Upon further increase of θ , we obtain a Hopf bifurcation at θ_H with periodic orbits arising for $\theta > \theta_H$. For our further analysis, we assume the coexistence equilibrium exists, hence we assume throughout:

$$\theta > \gamma a. \tag{5}$$

1.4. Main results

We consider self-similar travelling wave solutions of (3) of the form

$$(C(x, t), I(x, t), V(x, t)) = (C(\xi), I(\xi), V(\xi)),$$

with $\xi = x + ct$, where c denotes the wave speed of an invasion front travelling to the left for $c > 0$. We consider a virus invasion front propagating into an established tumor, subject to the boundary conditions

$$(C(-\infty), I(-\infty), V(-\infty)) = (1, 0, 0) \quad \text{and} \quad \|(C(\infty), V(\infty), I(\infty))\|_\infty < \infty. \quad (6)$$

We call an invasion front to be *nontrivial*, if none of the components are identical to zero. The main existence result for nontrivial invasion fronts is as follows.

Theorem 1.1. *Assume (5). There exists a constant $\bar{c} > 0$ such that for each $c \geq \bar{c}$, the system (3), (6) admits a self-similar nontrivial travelling wave solution $(C(\xi), I(\xi), V(\xi))$ with $\xi = x + ct$. The solution connects to $(1, 0, 0)$ as $\xi \rightarrow -\infty$ and remains bounded as $\xi \rightarrow +\infty$. The explicit form of $\bar{c}$ is given in formula (34) in the proof.*

The proof relies on a carefully balanced construction of upper and lower solutions, presented in Section 2. The proof of Theorem 1.1 is then carried out in Section 3 via a Schauder fixed point argument on a suitably chosen set Γ .

We also establish non-existence for small wave speeds.

Theorem 1.2. *Assume (5). For $0 < c < c_m$, there exists no nontrivial travelling front of (3) satisfying the boundary conditions (6), where c_m is defined in (20).*

The non-existence of travelling waves for $c < c_m$ is established in Section 4 using a negative one-sided Laplace transform method. This approach was developed in [40,41] for a related travelling wave problem.

Remark 1.3.

1. We find that in most biologically relevant cases we have $c_m = \bar{c}$, and Theorem 1.1 and 1.2 give a complete result. However, in extreme cases, as outlined below, we can also have $c_m < \bar{c}$ and there is a gap $c \in (c_m, \bar{c})$, where the existence of invasion fronts is an open question.
2. For $c < -2\sqrt{D}$, invasion fronts of (3) with boundary conditions (6) do exist in our model. These correspond to cancer-only fronts travelling to the right at the Fisher–KPP wave speed (see also Appendix A). Note that c is negative in this case, since we consider the wave variable $x + ct$.

2. Construction of upper- and lower solutions

Since our interest lies in describing the invasion of the viral treatment into an established tumor, it is convenient to shift the equilibrium corresponding to the untreated cancer to the origin. To do so, we introduce the change of variables

$$B := 1 - C.$$

Under this transformation, the relevant equilibrium E_1 becomes $(0, 0, 0)$, allowing us to apply standard techniques for studying invasion into the “disease-free” state $(0, 0, 0)$. Substituting $C = 1 - B$ into (3) yields the equivalent system

$$\begin{aligned}\frac{dB}{dt} &= D\Delta B + (B-1)(B-I) + (1-B)V, \\ \frac{dI}{dt} &= D\Delta I + (1-B)V - aI, \\ \frac{dV}{dt} &= \Delta V + \theta I - \gamma V.\end{aligned}\tag{7}$$

To derive the travelling wave equations, we introduce the coordinates $\xi = x + ct$. A travelling wave solution has the form

$$(B(x, t), I(x, t), V(x, t)) = (B(\xi), I(\xi), V(\xi)),$$

and substituting this into (7) leads to the ODE system

$$\begin{aligned}cB' &= DB'' - B + B^2 - BI - BV + I + V, \\ cI' &= DI'' - aI + (1-B)V, \\ cV' &= V'' + \theta I - \gamma V.\end{aligned}\tag{8}$$

The boundary conditions require special attention. At the leading edge of the wave, corresponding to the pre-invasion cancer state, we impose

$$(C(-\infty), I(-\infty), V(-\infty)) = (1, 0, 0).$$

Behind the wave, i.e. as $\xi \rightarrow \infty$, several behaviours are possible. The solution may converge to the equilibrium E_2 , but as illustrated in Fig. 1, E_2 undergoes a Hopf bifurcation for sufficiently large θ . In such cases, the wave approaches a periodic orbit surrounding E_2 rather than the equilibrium itself. Accordingly, we impose the general boundedness condition

$$\|(C(\infty), I(\infty), V(\infty))\| < \infty.$$

The associated boundary conditions, expressed in (B, I, V) variables, are

$$(B(-\infty), I(-\infty), V(-\infty)) = (0, 0, 0), \quad \text{and} \quad \|(B(\infty), I(\infty), V(\infty))\| < \infty.\tag{9}$$

From this point forward, we work exclusively with the transformed system (8), as it simplifies the analytical treatment of the travelling wave problem.

The system (8) is a second-order inhomogeneous ODE and we find an alternative formulation using the method of variation of parameters. This transformation has shown to be useful for travelling wave analysis in the work of Fang and Zhao [8]. We define $U := (B, I, V)$, the matrix $\mathcal{D}$, and the function f as

$$f(U) = f(B, I, V) = \begin{pmatrix} f_1(B, I, V) \\ f_2(B, I, V) \\ f_3(B, I, V) \end{pmatrix} = \begin{pmatrix} -B + B^2 - BI - BV + I + V \\ -aI + (1-B)V \\ \theta I - \gamma V \end{pmatrix},$$

$$\mathcal{D} := \begin{pmatrix} D & 0 & 0 \\ 0 & D & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (10)$$

so system (8) can be written as

$$\mathcal{D}U'' - cU' = -f(U). \quad (11)$$

Let

$$F(U) = \begin{pmatrix} F_1(U) \\ F_2(U) \\ F_3(U) \end{pmatrix} = \alpha U + f(U),$$

where α is a positive free parameter. This definition allows us to rewrite (11) as

$$\mathcal{D}U'' - cU' - \alpha U = -F(U).$$

We apply the variation of parameters such that

$$B(\xi) = \frac{1}{\sqrt{c^2 + 4\alpha D}} \times \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} F_1(B(\eta), I(\eta), V(\eta)) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} F_1(B(\eta), I(\eta), V(\eta)) d\eta \right), \quad (12)$$

where

$$\delta_1 = \frac{c - \sqrt{c^2 + 4\alpha D}}{2D}, \quad \delta_2 = \frac{c + \sqrt{c^2 + 4\alpha D}}{2D}. \quad (13)$$

$$I(\xi) = \frac{1}{\sqrt{c^2 + 4\alpha D}} \times \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} F_2(B(\eta), I(\eta), V(\eta)) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\sigma_2} F_2(B(\eta), I(\eta), V(\eta)) d\eta \right), \quad (14)$$

where

$$\sigma_1 = \frac{c - \sqrt{c^2 + 4\alpha D}}{2D}, \quad \sigma_2 = \frac{c + \sqrt{c^2 + 4\alpha D}}{2D}. \quad (15)$$

$$V(\xi) = \frac{1}{\sqrt{c^2 + 4\alpha}} \times \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} F_3(B(\eta), I(\eta), V(\eta)) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\zeta_2} F_3(B(\eta), I(\eta), V(\eta)) d\eta \right), \quad (16)$$

where

$$\zeta_1 = \frac{c - \sqrt{c^2 + 4\alpha}}{2}, \quad \zeta_2 = \frac{c + \sqrt{c^2 + 4\alpha}}{2}. \quad (17)$$

These integral formulations will allow us to define an operator framework, the fixed points of which correspond to travelling wave solutions.

2.1. Dispersion relation for the invasion wave speed

We first analyze the linearization of (8) in order to characterize the invasion speed and the exponential growth rate at the leading edge of the wave. This analysis leads to a set of conditions that must be satisfied for travelling wave solutions to exist.

We linearize the system (8) about the $(0, 0, 0)$ equilibrium

$$cB' = DB'' - B + I + V,$$

$$cI' = DI'' - aI + V,$$

$$cV' = V'' + \theta I - \gamma V.$$

At the leading edge of the wave, we assume exponential growth as $B = v_b e^{\rho\xi}$, $I = v_i e^{\rho\xi}$, and $V = v_v e^{\rho\xi}$ for a constant vector $v = (v_b, v_i, v_v)$, and a growth rate $\rho > 0$. Substituting the solution into the above equation yields the algebraic system

$$c\rho v = \underbrace{\begin{pmatrix} D & 0 & 0 \\ 0 & D & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{=: \mathcal{D}} \rho^2 v + \underbrace{\begin{pmatrix} -1 & 1 & 1 \\ 0 & -a & 1 \\ 0 & \theta & -\gamma \end{pmatrix}}_{=: f'(0,0,0)} v,$$

where $f'(0, 0, 0)$ is the Jacobian of f from (10) at equilibrium $(0, 0, 0)$. Defining the matrix $A(\rho) = \mathcal{D}\rho^2 + f'(0, 0, 0)$, we retrieve an eigenvalue problem

$$(A(\rho) - c\rho\mathbb{I})v = 0,$$

with the eigenvalue

$$\lambda = c\rho.$$

We explicitly compute the eigenvalues and the eigenvectors of the matrix A . The matrix A is given by

$$A(\rho) = \begin{pmatrix} D\rho^2 - 1 & +1 & +1 \\ 0 & D\rho^2 - a & 1 \\ 0 & \theta & \rho^2 - \gamma \end{pmatrix}.$$

Thus, the eigenvalues are given by

$$\begin{aligned} \lambda_1 &= D\rho^2 - 1, \\ \lambda_{2,3} &= \frac{\rho^2 + D\rho^2 - a - \gamma \pm \sqrt{(\rho^2 + D\rho^2 - a - \gamma)^2 - 4(D\rho^4 - D\rho^2\gamma - a\rho^2 + a\gamma - \theta)}}{2} \\ &= \frac{\rho^2 + D\rho^2 - a - \gamma \pm \sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2 + 4\theta}}{2}, \end{aligned} \quad (18)$$

and the eigenvectors

$$\begin{aligned} v_1 &= \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} \frac{(\lambda_2 + 1 + a - D\rho^2)}{(\lambda_2 + a - D\rho^2)(\lambda_2 + 1 - D\rho^2)} \\ 1 \\ \frac{\lambda_2 + a - D\rho^2}{1} \end{pmatrix}, \\ v_3 &= \begin{pmatrix} \frac{(\lambda_3 + 1 + a - D\rho^2)}{(\lambda_3 + a - D\rho^2)(\lambda_3 + 1 - D\rho^2)} \\ 1 \\ \frac{\lambda_3 + a - D\rho^2}{1} \end{pmatrix}. \end{aligned} \quad (19)$$

Remark 2.1. The eigenvalue $\lambda_1(\rho)$ in (18) has the associated eigenvector $v_1 = (1, 0, 0)$, which corresponds to a direction purely in the cancer component: it involves neither infected cells nor virus. In fact, λ_1 is determined entirely by the B -equation and represents the spatial decay rate of the stable manifold of the classical Fisher–KPP equation at the equilibrium $C = 1$ (equivalently, $B = 0$). It is well known that this mode cannot generate a bounded travelling wave solution, and we therefore exclude this case from further consideration (see also Appendix A).

Since our objective is to analyse viral invasion into a cancer population, we restrict attention to the remaining two eigenvalues, $\lambda_2(\rho)$ and $\lambda_3(\rho)$. Because $\lambda_2 > \lambda_3$ for all $\rho > 0$, the relevant branch for constructing travelling-wave solutions is λ_2 .

The wave speed associated with the eigenvalue $\lambda_2(\rho)$ is

$$S(\rho) = \frac{\lambda_2(\rho)}{\rho} = \frac{\rho^2 + D\rho^2 - a - \gamma}{2\rho} + \frac{\sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2 + 4\theta}}{2\rho}.$$

We then introduce the quantity

$$c_m = \inf_{\rho > 0} S(\rho), \quad (20)$$

which represents the minimum of this speed curve.

Lemma 2.2. *The speed curve $S(\rho)$ decreases as ρ is close to zero and tends to infinity as $\rho \rightarrow 0$. We have $\lim_{\rho \rightarrow \infty} S(\rho) = +\infty$ and $S(\rho)$ has at least one global minimum.*

Proof. The limit $\lim_{\rho \rightarrow 0} S(\rho) = \infty$ follows by applying *L'Hôpital's rule* to the defining expression of $S(\rho)$. Moreover, since $\lim_{\rho \rightarrow \infty} S(\rho) = \infty$ and S is continuous on $(0, \infty)$, it follows that S admits at least one global minimizer $\rho_m \in (0, \infty)$ such that

$$S(\rho_m) = \inf_{\rho > 0} S(\rho) =: c_m. \quad (21)$$

If multiple minimizers exist, we select the smallest ρ and denote it again by

$$\rho_m = \min\{\rho_i : S(\rho_i) = c_m\}. \quad (22)$$

Hence, the minimal value c_m is attained at $\rho = \rho_m$. $\square$

In the following lemma, we show that λ_2 is positive. This will show that the function $S(\rho)$ is positive and the minimum wave speed has positive values.

Lemma 2.3. *Assume (5). The eigenvalue $\lambda_2(\rho)$ is positive for $\rho > 0$.*

Proof. Using $\theta > a\gamma$, we prove that

$$\begin{aligned} \lambda_2 &= \frac{\rho^2 + D\rho^2 - a - \gamma}{2} + \frac{\sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2 + 4\theta}}{2} \\ &\geq \frac{\rho^2 + D\rho^2 - a - \gamma}{2} + \frac{\sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2}}{2}. \end{aligned} \quad (23)$$

Thus, we need to prove

$$\frac{\rho^2 + D\rho^2 - a - \gamma + \sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2 + 4a\gamma}}{2} > 0 \quad \text{for } \rho > 0.$$

We first note that when $D = 1$, we have

$$\lambda_2 > \frac{\rho^2 + \rho^2 - a - \gamma + \sqrt{(a - \gamma)^2 + 4a\gamma}}{2} = \frac{2\rho^2 - a - \gamma + \sqrt{(a + \gamma)^2}}{2} = \rho^2 > 0. \quad (24)$$

For $D \neq 1$, two possible cases exist that need to be handled separately.

1. $\rho^2 + D\rho^2 - a - \gamma > 0$,
2. $\rho^2 + D\rho^2 - a - \gamma < 0$.

In the first case, both terms in (23) are positive, which implies directly that $\lambda_2 > 0$. In the second case, noting that $D > 0$, we can rewrite the inequality as

$$\rho^2 + D\rho^2 - a - \gamma < 0 \quad \Longleftrightarrow \quad 0 < D\rho^2 < a + \gamma - \rho^2. \quad (25)$$

Under the assumption (25), we now determine conditions guaranteeing $\lambda_2 > 0$.

To ensure that

$$\lambda_2 = \frac{\sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2 + 4a\gamma} + (\rho^2 + D\rho^2 - a - \gamma)}{2} > 0,$$

we require

$$\sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2 + 4a\gamma} > -(\rho^2 + D\rho^2 - a - \gamma).$$

Squaring both sides yields

$$(\rho^2 - D\rho^2 + a - \gamma)^2 + 4a\gamma > (\rho^2 + D\rho^2 - a - \gamma)^2.$$

Expanding and simplifying, we obtain the required condition

$$a + D\gamma - D\rho^2 > 0 \quad \Longleftrightarrow \quad a + D(\gamma - \rho^2) > 0. \quad (26)$$

We now show that this inequality holds for all $\rho > 0$ in the two cases $D < 1$ and $D \geq 1$.

Since (25) implies $a + \gamma - \rho^2 > 0$, if $D < 1$, then immediately

$$a + D(\gamma - \rho^2) > 0. \quad (27)$$

If $D \geq 1$, then from (25) we have

$$D\rho^2 < a + \gamma - \rho^2.$$

Since $D\rho^2$ is smaller than $a + \gamma - \rho^2$, substituting the larger value into (26) yields

$$0 < a + D\gamma - (a + \gamma - \rho^2) < a + D\gamma - D\rho^2.$$

Simplifying, we obtain

$$a + D\gamma - a - \gamma + \rho^2 = \gamma(D - 1) + \rho^2 > 0, \quad (28)$$

which is always positive for $D \geq 1$.

Thus, by (24), (27) and (28), we conclude that

$$\lambda_2 > 0 \quad \text{for all } \rho > 0. \quad \square$$

We also prove an estimate for λ_2 in the following Lemma, which will be used in the proof of the main theorem.

Lemma 2.4. Assume (5). For all $\rho > 0$, we have that $\lambda_2 > D\rho^2 - a$.

Proof. Using $\theta > a\gamma$, we obtain

$$\begin{aligned}\lambda_2 &= \frac{\rho^2 + D\rho^2 - a - \gamma + \sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2 + 4\theta}}{2} \\ &> \frac{\rho^2 + D\rho^2 - a - \gamma + \sqrt{(\rho^2 - D\rho^2 + a - \gamma)^2}}{2}.\end{aligned}$$

We consider two cases. If we have $\rho^2 - D\rho^2 + a - \gamma \geq 0$ then

$$\lambda_2 \geq \frac{\rho^2 + D\rho^2 - a - \gamma + (\rho^2 - D\rho^2 + a - \gamma)}{2} = \rho^2 - \gamma \geq D\rho^2 - a.$$

Secondly, if $\rho^2 - D\rho^2 + a - \gamma \leq 0$,

$$\lambda_2 \geq \frac{\rho^2 + D\rho^2 - a - \gamma - (\rho^2 - D\rho^2 + a - \gamma)}{2} = D\rho^2 - a. \quad \square$$

2.2. Conditions for the construction of upper and lower solutions

We will use the eigenvector v_2 from (19) to define the upper and lower solutions. For this construction to work, we must ensure that all three components of v_2 are positive.

Lemma 2.5. Assume (5). Then all components of v_2 are positive in the following cases:

1. $a \leq 1$.
2. $a > 1$ and the curves

$$S(\rho) \quad \text{and} \quad H_1(\rho) = D\rho - \frac{1}{\rho}$$

do not intersect.

3. $a > 1$ and the curves $S(\rho)$ and $H_1(\rho)$ intersect, and the smallest intersection point ρ_* satisfies $\rho_* > \rho_m$.
4. $a > 1$ and the curves $S(\rho)$ and $H_1(\rho)$ intersect, and the smallest intersection point ρ_* satisfies $\rho_* < \rho_m$.

In cases (1)–(3) we have $\rho \leq \rho_m$, whereas in case (4) we restrict to $\rho \leq \rho_*$.

Proof. For the first component of v_2 in (19) we have

$$\frac{\lambda_2 + a + 1 - D\rho^2}{(\lambda_2 + a - D\rho^2)(\lambda_2 + a - D\rho^2)}.$$

Lemma 2.4 guarantees that

$$\lambda_2 + a - D\rho^2 > 0,$$

which in turn implies

$$\lambda_2 + a + 1 - D\rho^2 > 0.$$

Therefore, to ensure positivity of the first component, we further require

$$\lambda_2 + 1 - D\rho^2 > 0. \quad (29)$$

The second component of v_2 in (19) is always positive by Lemma 2.4.

To determine when condition (29) holds, we distinguish two cases depending on the value of a .

Case 1: $a \leq 1$. In this case, we have

$$\lambda_2 + 1 - D\rho^2 \geq \lambda_2 + a - D\rho^2 > 0,$$

and the positivity requirement is automatically satisfied. Fig. 2a illustrates an example of this case by plotting the speed curve $S(\rho) = \frac{\lambda_2(\rho)}{\rho}$ together with the curves $H_1(\rho) = D\rho - \frac{1}{\rho}$ and $H_2(\rho) = D\rho - \frac{a}{\rho}$. When $a \leq 1$, the graph of H_2 lies strictly above that of H_1 for all $\rho > 0$, and therefore neither curve intersects $S(\rho)$. This confirms that the inequality $\lambda_2 > D\rho^2 - 1$ holds, and hence the components of v_2 remain positive throughout this case.

Case 2: $a > 1$. When $a > 1$, there may exist an intersection point between $H_1(\rho)$ and $S(\rho)$. If no such intersection occurs, then no additional restriction is required. Otherwise, suppose that an intersection exists, and define the smallest intersection point (ρ_*, c_*) by

$$\rho_* := \{\rho > 0; H_1(\rho) = S(\rho)\}, \quad c_* := S(\rho_*). \quad (30)$$

The inequality

$$\lambda_2 + 1 - D\rho^2 > 0$$

may still hold if $\rho_m < \rho_*$. However, if the smallest intersection point satisfies $\rho_* < \rho_m$, then the inequality

$$\lambda_2 + 1 - D\rho^2 > 0$$

fails for $\rho \in (\rho_*, \rho_m)$. In this case, the first component of v_2 becomes negative. Consequently, the construction of an upper solution is no longer valid, and the parameter regime $\rho \in (\rho_*, \rho_m)$ remains undecided. $\square$

We obtain two additional restrictions arising from the construction of upper and lower solutions, which will be used later in (45). Specifically, we require

$$c \geq D\rho =: H_3(\rho),$$

and

$$c \leq \frac{-a}{\rho} + \frac{\theta}{\gamma\rho} + D\rho =: H_4(\rho). \quad (31)$$

If $H_3(\rho)$ intersects $S(\rho)$ at a point $(\check{\rho}, \check{c})$, defined by

$$\check{\rho} := \{\rho > 0; H_3(\rho) = S(\rho)\}, \quad \check{c} := S(\check{\rho}), \quad (32)$$

then the admissible values of ρ satisfy $\rho \leq \check{\rho}$.

Similarly, if $H_4(\rho)$ intersects $S(\rho)$ at a point $(\tilde{\rho}, \tilde{c})$, defined by

$$\tilde{\rho} := \{\rho > 0; H_4(\rho) = S(\rho)\}, \quad \tilde{c} := S(\tilde{\rho}), \quad (33)$$

then the admissible values of ρ satisfy $\rho \leq \tilde{\rho}$.

Combining all the above restrictions (30), (32), and (33), the values of ρ for which we can establish the existence of positive travelling wave solutions satisfy

$$0 < \rho \leq \bar{\rho},$$

where

$$\bar{\rho} := \min\{\rho_m, \rho_*, \check{\rho}, \tilde{\rho}\}, \quad \bar{c} := S(\bar{\rho}). \quad (34)$$

Here ρ_m is defined in (22), ρ_* in (30), $\check{\rho}$ in (32), and $\tilde{\rho}$ in (33).

In Fig. 2, we present four different cases illustrating the relative alignment of the critical points ρ_m , ρ_* , $\check{\rho}$, and $\tilde{\rho}$. Figs. 2a and 2c display the curves $S(\rho)$, $H_1(\rho)$, $H_2(\rho)$, $H_3(\rho)$, and $H_4(\rho)$ in the parameter regime $a > 1$. In Fig. 2a, since $a > 1$, the curve $H_1(\rho)$ lies above $H_2(\rho)$. However, there is no intersection between $H_1(\rho)$ and $S(\rho)$ for $\rho < \rho_m$. Similarly, $H_3(\rho)$ and $S(\rho)$ do not intersect for $\rho < \rho_m$. The curve $H_4(\rho)$ intersects $S(\rho)$ for $\rho > \rho_m$, and the point (ρ_m, c_m) lies within the shaded region. Consequently, in this case we have $\bar{\rho} = \rho_m$. The value of γ used in Figs. 2a and 2c is taken from [2] and equals $\gamma = 40/3$. By contrast, Friedman et al. [10] reported a viral clearance rate of $\gamma = 1.25$. This smaller value of γ yields a less restrictive shaded region, as illustrated in Figs. 2b and 2d. Fig. 2b depicts a possible, though biologically unrealistic, parameter regime in which the curve $H_1(\rho)$ intersects $S(\rho)$, as does $H_3(\rho)$. In this situation, the smallest of these intersection points lying within the shaded region determines $\bar{\rho}$.

Figs. 2b and 2d correspond to the case $a < 1$. In both panels, the curve $H_2(\rho)$ lies strictly above $H_1(\rho)$, and consequently there are no intersections between $S(\rho)$ and $H_1(\rho)$. The effect of varying γ is particularly evident in these latter two cases. As shown in Fig. 2b, for larger values of γ the region between $H_4(\rho)$ and $S(\rho)$ becomes more restrictive, and $\bar{\rho}$ occurs before ρ_m , so that $\bar{\rho} = \tilde{\rho}$. In contrast, for smaller values of γ , $\bar{\rho}$ occurs beyond ρ_m , and hence $\bar{\rho} = \rho_m$.

Before stating the main theorem, we analyze the asymptotic behaviour of the curves $S(\rho)$ and $H_4(\rho)$ as $\rho \rightarrow 0^+$. As illustrated in Fig. 2, whenever these two curves intersect at a point $(\bar{\rho}, \bar{c})$, $\bar{\rho} < \rho_m$, no further intersections occur for $0 < \rho < \bar{\rho}$. To justify this observation analytically, we examine the leading-order behaviour of both curves for small ρ .




Fig. 2. Numerical illustration of $S(\rho)$, and its minimum value indicated by the orange ball, and $H_1(\rho)$, $H_2(\rho)$, $H_3(\rho)$, and $H_4(\rho)$ for different parameter values. The parameter values have been chosen to illustrate each of the regimes and are given by: (a) $a = \frac{10}{3}$, $\gamma = \frac{40}{3}$, $\theta = 250$, $D = 0.025$ (b) $a = 0.96$, $\gamma = \frac{40}{3}$, $\theta = 50$, $D = 0.025$, (c) $a = 5$, $\gamma = 2$, $\theta = 17$, $D = 6$ (d) $a = 0.96$, $\gamma = 1.25$, $\theta = 17$, $D = 0.025$. Note that case (c) may not arise in biologically relevant cases. (For interpretation of the colours in the figure(s), the reader is referred to the web version of this article.)

Lemma 2.6. Under assumption (5), for ρ sufficiently small we have

$$H_4(\rho) > S(\rho).$$

Proof. For ρ sufficiently small, we may neglect all terms of order $\mathcal{O}(\rho)$ or higher. Under this approximation, $H_4(\rho)$ reduces to

$$H_4(\rho) \sim \frac{1}{\rho} \left(\frac{\theta}{\gamma} - a \right),$$

while $S(\rho)$ becomes

$$S(\rho) \sim \frac{1}{\rho} \left(-\frac{a}{2} - \frac{\gamma}{2} + \frac{1}{2} \sqrt{(a - \gamma)^2 + 4\theta} \right).$$

Since both expressions scale like $\mathcal{O}(1/\rho)$, the ordering of the two curves for small ρ is determined by comparing their coefficients. To ensure that $H_4(\rho)$ lies strictly above $S(\rho)$ as $\rho \rightarrow 0^+$, we require

$$\frac{\theta}{\gamma} - a > -\frac{a}{2} - \frac{\gamma}{2} + \frac{1}{2}\sqrt{(a-\gamma)^2 + 4\theta}. \quad (35)$$

By simplifying (35) and squaring both sides, we obtain the equivalent condition

$$\left(\frac{\theta}{\gamma} - \frac{a}{2} + \frac{\gamma}{2}\right)^2 > \frac{1}{4}((a-\gamma)^2 + 4\theta),$$

which expands to

$$\frac{\theta^2}{\gamma^2} + \frac{a^2}{4} + \frac{\gamma^2}{4} - \frac{\theta a}{\gamma} + \theta - \frac{a\gamma}{2} > \frac{a^2}{4} + \frac{\gamma^2}{4} + \theta - \frac{a\gamma}{2}.$$

Cancelling common terms yields the condition

$$\theta \left(\frac{\theta}{\gamma^2} - \frac{a}{\gamma} \right) > 0. \quad (36)$$

Since our standing assumption is $\theta > a\gamma$, the right-hand side of (36) satisfies

$$\frac{\theta}{\gamma} \left(\frac{\theta}{\gamma} - a \right) > 0.$$

Thus condition (36) holds automatically. Consequently, we conclude that

$$H_4(\rho) > S(\rho) \quad \text{for all sufficiently small } \rho > 0,$$

which confirms that the two curves cannot intersect again as $\rho \rightarrow 0^+$. $\square$

Note that $H_4(\rho)$ is convex for $\rho > 0$, since

$$H_4''(\rho) = \frac{2\left(\frac{\theta}{\gamma} - a\right)}{\rho^3} > 0 \quad (\rho > 0).$$

Because of the algebraic complexity of the function $S(\rho)$, establishing its convexity on the interval $0 < \rho < \rho_m$ is not straightforward. Nevertheless, by Lemma 2.6, we know that $H_4(\rho) > S(\rho)$ as $\rho \rightarrow 0^+$. Although our numerical simulations did not reveal any secondary intersections between the graphs of $H_4(\rho)$ and $S(\rho)$, we adopt a conservative formulation and define

$$\tilde{\rho} = \min_{0 < \rho < \rho_m} \{\rho : H_4(\rho) = S(\rho)\},$$

that is, $\tilde{\rho}$ is the smallest positive value of ρ at which the two curves intersect.




Fig. 3. Left: Wave-speed function $S(\rho)$ together with the values $\bar{c}$, c , and c_ϵ . The point $(\bar{\rho}, \bar{c})$ is marked in red. The points $\rho = \bar{\rho} - \tau$ and $\rho_\epsilon = \rho + \epsilon$, with $\tau = 1$ and $\epsilon = 0.5$, are also shown, along with their corresponding wave speeds (c, c_ϵ) . Right: Sketch of the upper solutions and their transition points $\bar{\xi}_1$, $\bar{\xi}_2$, and $\bar{\xi}_3$.

2.3. Upper and lower solutions

For the construction of upper and lower solutions, note that for any $c > \bar{c}$ there exists a $\rho \in (0, \bar{\rho})$ such that $S(\rho) = c$, where $\bar{\rho}$ is defined in (34). Moreover, since $S(\rho)$ is increasing on $(0, \bar{\rho}]$, we have $S(\bar{\rho}) = \bar{c} < c$. For $\epsilon > 0$, define

$$c_\epsilon := S(\rho + \epsilon).$$

If ϵ is chosen sufficiently small, then

$$\bar{c} < c_\epsilon < c.$$

We now assume that $\lambda_2(\rho)$ denotes the eigenvalue given in (18), which is associated with the wave speed c and the decay rate ρ .

The wave-speed curve $S(\rho)$ is shown in Fig. 3 (left panel). The pair $(\bar{\rho}, \bar{c})$ is indicated on the curve by an orange marker. We also introduce the points

$$\rho := \bar{\rho} - \tau, \quad \rho_\epsilon := \rho + \epsilon. \quad (37)$$

Here $\tau > 0$ is arbitrary, chosen so that $\bar{\rho} - \tau > 0$, and $\epsilon > 0$ is sufficiently small to ensure that

$$\rho < \rho_\epsilon < \bar{\rho}.$$

We now define our upper and lower solution functions. The upper solutions $(\bar{B}, \bar{I}, \bar{V})$ are given by

$$\begin{aligned} \bar{B}(\xi) &= \min \left\{ e^{\rho\xi} \frac{(c\rho + a + 1 - D\rho^2)}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)}, 1 \right\}, \\ \bar{I}(\xi) &= \min \left\{ e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2}, 1 \right\}, \\ \bar{V}(\xi) &= \min \left\{ e^{\rho\xi}, \frac{\theta}{\gamma} \right\}, \end{aligned} \quad (38)$$

and the lower solutions $(\underline{B}, \underline{I}, \underline{V})$ are given by

$$\begin{aligned}\underline{B}(\xi) &= 0, \\ \underline{I}(\xi) &= \max \left\{ \kappa e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2} - e^{(\rho+\epsilon)\xi} \frac{1}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2}, 0 \right\}, \\ \underline{V}(\xi) &= \max \left\{ \kappa e^{\rho\xi} - e^{(\rho+\epsilon)\xi}, 0 \right\},\end{aligned}\quad (39)$$

where $0 < \kappa < 1$ will be specified later.

We denote by $\bar{\xi}_1, \bar{\xi}_2$, and $\bar{\xi}_3$ the points at which the minimum functions in the definition of the upper solutions (38) switch from their exponential form to their corresponding constant values. And similarly, we use $\underline{\xi}_2$ and $\underline{\xi}_3$ to denote the points at which the maximum functions in the definition of the lower solutions (39) switch from their exponential form to their corresponding constant values. We show the upper solutions in Fig. 3 on the right.

The transition points are given by

$$\begin{aligned}\bar{\xi}_1 &= \frac{1}{\rho} \ln \frac{(D\rho^2 - a - c\rho)(D\rho^2 - a - c\rho)}{D\rho^2 - a - 1 - c\rho}, \\ \bar{\xi}_2 &= \frac{1}{\rho} \ln (c\rho + a - D\rho^2), \\ \bar{\xi}_3 &= \frac{1}{\rho} \ln \frac{\theta}{\gamma}, \\ \underline{\xi}_2 &= \frac{1}{\epsilon} \ln \kappa \frac{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2}{c\rho + a - D\rho^2}, \\ \underline{\xi}_3 &= \frac{1}{\epsilon} \ln \kappa.\end{aligned}$$

Thus, we can rewrite the constructed upper solutions as

$$\begin{aligned}\bar{B}(\xi) &= \begin{cases} e^{\rho\xi} \frac{(c\rho + a + 1 - D\rho^2)}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)}, & \xi \leq \bar{\xi}_1 \\ 1, & \xi > \bar{\xi}_1 \end{cases} \\ \bar{I}(\xi) &= \begin{cases} e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2}, & \xi \leq \bar{\xi}_2 \\ 1, & \xi > \bar{\xi}_2 \end{cases} \\ \bar{V}(\xi) &= \begin{cases} e^{\rho\xi}, & \xi \leq \bar{\xi}_3 \\ \frac{\theta}{\gamma}, & \xi > \bar{\xi}_3, \end{cases}\end{aligned}\quad (40)$$

and the lower solutions as

$$\underline{B}(\xi) = 0,$$

$$\begin{aligned}
 \underline{I}(\xi) &= \begin{cases} \kappa e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2} - e^{(\rho+\epsilon)\xi} \frac{1}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2}, & \xi \geq \underline{\xi}_2 \\ 0, & \xi < \underline{\xi}_2 \end{cases} \\
 \underline{V}(\xi) &= \begin{cases} \kappa e^{\rho\xi} - e^{(\rho+\epsilon)\xi}, & \xi \leq \underline{\xi}_3 \\ 0, & \xi > \underline{\xi}_3. \end{cases}
 \end{aligned} \quad (41)$$

In our proofs, we make use of the following ordering

$$\bar{\xi}_1 \leq \bar{\xi}_2 \leq \bar{\xi}_3,$$

which is true due to our choice of $\bar{\rho}$ in (34).

Indeed, we rewrite the upper solutions as

$$\bar{B}(\xi) = \min \{A_1 e^{\rho\xi}, 1\}, \quad \bar{I}(\xi) = \min \{A_2 e^{\rho\xi}, 1\}, \quad \bar{V}(\xi) = \min \left\{ e^{\rho\xi}, \frac{\theta}{\gamma} \right\}.$$

We first observe that since a has a positive value, we have

$$\begin{aligned}
 A_1 &= \frac{c\rho + a + 1 - D\rho^2}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)} \geq \frac{c\rho + 1 - D\rho^2}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)} = \frac{1}{c\rho + a - D\rho^2} \\
 &= A_2.
 \end{aligned}$$

Since $A_1 \geq A_2$, it follows that

$$\bar{\xi}_1 = \frac{1}{\rho} \ln\left(\frac{1}{A_1}\right) \leq \frac{1}{\rho} \ln\left(\frac{1}{A_2}\right) = \bar{\xi}_2.$$

Next we determine the condition under which $\bar{\xi}_2 \leq \bar{\xi}_3$. This requires

$$\bar{\xi}_2 = \frac{1}{\rho} \ln\left(\frac{1}{A_2}\right) \leq \frac{1}{\rho} \ln\left(\frac{\theta}{\gamma}\right) = \bar{\xi}_3.$$

Equivalently,

$$\frac{1}{A_2} \leq \frac{\theta}{\gamma},$$

which reduces to

$$c\rho + a - D\rho^2 < \frac{\theta}{\gamma}.$$

This is precisely the assumption (31).

Lemma 2.7. For $c > \bar{c}$ and $\bar{B}$, $\bar{I}$ and $\bar{V}$ as given above (40), the following inequalities are satisfied.

$$\begin{aligned} c\bar{B}' - D\bar{B}'' &\geq -\bar{B} + \bar{I} + \bar{B}^2 - \bar{B}\bar{I} + \bar{V} - \bar{B}\bar{V}, \\ c\bar{I}' - D\bar{I}'' &\geq -a\bar{I} + (1 - \bar{B})\bar{V}, \\ c\bar{V}' - \bar{V}'' &\geq \theta\bar{I} - \gamma\bar{V}. \end{aligned} \quad (42)$$

Proof. The proof uses 5 steps.

Step 1. We first show that under the condition $c > D\rho$, the upper solution $\bar{U} = (\bar{B}, \bar{I}, \bar{V})$ satisfies the linearization inequality

$$cU' - \mathcal{D}U'' - f(U) \geq cU' - \mathcal{D}U'' - Uf'(0). \quad (43)$$

For the components $\bar{I}$ and $\bar{V}$ the inequality follows directly. For $\bar{I}$ we compute

$$\begin{aligned} c\bar{I}' - D\bar{I}'' - f(\bar{I}) &= c\bar{I}' - D\bar{I}'' + a\bar{I} + (\bar{B} - 1)\bar{V} \\ &\geq c\bar{I}' - D\bar{I}'' + a\bar{I} - \bar{V} = c\bar{I}' - D\bar{I}'' - \bar{I}f'(0), \end{aligned}$$

since $\bar{B} \geq 0$. For $\bar{V}$ we similarly obtain

$$c\bar{V}' - \bar{V}'' - f(\bar{V}) = c\bar{V}' - \bar{V}'' + \gamma\bar{V} - \theta\bar{I} = c\bar{V}' - \bar{V}'' - \bar{V}f'(0).$$

For the component $\bar{B}$, we have

$$\begin{aligned} c\bar{B}' - D\bar{B}'' - f(\bar{B}) &= c\bar{B}' - D\bar{B}'' + \bar{B} - \bar{I} - \bar{B}^2 + \bar{B}\bar{I} - \bar{V} + \bar{B}\bar{V} \\ &= c\bar{B}' - D\bar{B}'' + \bar{B}(-\bar{B} + \bar{I} + \bar{V}) + (\bar{B} - \bar{I} - \bar{V}) \\ &\geq c\bar{B}' - D\bar{B}'' + \bar{B} - \bar{I} - \bar{V} = c\bar{B}' - D\bar{B}'' - \bar{B}f'(0), \end{aligned}$$

which holds provided that

$$\bar{B} \leq \bar{I} + \bar{V}. \quad (44)$$

For $\xi < \bar{\xi}_1$, substituting the expressions from (40) into (44) gives

$$\bar{I} + \bar{V} = \frac{c\rho^2 + a + 1 - D\rho^2}{c\rho + a - D\rho^2} \geq \frac{(c\rho + a + 1 - D\rho^2)}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)} = \bar{B}.$$

Since we have

$$c\rho + a - D\rho > 0, \quad c\rho + 1 - D\rho^2 > 0,$$

the inequality reduces to the requirement

$$0 \leq c\rho - D\rho^2, \quad (45)$$

which is equivalent to the condition $c \geq D\rho$ for $\rho > 0$.

On the interval $\bar{\xi}_1 < \xi < \bar{\xi}_2$, only the condition at $\xi = \bar{\xi}_1$ needs to be verified, because $\bar{B}$ is identically equal to 1 on this interval, while both $\bar{I}$ and $\bar{V}$ are increasing functions. At $\xi = \bar{\xi}_1$ we have

$$\bar{B} = e^{\rho\bar{\xi}_1} \left(\frac{(c\rho + a + 1 - D\rho^2)}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)} \right) = 1,$$

which simplifies to

$$e^{\rho\bar{\xi}_1} \left(\frac{(c\rho + a + 1 - D\rho^2)}{(c\rho + a - D\rho^2)} \right) = c\rho - D\rho^2 + 1 \geq 1,$$

since $c > D\rho$. Thus,

$$\bar{I} + \bar{V} = e^{\rho\bar{\xi}_1} \left(\frac{(c\rho + a + 1 - D\rho^2)}{(c\rho + a - D\rho^2)} \right) \geq 1.$$

For all $\xi > \bar{\xi}_2$, the inequality $\bar{I} + \bar{V} \geq \bar{B}$ holds trivially because $\bar{I} = 1$ and $\bar{V} \geq 0$.

Since all components $\bar{B}$, $\bar{I}$, and $\bar{V}$ satisfy the linearization inequality $cU' - \mathcal{D}U'' - f(U) \geq cU' - \mathcal{D}U'' - Uf'(0)$, instead of proving the inequalities (42), we can equivalently prove

$$\begin{aligned} c\bar{B}' - D\bar{B}'' &\geq -\bar{B} + \bar{I} + \bar{V}, \\ c\bar{I}' - D\bar{I}'' &\geq -a\bar{I} + \bar{V}, \\ c\bar{V}' - \bar{V}'' &\geq \theta\bar{I} - \gamma\bar{V}. \end{aligned}$$

Thus, in more complicated cases, it is sufficient to verify the linearized inequality

$$cU' - \mathcal{D}U'' - Uf'(0) \geq 0,$$

instead of the full nonlinear inequality

$$cU' - \mathcal{D}U'' - f(U) \geq 0.$$

In the following steps, we prove the inequalities (42) for all possible positions of ξ within the ordering

$$\bar{\xi}_1 \leq \bar{\xi}_2 \leq \bar{\xi}_3.$$

Step 2. We first consider the region $\xi < \bar{\xi}_1$, where the upper solutions are given by

$$\bar{B}(\xi) = e^{\rho\xi} \frac{(c\rho + a + 1 - D\rho^2)}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)}, \quad \bar{I}(\xi) = e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2}, \quad \bar{V}(\xi) = e^{\rho\xi}.$$

Substituting these into the linearized equation

$$cU' - \mathcal{D}U'' - Uf'(0),$$

we compute each component. For $\bar{B}$, we obtain

$$c\bar{B}' - D\bar{B}'' + \bar{B} - \bar{I} - \bar{V} = e^{\rho\xi} \left[\frac{(c\rho + 1 - D\rho^2)(c\rho + a + 1 - D\rho^2)}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)} - \frac{1}{c\rho + a - D\rho^2} - 1 \right] = 0.$$

For $\bar{I}$, we have

$$c\bar{I}' - D\bar{I}'' + a\bar{I} - \bar{V} = e^{\rho\xi} \left[\frac{D\rho^2 - a - c\rho}{D\rho^2 - a - c\rho} - 1 \right] = 0.$$

For $\bar{V}$, substitution yields

$$c\bar{V}' - \bar{V}'' - \theta\bar{I} + \gamma\bar{V} = e^{\rho\xi} \left[(c\rho - \rho^2 + \gamma)(D\rho^2 - a - c\rho) + \theta \right] = 0.$$

The last equality follows from the relation

$$0 = \det(A(\rho) - c\rho\mathbb{I}) = (D\rho^2 - 1) \left[(c\rho - \rho^2 - \gamma)(D\rho^2 - a - c\rho) + \theta \right], \quad (46)$$

since $c\rho$ is the eigenvalue of $A(\rho)$.

Step 3. Next, consider $\xi_1 \leq \xi < \bar{\xi}_2 < \bar{\xi}_3$, where

$$\bar{B} = 1, \quad \bar{I} = e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2}, \quad \bar{V} = e^{\rho\xi}.$$

For $\bar{B}$, we have

$$c\bar{B}' - D\bar{B}'' + \bar{B} - \bar{B}^2 + \bar{B}\bar{I} + \bar{B}\bar{V} - \bar{I} - \bar{V} = 1 - 1 + \bar{I} + \bar{V} - \bar{I} - \bar{V} = 0.$$

For $\bar{I}$, we obtain

$$c\bar{I}' - D\bar{I}'' + a\bar{I} - \bar{V} = e^{\rho\xi} \left[(c\rho + a - D\rho^2) \frac{1}{c\rho + a - D\rho^2} - 1 \right] = 0.$$

For $\bar{V}$, we compute

$$c\bar{V}' - \bar{V}'' - \theta\bar{I} + \gamma\bar{V} = e^{\rho\xi} \left[(c\rho - 1 + \gamma) + \frac{\theta}{D\rho^2 - a - c\rho} \right] = 0.$$

Once again, the final equality results from the identity $\det(A(\rho) - c\rho\mathbb{I}) = 0$.

Step 4. In the next interval, $\bar{\xi}_1 < \bar{\xi}_2 \leq \xi < \bar{\xi}_3$, the upper solutions are

$$\bar{B} = 1, \quad \bar{I} = 1, \quad \bar{V} = e^{\rho\xi}.$$

For $\bar{B}$ we again have

$$c\bar{B}' - D\bar{B}'' + \bar{B} - \bar{B}^2 + \bar{B}\bar{I} + \bar{B}\bar{V} - \bar{I} - \bar{V} = 1 - 1 + \bar{I} + \bar{V} - \bar{I} - \bar{V} = 0.$$

For $\bar{I}$,

$$c\bar{I}' - D\bar{I}'' + a\bar{I} - (1 - \bar{B})\bar{V} = a \geq 0,$$

and this inequality holds since $a \geq 0$.

For $\bar{V}$, noting that (46) gives $(c\rho - \rho^2 - \gamma) = \frac{\theta}{(c\rho + a - D\rho^2)}$, we write

$$\begin{aligned} \bar{V}' - D\bar{V}'' - \theta\bar{I} + \gamma\bar{V} &= e^{\rho\xi} (c\rho - \rho^2 + \gamma) - \theta \\ &= e^{\rho\xi} \left(\frac{\theta}{c\rho + a - D\rho^2} \right) - \theta = \theta \left(e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2} - 1 \right) \geq 0, \end{aligned}$$

because in this region $\bar{I} = 1$ which means $1 \leq e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2}$.

Step 5. Finally, consider the last interval, $\xi \geq \bar{\xi}_3$, where

$$\bar{B} = 1, \quad \bar{I} = 1, \quad \bar{V} = \frac{\theta}{\gamma}.$$

For $\bar{B}$, we again have

$$c\bar{B}' - D\bar{B}'' + \bar{B} - \bar{B}^2 + \bar{B}\bar{I} + \bar{B}\bar{V} - \bar{I} - \bar{V} = 0.$$

For $\bar{V}$,

$$\bar{V}' - D\bar{V}'' - \theta\bar{I} + \gamma\bar{V} = \gamma \left(\frac{\theta}{\gamma} \right) - \theta \geq 0.$$

For $\bar{I}$,

$$c\bar{I}' - D\bar{I}'' + a\bar{I} - (1 - \bar{B})\bar{V} = a - 0 = a \geq 0. \quad \square$$

Now for the lower solutions, we set

$$\begin{aligned} \underline{B}(\xi) &= 0, \\ \underline{I}(\xi) &= \begin{cases} \kappa e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2} - e^{(\rho+\epsilon)\xi} \frac{1}{c_\epsilon(\rho+\epsilon) + a - D(\rho+\epsilon)^2}, & \xi \geq \underline{\xi}_2 \\ 0, & \xi < \underline{\xi}_2 \end{cases} \\ \underline{V}(\xi) &= \begin{cases} \kappa e^{\rho\xi} - e^{(\rho+\epsilon)\xi}, & \xi \leq \underline{\xi}_3 \\ 0, & \xi > \underline{\xi}_3. \end{cases} \end{aligned} \quad (47)$$

Lemma 2.8. Let $\underline{B}$, $\underline{I}$, and $\underline{V}$ be defined as in (47), and let ρ and ρ_ϵ be as in (37), with corresponding wave speeds c and c_ϵ . If

$$0 < \kappa < \min \left\{ 1, \frac{c\rho + a - D\rho^2}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \right\},$$

then the following inequalities are satisfied

$$\begin{aligned} c\underline{B}' - D\underline{B}'' &\leq (\underline{B} - 1)(\underline{B} - \underline{I}) + (1 - \underline{B})\underline{V}, \\ c\underline{I}' - \underline{I}'' &\leq -a\underline{I} + (1 - \underline{B})\underline{V}, \\ c\underline{V}' - \underline{V}'' &\leq \theta\underline{I} - \gamma\underline{V}. \end{aligned} \quad (48)$$

Proof. We first observe that, since $\underline{B} \equiv 0$, there are two possible alignments:

$$\underline{\xi}_3 < \underline{\xi}_2 \quad \text{or} \quad \underline{\xi}_2 < \underline{\xi}_3. \quad (49)$$

In what follows, we show that the inequalities in (48) hold in both cases. Also with this choice of κ , $\underline{\xi}_2$ and $\underline{\xi}_3$ are both negative.

Step 1. We begin with the first inequality in (48). Observe that in both cases of (49), since $\underline{B} = 0$ and $\underline{I} \geq 0$, $\underline{V} \geq 0$, in all the possible intervals we have

$$c\underline{B}' - D\underline{B}'' + \underline{B} - \underline{B}^2 + \underline{B}\underline{I} - \underline{I} + \underline{B}\underline{V} - \underline{V} = -\underline{I} - \underline{V} \leq 0.$$

Thus, the first inequality holds automatically, and we need only to verify the second and third inequalities in (48).

Step 2. Note that if

$$\max\{\underline{\xi}_3, \underline{\xi}_2\} \leq \xi,$$

then $\underline{B} = \underline{I} = \underline{V} = 0$, and therefore all inequalities in (48) hold trivially.

Step 3. We now consider the case $\underline{\xi}_3 < \underline{\xi}_2$. On the interval

$$\underline{\xi}_3 \leq \xi \leq \underline{\xi}_2,$$

the lower solutions take the form

$$\underline{I}(\xi) = \kappa e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2} - e^{(\rho+\epsilon)\xi} \frac{1}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2}, \quad \underline{B} = \underline{V} = 0.$$

For the second inequality in (48), we compute

$$\begin{aligned} c\underline{I}' - D\underline{I}'' + a\underline{I} - (1 - \underline{B})\underline{V} &= c\underline{I}' - D\underline{I}'' + a\underline{I} \\ &= (c\rho + a - D\rho^2)\kappa e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2} \\ &\quad - (c(\rho + \epsilon) + a - D(\rho + \epsilon)^2)e^{(\rho+\epsilon)\xi} \frac{1}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \end{aligned} \quad (50)$$

$$\begin{aligned}
&= \kappa e^{\xi \rho} - e^{(\rho+\epsilon)\xi} \left(\frac{(c(\rho+\epsilon) + a - D(\rho+\epsilon)^2)}{c_\epsilon(\rho+\epsilon) + a - D(\rho+\epsilon)^2} \right) \\
&= e^{\xi \rho} \left(\kappa - e^{\epsilon \xi} \left(\frac{(c(\rho+\epsilon) + a - D(\rho+\epsilon)^2)}{c_\epsilon(\rho+\epsilon) + a - D(\rho+\epsilon)^2} \right) \right) = e^{\xi \rho} \Lambda_1.
\end{aligned}$$

Since $c > c_\epsilon$, we observe that

$$c(\rho + \epsilon) > c_\epsilon(\rho + \epsilon) > D(\rho + \epsilon)^2 - a,$$

and therefore

$$\frac{c(\rho + \epsilon) + a - D(\rho + \epsilon)^2}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \geq 1.$$

Using that $e^{\epsilon \xi} \geq e^{\epsilon \xi_3} = \kappa$, we have

$$\begin{aligned}
\Lambda_1 &\leq \left(\kappa - e^{\epsilon \xi_3} \left(\frac{(c(\rho + \epsilon) + a - D(\rho + \epsilon)^2)}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \right) \right) \leq \left(\kappa - \kappa \left(\frac{(c(\rho + \epsilon) + a - D(\rho + \epsilon)^2)}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \right) \right) \\
&= \kappa \left(1 - \left(\frac{(c(\rho + \epsilon) + a - D(\rho + \epsilon)^2)}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \right) \right) \leq 0.
\end{aligned}$$

For the third equation, we obtain

$$c \underline{V}' - \underline{V}'' + \gamma \underline{V} - \theta \underline{I} = -\theta \underline{I} \leq 0,$$

which follows directly from the fact that $\underline{I} \geq 0$.

Step 4. We now consider the next case in (49), namely

$$\underline{\xi}_2 \leq \underline{\xi}_3.$$

On the interval

$$\underline{\xi}_2 \leq \xi \leq \underline{\xi}_3,$$

the lower solutions are given by

$$\underline{B} = \underline{I} = 0, \quad \underline{V}(\xi) = \kappa e^{\rho \xi} - e^{(\rho+\epsilon)\xi}.$$

For the second equation in (48), we obtain

$$c \underline{I}' - D \underline{I}'' + a \underline{I} - (1 - \underline{B}) \underline{V} = -\underline{V} \leq 0,$$

which holds since $\underline{V} \leq 0$.

For the third equation, we compute

$$c \underline{V}' - \underline{V}'' + \gamma \underline{V} - \theta \underline{I} = \kappa e^{\rho \xi} (c\rho - \rho^2 + \gamma) - e^{(\rho+\epsilon)\xi} (c(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma). \quad (51)$$

Using the characteristic relation for ρ ,

$$0 = \det(A(\rho) - c\rho \mathbb{I}) = (D\rho^2 - 1) \left[(c\rho - \rho^2 + \gamma)(c\rho + a - D\rho^2) - \theta \right], \quad (52)$$

and noting $c\rho + a - D\rho^2 > 0$, we obtain the identity

$$c\rho - \rho^2 + \gamma = \frac{\theta}{c\rho + a - D\rho^2} \geq 0.$$

Moreover, by the characteristic relation for $\rho + \epsilon$ and $c > c_\epsilon$ we have

$$c(\rho + \epsilon) + \gamma - (\rho + \epsilon)^2 \geq c_\epsilon(\rho + \epsilon) + \gamma - (\rho + \epsilon)^2 = \frac{\theta}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \geq 0.$$

Using $c > c_\epsilon$, rewrite the left-hand side of (51) as

$$\begin{aligned} c \underline{V}' - \underline{V}'' + \gamma \underline{V} - \theta \underline{I} &= \kappa e^{\rho\xi} (c\rho - \rho^2 + \gamma) - e^{(\rho+\epsilon)\xi} (c(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma) \\ &\leq e^{\rho\xi} \left(\kappa (c\rho - \rho^2 + \gamma) - e^{\epsilon\xi} (c_\epsilon(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma) \right) = e^{\xi\rho} \Lambda_2, \end{aligned}$$

where

$$\Lambda_2 := \kappa (c\rho - \rho^2 + \gamma) - e^{\epsilon\xi} (c_\epsilon(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma).$$

On the interval under consideration, we have the estimate

$$e^{\epsilon\xi} \geq e^{\epsilon\xi_2} = \kappa \frac{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2}{c\rho + a - D\rho^2}.$$

Hence

$$\begin{aligned} \Lambda_2 &\leq \kappa (c\rho - \rho^2 + \gamma) - \left(\kappa \frac{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2}{c\rho + a - D\rho^2} \right) (c_\epsilon(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma) \quad (53) \\ &= \kappa \left[(c\rho - \rho^2 + \gamma) - \left(\frac{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2}{c\rho + a - D\rho^2} \right) (c_\epsilon(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma) \right]. \end{aligned}$$

By the characteristic relation for $\rho + \epsilon$ with speed c_ϵ , we have

$$\begin{aligned} 0 &= \det(A(\rho + \epsilon) - c_\epsilon(\rho + \epsilon) \mathbb{I}) \\ &= (D(\rho + \epsilon)^2 - 1) \left[(c_\epsilon(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma)(c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2) - \theta \right], \end{aligned}$$

so

$$(c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2)(c_\epsilon(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma) = \theta.$$

Substituting this into the expression (53) yields

$$\Lambda_2 \leq \kappa \left[(c\rho - \rho^2 + \gamma) - \frac{\theta}{c\rho + a - D\rho^2} \right] = \kappa \left(\frac{(c\rho - \rho^2 + \gamma)(c\rho + a - D\rho^2) - \theta}{c\rho + a - D\rho^2} \right) = 0.$$

By (52), the numerator vanishes, therefore $\Lambda_2 \leq 0$, and hence

$$c \underline{V}' - \underline{V}'' + \gamma \underline{V} - \theta \underline{I} = e^{\rho\xi} \Lambda_2 \leq 0.$$

Thus, the third inequality is satisfied on this interval.

Step 5. Finally, if

$$\xi < \min\{\underline{\xi}_2, \underline{\xi}_3\},$$

the lower solutions have the form

$$\begin{aligned} \underline{B} &= 0, \quad \underline{I}(\xi) = \kappa e^{\rho\xi} \frac{1}{c\rho + a - D\rho^2} - e^{(\rho+\epsilon)\xi} \frac{1}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2}, \\ \underline{V}(\xi) &= \kappa e^{\rho\xi} - e^{(\rho+\epsilon)\xi}. \end{aligned}$$

The first inequality holds exactly as before, since $\underline{B} = 0$ and $\underline{I} \geq 0$, $\underline{V} \geq 0$.

For the second equation we compute

$$c \underline{I}' - D \underline{I}'' + a \underline{I} + (\underline{B} - 1) \underline{V} \leq c \underline{I}' - D \underline{I}'' + a \underline{I},$$

since $\underline{B} = 0$. The argument from the previous interval in (50) shows that

$$c \underline{I}' - D \underline{I}'' + a \underline{I} \leq 0,$$

and therefore the second inequality of (48) is also satisfied in this region.

For the third inequality we obtain

$$\begin{aligned} c \underline{V}' - \underline{V}'' + \gamma \underline{V} - \theta \underline{I} &= \kappa e^{\rho\xi} \left((c\rho - \rho^2 + \gamma) - \frac{\theta}{c\rho + a - D\rho^2} \right) - \\ &\quad - e^{(\rho+\epsilon)\xi} \left((c(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma) - \frac{\theta}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \right) \\ &= \kappa e^{\rho\xi} \left(\frac{(c\rho - \rho^2 + \gamma)(c\rho + a - D\rho^2) - \theta}{c\rho + a - D\rho^2} \right) \\ &\quad - e^{(\rho+\epsilon)\xi} \left(\frac{(c(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma)(c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2) - \theta}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \right) \\ &= 0 - e^{(\rho+\epsilon)\xi} \left(\frac{(c(\rho + \epsilon) - \rho^2 + \gamma)(c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2) - \theta}{c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2} \right) \\ &\leq 0, \end{aligned}$$

since we have

$$(c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2) > 0,$$

and

$$\begin{aligned} & (c(\rho + \epsilon) - \rho^2 + \gamma)(c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2) \\ & > ((c_\epsilon(\rho + \epsilon) - \rho^2 + \gamma)(c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2) = \theta, \end{aligned}$$

consequently,

$$(c(\rho + \epsilon) - (\rho + \epsilon)^2 + \gamma)(c_\epsilon(\rho + \epsilon) + a - D(\rho + \epsilon)^2) - \theta \geq 0,$$

which shows that the required inequalities hold.

Therefore, the proof is complete. $\square$

3. Proof of Theorem 1.1

Before we begin the proof of Theorem 1.1, we formally define the upper and lower solutions, as well as the function spaces mentioned earlier. We work in the space

$$\chi := \left\{ \Phi = (\phi_1, \phi_2, \phi_3) \in C(\mathbb{R}, \mathbb{R}^3) : \|\Phi\|_\mu < \infty \right\},$$

where the norm $\|\Phi\|_\mu$ is defined as the maximum of the weighted supremum norms of the three components; that is,

$$\|\Phi\|_\mu = \max \left\{ \sup_{\xi \in \mathbb{R}} |\phi_1(\xi)| e^{-\mu|\xi|}, \sup_{\xi \in \mathbb{R}} |\phi_2(\xi)| e^{-\mu|\xi|}, \sup_{\xi \in \mathbb{R}} |\phi_3(\xi)| e^{-\mu|\xi|} \right\}, \quad (54)$$

where $\mu > 0$ is a fixed weight parameter. Note that χ is a Banach space since we can define a linear isometry from $C_0(\mathbb{R}, \mathbb{R}^3)$ into $C_\mu(\mathbb{R}, \mathbb{R}^3)$ by the map

$$A : C_0(\mathbb{R}, \mathbb{R}^3) \rightarrow C_\mu(\mathbb{R}, \mathbb{R}^3), \quad A(f)(\xi) = f(\xi) e^{\mu|\xi|}.$$

We now introduce the subset of χ on which we will apply Schauder's fixed point theorem:

$$\begin{aligned} \Gamma := \{ (B, I, V) \in \chi : (\underline{B}(\xi), \underline{I}(\xi), \underline{V}(\xi)) \leq (B(\xi), I(\xi), V(\xi)) \leq (\bar{B}(\xi), \bar{I}(\xi), \bar{V}(\xi)) \\ \text{for all } \xi \in \mathbb{R} \}, \end{aligned}$$

where $(\underline{B}, \underline{I}, \underline{V})$ denotes the lower solution defined in (39), and $(\bar{B}, \bar{I}, \bar{V})$ denotes the upper solution defined in (38). Clearly,

$$(\underline{B}, \underline{I}, \underline{V}) \in \Gamma, \quad (\bar{B}, \bar{I}, \bar{V}) \in \Gamma.$$

Of course, it remains to verify that the functions constructed above do indeed form valid upper and lower solutions. This will be established in Section 3.1, where we show that Γ is a bounded,

convex, and invariant subset of χ . Now, we will use the operators defined above from (12)–(17) which are given by

$$T_1[(B, I, V)] = \frac{1}{\sqrt{c^2 + 4\alpha D}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} \underbrace{((\alpha-1)B + B^2 - BI - BV + I + V)}_{F_1(B, I, V)} d\eta \right. \\ \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} \underbrace{((\alpha-1)B + B^2 - BI - BV + I + V)}_{F_1(B, I, V)} d\eta \right),$$

where λ_1 and λ_2 solve the characteristic equation

$$D\delta^2 - c\delta - \alpha = 0,$$

and are given by

$$\delta_1 = \frac{c - \sqrt{c^2 + 4D\alpha}}{2D}, \quad \delta_2 = \frac{c + \sqrt{c^2 + 4D\alpha}}{2D}. \quad (55)$$

The second component is given by

$$T_2[(B, I, V)] = \frac{1}{\sqrt{c^2 + 4\alpha D}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} \underbrace{((\alpha-a)I + (1-B)V)}_{F_2(B, I, V)} d\eta \right. \\ \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\sigma_2} \underbrace{((\alpha-a)I + (1-B)V)}_{F_2(B, I, V)} d\eta \right),$$

where σ_1 and σ_2 solve the characteristic equation

$$D\sigma^2 - c\sigma - \alpha = 0,$$

and are given by

$$\sigma_1 = \frac{c - \sqrt{c^2 + 4D\alpha}}{2D}, \quad \sigma_2 = \frac{c + \sqrt{c^2 + 4D\alpha}}{2D}. \quad (56)$$

And the final component is given by

$$T_3[(B, I, V)] = \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} \underbrace{((\alpha-\gamma)V + \theta I)}_{F_3(B, I, V)} d\eta \right.$$

$$+ \int_{\xi}^{\infty} e^{(\xi-\eta)\zeta_2} \underbrace{((\alpha - \gamma)V + \theta I)}_{F_3(B, I, V)} d\eta \Bigg),$$

where ζ_1 and ζ_2 solve the characteristic equation

$$\zeta^2 - c\zeta - \alpha = 0,$$

and are given by

$$\zeta_1 = \frac{c - \sqrt{c^2 + 4\alpha}}{2}, \quad \zeta_2 = \frac{c + \sqrt{c^2 + 4\alpha}}{2}. \quad (57)$$

We choose

$$\alpha = \max \left\{ \gamma, a, 2 + \frac{\theta}{\gamma} \right\}, \quad (58)$$

which ensures that each component of F is nondecreasing with respect to its own variable: $F_1(B, I, V)$ is nondecreasing in B , $F_2(B, I, V)$ is nondecreasing in I , and $F_3(B, I, V)$ is non-decreasing in V . We then define the operator

$$T[(B, I, V)] = (T_1[(B, I, V)], T_2[(B, I, V)], T_3[(B, I, V)]),$$

and we will demonstrate that the fixed points of T correspond to travelling wave solutions of the system (2).

3.1. Existence of travelling wave solution for $c \geq \bar{c}$

We start with showing that the operator T is invariant in Γ in the following lemma.

Lemma 3.1. *Let $\bar{c}$ be defined as in (20), and let $c > \bar{c}$. Then the operator*

$$T : \Gamma \longrightarrow \Gamma$$

is invariant on Γ , i.e., it maps the set Γ into itself.

Proof. Let $(B, I, V) \in \Gamma$. For the upper bound of T_1 , using (43), we estimate

$$T_1[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4\alpha D}} \times \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} (\alpha B - B + I + V) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} ((\alpha B - B + I + V) d\eta \right).$$

With the choice of α (58), the above expression is non-decreasing in B , I , and V , thus, we can replace B , I and V by their upper bounds $(\bar{B}, \bar{I}, \bar{V})$

$$T_1[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \times \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} (\alpha \bar{B} - \bar{B} + \bar{I} + \bar{V}) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} (\alpha \bar{B} - \bar{B} + \bar{I} + \bar{V}) d\eta \right).$$

Step 1: For $\xi \leq \bar{\xi}_1$

$$T_1[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} (\alpha \bar{B} - \bar{B} + \bar{I} + \bar{V}) d\eta + \int_{\xi}^{\bar{\xi}_1} e^{(\xi-\eta)\delta_2} (\alpha \bar{B} - \bar{B} + \bar{I} + \bar{V}) d\eta + \int_{\bar{\xi}_1}^{\infty} e^{(\xi-\eta)\delta_2} (\alpha \bar{B} - \bar{B} + \bar{I} + \bar{V}) d\eta \right).$$

Using the fact that $\bar{B}$ satisfies the inequality (40), we obtain

$$T_1[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} (\alpha \bar{B} + c\bar{B}' - D\bar{B}'') d\eta + \int_{\xi}^{\bar{\xi}_1} e^{(\xi-\eta)\delta_2} (\alpha \bar{B} + c\bar{B}' - D\bar{B}'') d\eta + \int_{\bar{\xi}_1}^{\infty} e^{(\xi-\eta)\delta_2} (\alpha \bar{B} + c\bar{B}' - D\bar{B}'') d\eta \right).$$

Using the integration by parts on the terms $c\bar{B}'e^{\delta_1\eta}$, $c\bar{B}'e^{\delta_2\eta}$, $D\bar{B}''e^{\delta_1\eta}$, and $D\bar{B}''e^{\delta_2\eta}$, yields

$$T_1[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \times \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} \underbrace{(\alpha + \delta_1 c - \delta_1^2)}_{=0} \bar{B} d\eta + (c\bar{B} - D\bar{B}' - \delta_1 D\bar{B}) e^{\delta_1(\xi-\eta)} \Big|_{-\infty}^{\xi} \right)$$

$$\begin{aligned}
 & + \int_{\xi}^{\bar{\xi}_1} e^{(\xi-\eta)\delta_2} \underbrace{(\alpha + \delta_2 c - D\delta_2^2)}_{=0} \bar{B} d\eta + (c\bar{C} - D\bar{B}' - \delta_2 D\bar{B}) e^{\delta_2(\xi-\eta)} \Big|_{\xi}^{\bar{\xi}_1} \\
 & + \int_{\bar{\xi}_1}^{\infty} e^{(\xi-\eta)\delta_2} \underbrace{(\alpha + \delta_2 c - D\delta_2^2)}_{=0} \bar{B} d\eta + (c\bar{B} - D\bar{B}' - \delta_2 D\bar{B}) e^{\delta_2(\xi-\eta)} \Big|_{\bar{\xi}_1}^{\infty} \Bigg),
 \end{aligned}$$

where we used the fact that δ_1 and δ_2 are solutions of (55). Noting that $\delta_2 - \delta_1 = \frac{\sqrt{c^2 + 4D\alpha}}{D}$, further simplifying yields

$$\begin{aligned}
 T_1[(B, I, V)] & \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left((\delta_2 - \delta_1) D\bar{B} - D\bar{B}'(\bar{\xi}_1 -) e^{\delta_2(\xi - \bar{\xi}_1)} + D\bar{B}'(\bar{\xi}_1 +) e^{\delta_2(\xi - \bar{\xi}_1)} \right) \\
 & = \bar{B}(\xi) - \frac{e^{\delta_2(\xi - \bar{\xi}_1)}}{\sqrt{c^2 + 4D\alpha}} \left(\underbrace{\bar{B}'(\bar{\xi}_1 -)}_{>0} - \underbrace{\bar{B}'(\bar{\xi}_1 +)}_{=0} \right) \leq \bar{B}(\xi).
 \end{aligned}$$

Step 2: For $\bar{\xi}_1 < \xi$

$$\begin{aligned}
 T_1[(B, I, V)] & \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\bar{\xi}_1} e^{(\xi-\eta)\delta_1} (\alpha\bar{B} - \bar{B} + \bar{I} + \bar{V}) d\eta \right. \\
 & \quad + \int_{\bar{\xi}_1}^{\xi} e^{(\xi-\eta)\delta_1} ((\alpha\bar{B} - \bar{B} + \bar{I} + \bar{V})) d\eta \\
 & \quad \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} ((\alpha\bar{B} - \bar{B} + \bar{I} + \bar{V})) d\eta \right).
 \end{aligned}$$

Inserting the equality from (40), and applying similar calculations as the last step, yields

$$\begin{aligned}
 T_1[(B, I, V)] & \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\bar{\xi}_1} e^{(\xi-\eta)\delta_1} (\alpha\bar{B} + c\bar{B}' - D\bar{B}'') d\eta \right. \\
 & \quad + \int_{\bar{\xi}_1}^{\xi} e^{(\xi-\eta)\delta_1} ((\alpha\bar{B} + c\bar{B}' - D\bar{B}'')) d\eta \\
 & \quad \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} ((\alpha\bar{B} + c\bar{B}' - D\bar{B}'')) d\eta \right)
 \end{aligned}$$

$$= \bar{B}(\xi) - \frac{e^{\delta_2(\xi - \bar{\xi}_1)}}{\sqrt{c^2 + 4D\alpha}} \left(\underbrace{\bar{B}'(\bar{\xi}_1 -)}_{>0} - \underbrace{\bar{B}'(\bar{\xi}_1 +)}_{=0} \right) \leq \bar{B}(\xi).$$

For the upper bound of T_2 we estimate

$$T_2[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha I - aI + V) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\sigma_2} ((\alpha I - aI + V)) d\eta \right).$$

The above expression is non-decreasing in B , I , and V ; thus, we can replace B , I and V by their upper bounds

$$T_2[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha \bar{I} - a\bar{I} + \bar{V}) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\sigma_2} ((\alpha \bar{I} - a\bar{I} + \bar{V})) d\eta \right).$$

Step 3: For $\xi \leq \bar{\xi}_2$

$$\begin{aligned} T_2[(B, I, V)] &\leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha \bar{I} - a\bar{I} + \bar{V}) d\eta \right. \\ &\quad + \int_{\xi}^{\bar{\xi}_2} e^{(\xi-\eta)\sigma_2} ((\alpha \bar{I} - a\bar{I} + \bar{V})) d\eta \\ &\quad \left. + \int_{\bar{\xi}_2}^{\infty} e^{(\xi-\eta)\sigma_2} ((\alpha \bar{I} - a\bar{I} + \bar{V})) d\eta \right). \end{aligned}$$

Using that $\bar{I}$ satisfies the inequality (40) we have

$$\begin{aligned} T_2[(B, I, V)] &\leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha \bar{I} + c\bar{I}' - D\bar{I}'') d\eta \right. \\ &\quad + \int_{\xi}^{\bar{\xi}_2} e^{(\xi-\eta)\sigma_2} ((\alpha \bar{I} + c\bar{I}' - D\bar{I}'') d\eta \\ &\quad \left. + \int_{\bar{\xi}_2}^{\infty} e^{(\xi-\eta)\sigma_2} ((\alpha \bar{I} + c\bar{I}' - D\bar{I}'') d\eta \right). \end{aligned}$$

Using the integration by parts on the terms $c\bar{I}'e^{\sigma_1\eta}$, $c\bar{I}'e^{\sigma_2\eta}$, $D\bar{I}''e^{\sigma_1\eta}$, and $D\bar{I}''e^{\sigma_2\eta}$ yields

$$\begin{aligned}
T_2[(B, I, V)] &\leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \\
&\times \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} \underbrace{(\alpha + \sigma_1 c - D\sigma_1^2)}_{=0} \bar{I} d\eta + (c\bar{I} - D\bar{I}' - \lambda_1 D\bar{I}) e^{\sigma_1(\xi-\eta)} \right) \Big|_{-\infty}^{\xi} \\
&+ \int_{\xi}^{\bar{\xi}_2} e^{(\xi-\eta)\sigma_2} \underbrace{(\alpha + \sigma_2 c - D\sigma_2^2)}_{=0} \bar{I} d\eta + (c\bar{I} - D\bar{I}' - \sigma_2 D\bar{I}) e^{\sigma_2(\xi-\eta)} \Big|_{\xi}^{\bar{\xi}_2} \\
&+ \int_{\bar{\xi}_2}^{\infty} e^{(\xi-\eta)\sigma_2} \underbrace{(\alpha + \sigma_2 c - D\sigma_2^2)}_{=0} \bar{I} d\eta + (c\bar{I} - D\bar{I}' - \sigma_2 D\bar{I}) e^{\sigma_2(\xi-\eta)} \Big|_{\bar{\xi}_2}^{\infty} \Bigg),
\end{aligned}$$

where we used that σ_1 and σ_2 are solutions to (56). Noting that $\sigma_2 - \sigma_1 = \frac{\sqrt{c^2 + 4D\alpha}}{D}$ and further simplifying yields

$$\begin{aligned}
T_2[(B, I, V)] &\leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left((\sigma_2 - \sigma_1) D\bar{I} - D\bar{I}'(\bar{\xi}_2 -) e^{\sigma_2(\xi - \bar{\xi}_2)} + D\bar{I}'(\bar{\xi}_2 +) e^{\sigma_2(\xi - \bar{\xi}_2)} \right) \\
&= \bar{I}(\xi) - \frac{e^{\sigma_2(\xi - \bar{\xi}_1)}}{\sqrt{c^2 + 4D\alpha}} \left(\underbrace{\bar{I}'(\bar{\xi}_2 -)}_{>0} - \underbrace{\bar{I}'(\bar{\xi}_2 +)}_{=0} \right) \leq \bar{I}(\xi).
\end{aligned}$$

Step 4: For $\bar{\xi}_2 < \xi$

$$\begin{aligned}
T_2[(B, I, V)] &\leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\bar{\xi}_2} e^{(\xi-\eta)\sigma_1} (\alpha\bar{I} - a\bar{I} + \bar{V}) d\eta \right. \\
&\quad + \int_{\bar{\xi}_2}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha\bar{I} - a\bar{I} + \bar{V}) d\eta \\
&\quad \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\sigma_2} (\alpha\bar{I} - a\bar{I} + \bar{V}) d\eta \right).
\end{aligned}$$

Inserting the equality from (40) and applying similar calculations as the last step yields

$$T_2[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\bar{\xi}_2} e^{(\xi-\eta)\sigma_1} (\alpha\bar{I} + c\bar{I}' - D\bar{I}'') d\eta \right.$$

$$\begin{aligned}
 & + \int_{\bar{\xi}_2}^{\xi} e^{(\xi-\eta)\sigma_1} ((\alpha \bar{I} + c \bar{I}' - D \bar{I}'') d\eta \\
 & + \int_{\xi}^{\infty} e^{(\xi-\eta)\sigma_2} ((\alpha \bar{I} + c \bar{I}' - D \bar{I}'') d\eta \Bigg) \\
 & = \bar{I}(\xi) - \frac{e^{\sigma_2(\xi-\bar{\xi}_1)}}{\sqrt{I^2 + 4D\alpha}} \left(\underbrace{\bar{I}'(\bar{\xi}_2-)}_{>0} - \underbrace{\bar{I}'(\bar{\xi}_2+)}_{=0} \right) \leq \bar{I}(\xi).
 \end{aligned}$$

For the upper bound of T_3 we estimate

$$T_3[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha I - \gamma V + \theta I) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\zeta_2} ((\alpha I - \gamma V + \theta I) d\eta \right).$$

The above expression is non-decreasing in B , I , and V , thus we can replace B , I and V by their upper bounds

$$T_3[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha \bar{I} - \gamma \bar{V} + \theta \bar{I}) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\zeta_2} ((\alpha \bar{I} - \gamma \bar{V} + \theta \bar{I}) d\eta \right).$$

Step 5: For $\xi \leq \bar{\xi}_3$

$$\begin{aligned}
 T_3[(B, I, V)] & \leq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha \bar{I} - \gamma \bar{V} + \theta \bar{I}) d\eta \right. \\
 & \quad + \int_{\xi}^{\bar{\xi}_3} e^{(\xi-\eta)\zeta_2} ((\alpha \bar{I} - \gamma \bar{V} + \theta \bar{I}) d\eta \\
 & \quad \left. + \int_{\bar{\xi}_3}^{\infty} e^{(\xi-\eta)\zeta_2} ((\alpha \bar{I} - \gamma \bar{V} + \theta \bar{I}) d\eta \right).
 \end{aligned}$$

Using that $\bar{V}$ satisfies the inequality (40) we have

$$T_3[(B, I, V)] \leq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha \bar{V} + c \bar{V}' - \bar{V}'') d\eta \right.$$

$$\begin{aligned}
& + \int_{\bar{\xi}}^{\bar{\xi}_3} e^{(\xi-\eta)\zeta_2} ((\alpha \bar{V} + c \bar{V}' - \bar{V}'') d\eta \\
& + \int_{\bar{\xi}_3}^{\infty} e^{(\xi-\eta)\zeta_2} ((\alpha \bar{V} + c \bar{V}' - \bar{V}'') d\eta \Bigg).
\end{aligned}$$

Using the integration by parts on the terms $c \bar{V}' e^{\zeta_1 \eta}$, $c \bar{V}' e^{\zeta_2 \eta}$, $\bar{V}'' e^{\zeta_1 \eta}$, and $\bar{V}'' e^{\zeta_2 \eta}$ yields

$$\begin{aligned}
T_3[(B, I, V)] \leq & \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} \underbrace{(\alpha + \zeta_1 c - \zeta_1^2)}_{=0} \bar{V} d\eta + (c\bar{C} - \bar{V}' - \zeta_1 \bar{V}) e^{\zeta_1(\xi-\eta)} \Bigg|_{-\infty}^{\xi} \right. \\
& + \int_{\bar{\xi}}^{\bar{\xi}_3} e^{(\xi-\eta)\zeta_2} \underbrace{(\alpha + \zeta_2 c - \zeta_2^2)}_{=0} \bar{V} d\eta + (c\bar{V} - \bar{V}' - \zeta_2 \bar{V}) e^{\zeta_2(\xi-\eta)} \Bigg|_{\bar{\xi}}^{\bar{\xi}_3} \\
& \left. + \int_{\bar{\xi}_3}^{\infty} e^{(\xi-\eta)\zeta_2} \underbrace{(\alpha + \zeta_2 c - \zeta_2^2)}_{=0} \bar{V} d\eta + (c\bar{V} - \bar{V}' - \zeta_2 \bar{V}) e^{\zeta_2(\xi-\eta)} \Bigg|_{\bar{\xi}_3}^{\infty} \right)
\end{aligned}$$

where we used that ζ_1 and ζ_2 are solutions to (57). Nothing that $\zeta_2 - \zeta_1 = \frac{\sqrt{c^2 + 4\alpha}}{2}$, and further simplifying yields

$$\begin{aligned}
T_3[(B, I, V)] & \leq \frac{1}{\sqrt{c^2 + 4\alpha}} \left((\zeta_2 - \zeta_1) \bar{V} - \bar{V}'(\bar{\xi}_3 -) e^{\zeta_2(\xi - \bar{\xi}_3)} + \bar{V}'(\bar{\xi}_3 +) e^{\zeta_2(\xi - \bar{\xi}_3)} \right) \\
& = \bar{V}(\xi) - \frac{e^{\zeta_2(\xi - \bar{\xi}_3)}}{\sqrt{c^2 + 4\alpha}} \left(\underbrace{\bar{V}'(\bar{\xi}_3 -)}_{>0} - \underbrace{\bar{V}'(\bar{\xi}_3 +)}_{=0} \right) \leq \bar{V}(\xi).
\end{aligned}$$

Step 6: For $\bar{\xi}_3 < \xi$

$$\begin{aligned}
T_3[(B, I, V)] \leq & \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\bar{\xi}_3} e^{(\xi-\eta)\zeta_1} (\alpha \bar{V} + \theta \bar{I} - \gamma \bar{V}) d\eta \right. \\
& \left. + \int_{\bar{\xi}_3}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha \bar{V} + \theta \bar{I} - \gamma \bar{V}) d\eta \right)
\end{aligned}$$

$$+ \int_{\xi}^{\infty} e^{(\xi-\eta)\zeta_2} ((\alpha \bar{V} + \theta \bar{I} - \gamma \bar{V}) d\eta) \Bigg).$$

Inserting the equality from (40) and applying similar calculations as the last step, yields

$$\begin{aligned} T_3[(B, I, V)] &\leq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\bar{\xi}_3} e^{(\xi-\eta)\zeta_1} (\alpha \bar{V} + c \bar{V}' - \bar{V}'') d\eta \right. \\ &\quad + \int_{\bar{\xi}_3}^{\xi} e^{(\xi-\eta)\zeta_1} ((\alpha \bar{V} + c \bar{V}' \bar{V}'') d\eta \\ &\quad \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\zeta_2} ((\alpha \bar{V} + c \bar{V}' - \bar{V}'') d\eta) \right) \\ &= \bar{V}(\xi) - \frac{e^{\zeta_2(\xi-\bar{\xi}_3)}}{\sqrt{V^2 + 4\alpha}} \left(\underbrace{\bar{V}'(\bar{\xi}_3-)}_{>0} - \underbrace{\bar{V}'(\bar{\xi}_3+)}_{=0} \right) \leq \bar{V}(\xi). \end{aligned}$$

Step 7: For the lower bound of T_1 we again note that the condition (58) makes T_1 non-decreasing in B , thus

$$\begin{aligned} T_1[(B, I, V)] &= \frac{1}{\sqrt{c^2 + 4\alpha D}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} (\alpha B - B + B^2 - BI - BV + I + V) d\eta \right. \\ &\quad \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} ((\alpha B - B + B^2 - BI - BV + I + V) d\eta) \right) \\ &\geq \frac{1}{\sqrt{c^2 + 4\alpha D}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} (\alpha \underline{B} - \underline{B} + \underline{B}^2 - \underline{B}\underline{I} - \underline{B}\underline{V} + \underline{I} + \underline{V}) d\eta \right. \\ &\quad \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} ((\alpha \underline{B} - \underline{B} + \underline{B}^2 - \underline{B}\underline{I} - \underline{B}\underline{V} + \underline{I} + \underline{V}) d\eta) \right). \end{aligned}$$

Inserting the inequality (41) yields

$$T_1[(B, I, V)] \geq \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\delta_1} (\alpha \underline{B} + c \underline{B}' - D \underline{B}'') d\eta \right.$$

$$\begin{aligned}
& + \int_{\xi}^{\infty} e^{(\xi-\eta)\delta_2} ((\alpha \underline{B} + c \underline{B}' - D \underline{B}'') d\eta \\
& = 0 \geq \underline{B}(\xi).
\end{aligned}$$

Step 8: For the lower bound of T_2 , let $\xi < \underline{\xi}_2$, using the inequality (48) yields

$$\begin{aligned}
T_2[(B, I, V)] &= \frac{1}{\sqrt{c^2 + 4\alpha D}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha I - aI + (1-B)V) d\eta \right. \\
&\quad \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\sigma_2} ((\alpha I - aI + (1-B)V) d\eta \right) \\
&\geq \frac{1}{\sqrt{c^2 + D4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha \underline{I} - a\underline{I} + (1-\underline{B})\underline{V}) d\eta \right. \\
&\quad \left. + \int_{\xi}^{\underline{\xi}_2} e^{(\xi-\eta)\sigma_2} ((\alpha \underline{I} - a\underline{I} + (1-\underline{B})\underline{V}) d\eta \right) \\
&\geq \frac{1}{\sqrt{c^2 + D4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha \underline{I} + c \underline{I}' - D \underline{I}'') d\eta \right. \\
&\quad \left. + \int_{\xi}^{\underline{\xi}_2} e^{(\xi-\eta)\sigma_2} ((\alpha \underline{I} + c \underline{I}' - D \underline{I}'') d\eta \right) \\
&= \underline{I} + \frac{e^{\sigma_2(\xi-\underline{\xi}_2)}}{\sqrt{c^2 + D4\alpha}} \left(\underbrace{c \underline{I}(\underline{\xi}_2)}_{=0} - \underbrace{D \underline{I}'(\underline{\xi}_2^-)}_{<0} - \underbrace{\sigma_2 D \underline{I}(\underline{\xi}_2)}_{=0} \right) \geq \underline{I}.
\end{aligned}$$

Step 9: Now let $\xi \geq \underline{\xi}_2$

$$\begin{aligned}
T_2[(B, I, V)] &\geq \frac{1}{\sqrt{c^2 + D4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\sigma_1} (\alpha \underline{I} - a\underline{I} + (1-\underline{B})\underline{V}) d\eta \right. \\
&\quad \left. + \int_{\xi}^{\infty} e^{(\xi-\eta)\sigma_2} ((\alpha \underline{I} - a\underline{I} + (1-\underline{B})\underline{V}) d\eta \right)
\end{aligned}$$

$$\begin{aligned} &\geq \frac{1}{\sqrt{c^2 + D4\alpha}} \left(\int_{-\infty}^{\xi_1} e^{(\xi-\eta)\sigma_1} (\alpha \underline{I} - a \underline{I} + (1 - \underline{B}) \underline{V}) d\eta \right) \\ &= \frac{1}{\sqrt{c^2 + D4\alpha}} \left(\underbrace{c \underline{I}(\xi_2)}_{=0} - \underbrace{D \underline{I}'(\xi_2)}_{<0} - \sigma_1 D \underbrace{\underline{I}(\xi_2)}_{=0} \right) \geq 0 = \underline{I}(\xi). \end{aligned}$$

Step 10: For the lower bound of T_3 , let $\xi \leq \underline{\xi}_3$, we can use the relation in (41) and obtain

$$\begin{aligned} &T_3[(B, I, V)] \\ &= \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha V + \theta I - \gamma V) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\zeta_2} ((\alpha V + \theta I - \gamma V) d\eta \right) \\ &\geq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha \underline{V} + \theta \underline{I} - \gamma \underline{V}) d\eta + \int_{\xi}^{\underline{\xi}_3} e^{(\xi-\eta)\zeta_2} ((\alpha \underline{V} + \theta \underline{I} - \gamma \underline{V}) d\eta \right) \\ &\geq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha \underline{V} + c \underline{V}' - \underline{V}'') d\eta + \int_{\xi}^{\underline{\xi}_3} e^{(\xi-\eta)\zeta_2} ((\alpha \underline{V} + c \underline{V}' - \underline{V}'') d\eta \right) \\ &= \underline{V} + \frac{e^{\zeta_2(\xi-\underline{\xi}_3)}}{\sqrt{c^2 + 4\alpha}} \left(\underbrace{c \underline{V}(\underline{\xi}_3)}_{=0} - \underbrace{\underline{V}'(\underline{\xi}_3-)}_{<0} - \zeta_2 \underbrace{\underline{V}(\underline{\xi}_3)}_{=0} \right) \geq \underline{V}(\xi). \end{aligned}$$

Step 11: Now let $\xi > \underline{\xi}_3$

$$\begin{aligned} &T_3[(B, I, V)] \\ &\geq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{(\xi-\eta)\zeta_1} (\alpha \underline{V} + \theta \underline{I} - \gamma \underline{V}) d\eta + \int_{\xi}^{\infty} e^{(\xi-\eta)\zeta_2} ((\alpha \underline{V} + \theta \underline{I} - \gamma \underline{V}) d\eta \right) \\ &\geq \frac{1}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\underline{\xi}_3} e^{(\xi-\eta)\zeta_1} (\alpha \underline{V} + c \underline{V}' - \underline{V}'') d\eta \right) \\ &\geq \frac{e^{\zeta_2(\xi-\underline{\xi}_3)}}{\sqrt{c^2 + 4\alpha}} \left(\underbrace{c \underline{V}(\underline{\xi}_3)}_{=0} - \underbrace{\underline{V}'(\underline{\xi}_3-)}_{<0} - \zeta_2 \underbrace{\underline{V}(\underline{\xi}_3)}_{=0} \right) \geq 0 = \underline{V}. \end{aligned}$$

Thus $\forall (B, I, V) \in \Gamma$, $T[(B, I, V)] \in \Gamma$. $\square$

Next, we show that T is continuous on Γ with respect to the norm $\|\cdot\|_\mu$ defined in (54).

Lemma 3.2. *Suppose $c > \bar{c}$. Then for sufficiently small μ , the operator $T : \Gamma \rightarrow \Gamma$ is continuous with respect to the weighted norm $\|\cdot\|_\mu$.*

Proof. Assume $\Phi = (\phi_1, \phi_2, \phi_3)$, $\Psi = (\psi_1, \psi_2, \psi_3) \in \Gamma$ and $\Phi \neq \Psi$. Consider

$$\begin{aligned} |F_1(\Phi) - F_1(\Psi)| &= |\alpha(\phi_1 - \psi_1) - (\phi_1 - \psi_1) + (\phi_1^2 - \psi_1^2) - \phi_1\phi_2 + \psi_1\psi_2 - \phi_1\phi_3 + \psi_1\psi_3| \\ &= |(\phi_1 - \psi_1)(\alpha - 1 + (\phi_1 + \psi_1)) - \phi_1\phi_2 + \phi_1\psi_2 - \phi_1\psi_2 + \psi_1\psi_2 - \phi_1\phi_3 + \phi_1\psi_3 - \phi_1\psi_3 + \psi_1\psi_3| \\ &= |(\phi_1 - \psi_1)(\alpha - 1 + (\phi_1 + \psi_1) - \psi_2 - \psi_3) + (\phi_2 - \psi_2)(-\phi_1) + (\phi_3 - \psi_3)(-\phi_1)| \\ &\leq |\alpha - 1 + (\phi_1 + \psi_1) - \psi_2 - \psi_3||\phi_1 - \psi_1| + |\phi_1||\phi_2 - \psi_2| + |\phi_1||\phi_3 - \psi_3|. \end{aligned}$$

Since Φ and Ψ belong to Γ , all of their components are bounded. Consequently, there exists a constant $\widehat{C}_1 > 0$ such that $|F_1(\Phi) - F_1(\Psi)|e^{-\mu|\xi|} \leq \widehat{C}_1\|\Phi - \Psi\|_\mu$. We have

$$\begin{aligned} |T_1[\Phi] - T_1[\Psi]|e^{-\mu|\xi|} &= \frac{e^{-\mu|\xi|}}{\sqrt{c^2 + 4D\alpha}} \left| \int_{-\infty}^{\xi} e^{\delta_1(\xi-\eta)} (F_1(\Phi) - F_1(\Psi)) d\eta \right. \\ &\quad \left. + \int_{\xi}^{\infty} e^{\delta_2(\xi-\eta)} (F_1(\Phi) - F_1(\Psi)) d\eta \right| \\ &\leq \frac{e^{-\mu|\xi|}}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{\delta_1(\xi-\eta)+\mu|\eta|} |F_1(\Phi) - F_1(\Psi)| e^{-\mu|\eta|} d\eta \right. \\ &\quad \left. + \int_{\xi}^{\infty} e^{\delta_2(\xi-\eta)+\mu|\eta|} |F_1(\Phi) - F_1(\Psi)| e^{-\mu|\eta|} d\eta \right) \\ &\leq \frac{\widehat{C}_1 e^{-\mu|\xi|}}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{\delta_1(\xi-\eta)+\mu|\eta|} d\eta + \int_{\xi}^{\infty} e^{\delta_2(\xi-\eta)+\mu|\eta|} d\eta \right) \|\Phi - \Psi\|_\mu. \end{aligned}$$

The integrals are bounded for $0 < \mu < \min\{-\delta_1, \delta_2\}$, which we assume henceforth. We estimate the integrals separately over the regions $\xi < 0$ and $\xi \geq 0$. If $\xi < 0$, then

$$\begin{aligned} |T_1[\Phi] - T_1[\Psi]|e^{-\mu|\xi|} &\leq \frac{\widehat{C}_1 e^{\mu\xi}}{\sqrt{c^2 + 4D\alpha}} \left(e^{\delta_1\xi} \int_{-\infty}^{\xi} e^{-(\delta_1+\mu)\eta} d\eta + e^{\delta_2\xi} \int_{\xi}^0 e^{-(\delta_2+\mu)\eta} d\eta + e^{\delta_2\xi} \int_0^{\infty} e^{(\mu-\delta_2)\eta} d\eta \right) \|\Phi - \Psi\|_\mu \\ &= \frac{\widehat{C}_1}{\sqrt{c^2 + 4D\alpha}} \left(-\frac{1}{\mu + \delta_1} + \frac{1 - e^{(\delta_2+\mu)\xi}}{\mu + \delta_2} + \frac{e^{(\delta_2+\mu)\xi}}{\delta_2 - \mu} \right) \|\Phi - \Psi\|_\mu \end{aligned}$$

$$\leq \frac{\widehat{C}_1}{\sqrt{c^2 + 4D\alpha}} \left(\underbrace{-\frac{1}{\mu + \delta_1} + \frac{1}{\mu + \delta_2} + \frac{1}{\delta_2 - \mu}}_{\tilde{C}_{\xi < 0}} \right) \|\Phi - \Psi\|_{\mu}.$$

Now, if $\xi \geq 0$, we have

$$\begin{aligned} & |T_1[\Phi] - T_1[\Psi]| e^{-\mu|\xi|} \\ & \leq \frac{\widehat{C}_1 e^{-\mu\xi}}{\sqrt{c^2 + 4D\alpha}} \left(e^{\delta_1\xi} \int_{-\infty}^0 e^{-(\delta_1 + \mu)\eta} d\eta + e^{\delta_1\xi} \int_0^{\xi} e^{(\mu - \delta_1) + \eta} d\eta + e^{\delta_2\xi} \int_{\xi}^{\infty} e^{(\mu - \delta_2)\eta} d\eta \right) \|\Phi - \Psi\|_{\mu} \\ & = \frac{\widehat{C}_1}{\sqrt{c^2 + 4D\alpha}} \left(-\frac{e^{(\delta_1 - \mu)\xi}}{\mu + \delta_1} + \frac{1 - e^{(\delta_1 - \mu)\xi}}{\mu - \delta_1} + \frac{1}{\delta_2 - \mu} \right) \|\Phi - \Psi\|_{\mu} \\ & \leq \frac{\widehat{C}_1}{\sqrt{c^2 + 4D\alpha}} \left(\underbrace{-\frac{1}{\mu + \delta_1} + \frac{1}{\mu - \delta_1} + \frac{1}{\delta_2 - \mu}}_{\tilde{C}_{\xi > 0}} \right) \|\Phi - \Psi\|_{\mu}. \end{aligned}$$

Let $C_1 = \frac{\widehat{C}_1}{\sqrt{c^2 + 4D\alpha}} \max\{\tilde{C}_{\xi > 0}, \tilde{C}_{\xi < 0}\}$, thus $|T_1[\Phi] - T_1[\Psi]| e^{-\mu|\xi|} \leq C_1 \|\Phi - \Psi\|_{\mu}$.

We now establish an analogous estimate for the second component T_2 . To begin, we examine the integrand

$$\begin{aligned} |F_2(\Phi) - F_2(\Psi)| &= |\alpha(\phi_1 - \psi_2) - (\phi_1\phi_3 - \psi_1\psi_3) - a(\phi_2 - \psi_2) + (\phi_3 - \psi_3)| \\ &= |(\alpha - a)(\phi_2 - \psi_2) - \phi_2\psi_3 + \phi_1\psi_3 - \phi_1\psi_3 + \psi_1\psi_3 + (\phi_3 - \psi_3)| \\ &= |(\phi_1 - \psi_1)(-\psi_3) + (\phi_2 - \psi_2)(\alpha - a) + (\phi_3 - \psi_3)(1 - \phi_1)| \\ &\leq |\phi_1 - \psi_1||\psi_3| + |\phi_2 - \psi_2||\alpha - a| + |\phi_3 - \psi_3||1 - \phi_1|. \end{aligned}$$

Thus, there exists a constant $\widehat{C}_2$ such that

$$|F_2(\Phi) - F_2(\Psi)| e^{-\mu|\xi|} \leq \widehat{C}_2 \|\Phi - \Psi\|_{\mu}.$$

Turning now to the operator T_2 , we obtain

$$\begin{aligned} |T_2[\Phi] - T_2[\Psi]| e^{-\mu|\xi|} &= \frac{e^{-\mu|\xi|}}{\sqrt{c^2 + 4D\alpha}} \left| \int_{-\infty}^{\xi} e^{\sigma_1(\xi - \eta)} (F_2(\Phi) - F_2(\Psi)) d\eta \right. \\ &\quad \left. + \int_{\xi}^{\infty} e^{\sigma_2(\xi - \eta)} (F_2(\Phi) - F_2(\Psi)) d\eta \right| \end{aligned}$$

$$\begin{aligned}
&\leq \frac{e^{-\mu|\xi|}}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{\sigma_1(\xi-\eta)+\mu|\eta|} |F_2(\Phi) - F_2(\Psi)| e^{-\mu|\eta|} d\eta \right. \\
&\quad \left. + \int_{\xi}^{\infty} e^{\sigma_2(\xi-\eta)+\mu|\eta|} |F_2(\Phi) - F_2(\Psi)| e^{-\mu|\eta|} d\eta \right) \\
&\leq \frac{\widehat{C}_2 e^{-\mu|\xi|}}{\sqrt{c^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{\sigma_1(\xi-\eta)+\mu|\eta|} d\eta + \int_{\xi}^{\infty} e^{\sigma_2(\xi-\eta)+\mu|\eta|} d\eta \right) \|\Phi - \Psi\|_{\mu}.
\end{aligned}$$

The integrals are bounded for

$$0 < \mu < \min\{-\sigma_1, \sigma_2\},$$

which we assume from now on. We estimate the integrals separately on the regions $\xi < 0$ and $\xi \geq 0$. If $\xi < 0$, then we have

$$\begin{aligned}
&|T_2[\Phi] - T_2[\Psi]| e^{-\mu|\xi|} \\
&\leq \frac{\widehat{C}_2 e^{\mu\xi}}{\sqrt{c^2 + 4D\alpha}} \left(e^{\sigma_1\xi} \int_{-\infty}^{\xi} e^{-(\sigma_1+\mu)\eta} d\eta + e^{\sigma_2\xi} \int_{\xi}^0 e^{-(\sigma_2+\mu)\eta} d\eta + e^{\sigma_2\xi} \int_0^{\infty} e^{(\mu-\sigma_2)\eta} d\eta \right) \|\Phi - \Psi\|_{\mu} \\
&= \frac{\widehat{C}_2}{\sqrt{c^2 + 4D\alpha}} \left(-\frac{1}{\mu + \sigma_1} + \frac{1 - e^{(\mu_2+\mu)\xi}}{\mu + \sigma_2} + \frac{e^{(\sigma_2+\mu)\xi}}{\sigma_2 - \mu} \right) \|\Phi - \Psi\|_{\mu} \\
&\leq \frac{\widehat{C}_2}{\sqrt{c^2 + 4D\alpha}} \left(\underbrace{-\frac{1}{\mu + \sigma_1} + \frac{1}{\mu + \sigma_2} + \frac{1}{\sigma_2 - \mu}}_{\widehat{C}_{\xi < 0}} \right) \|\Phi - \Psi\|_{\mu}.
\end{aligned}$$

Now, for the other case $\xi \geq 0$, it gives

$$\begin{aligned}
&|T_2[\Phi] - T_2[\Psi]| e^{-\mu|\xi|} \\
&\leq \frac{\widehat{C}_2 e^{-\mu\xi}}{\sqrt{c^2 + 4D\alpha}} \left(e^{\sigma_1\xi} \int_{-\infty}^0 e^{-(\sigma_1+\mu)\eta} d\eta + e^{\sigma_1\xi} \int_0^{\xi} e^{(\mu-\sigma_1)+\eta} d\eta + e^{\sigma_2\xi} \int_{\xi}^{\infty} e^{(\mu-\sigma_2)\eta} d\eta \right) \|\Phi - \Psi\|_{\mu} \\
&= \frac{\widehat{C}_2}{\sqrt{c^2 + 4D\alpha}} \left(-\frac{e^{(\sigma_1-\mu)\xi}}{\mu + \sigma_1} + \frac{1 - e^{(\sigma_1-\mu)\xi}}{\mu - \sigma_1} + \frac{1}{\sigma_2 - \mu} \right) \|\Phi - \Psi\|_{\mu}
\end{aligned}$$

$$\leq \frac{\widehat{C}_2}{\sqrt{c^2 + 4D\alpha}} \left(\underbrace{-\frac{1}{\mu + \sigma_1} + \frac{1}{\mu - \sigma_1} + \frac{1}{\sigma_2 - \mu}}_{\tilde{C}_{\xi > 0}} \right) \|\Phi - \Psi\|_\mu.$$

Let $C_2 = \frac{\widehat{C}_2}{\sqrt{c^2 + 4D\alpha}} \max\{\tilde{C}_{\xi > 0}, \tilde{C}_{\xi < 0}\}$, thus $|T_2[\Phi] - T_2[\Psi]|e^{-\mu|\xi|} \leq C_2 \|\Phi - \Psi\|_\mu$.

Again, we derive a corresponding estimate for the third component T_3 . We begin by examining the integrand

$$\begin{aligned} |F_3(\Phi) - F_3(\Psi)| &= |(\alpha + \theta)(\phi_2 - \psi_2) - \gamma(\phi_3 - \psi_3)| \\ &\leq |\phi_1 - \psi_1||0| + |\phi_2 - \psi_2||\theta| + |\phi_3 - \psi_3||\gamma| \end{aligned}$$

Thus, there exists a constant $\widehat{C}_3$ such that $|F_3(\Phi) - F_3(\Psi)|e^{-\mu|\xi|} \leq \widehat{C}_3 \|\Phi - \Psi\|_\mu$. Moving to the operator T_3 itself, we write

$$\begin{aligned} |T_3[\Phi] - T_3[\Psi]|e^{-\mu|\xi|} &= \frac{e^{-\mu|\xi|}}{\sqrt{c^2 + 4\alpha}} \left| \int_{-\infty}^{\xi} e^{\zeta_1(\xi-\eta)} (F_3(\Phi) - F_3(\Psi)) d\eta \right. \\ &\quad \left. + \int_{\xi}^{\infty} e^{\zeta_2(\xi-\eta)} (F_3(\Phi) - F_3(\Psi)) d\eta \right| \\ &= \frac{e^{-\mu|\xi|}}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{\zeta_1(\xi-\eta)+\mu|\eta|} |F_3(\Phi) - F_3(\Psi)| e^{-\mu|\eta|} d\eta + \right. \\ &\quad \left. \int_{\xi}^{\infty} e^{\zeta_2(\xi-\eta)+\mu|\eta|} |F_3(\Phi) - F_3(\Psi)| e^{-\mu|\eta|} d\eta \right) \\ &\leq \frac{\widehat{C}_3 e^{-\mu|\xi|}}{\sqrt{c^2 + 4\alpha}} \left(\int_{-\infty}^{\xi} e^{\zeta_1(\xi-\eta)+\mu|\eta|} d\eta + \int_{\xi}^{\infty} e^{\zeta_2(\xi-\eta)+\mu|\eta|} d\eta \right) \|\Phi - \Psi\|_\mu. \end{aligned}$$

The integrals are bounded for $0 < \mu < \min\{-\zeta_1, \zeta_2\}$ which we assume forthwith. We estimate the integrals separately for $\xi < 0$ and $\xi \geq 0$. If $\xi < 0$, it follows

$$\begin{aligned} |T_3[\Phi] - T_3[\Psi]|e^{-\mu|\xi|} &\leq \frac{\widehat{C}_3 e^{\mu\xi}}{\sqrt{c^2 + 4\alpha}} \left(e^{\zeta_1\xi} \int_{-\infty}^{\xi} e^{-(\zeta_1+\mu)\eta} d\eta + e^{\zeta_2\xi} \int_{\xi}^0 e^{-(\zeta_2+\mu)\eta} d\eta + e^{\zeta_2\xi} \int_0^{\infty} e^{(\mu-\zeta_2)\eta} d\eta \right) \|\Phi - \Psi\|_\mu \end{aligned}$$

$$\begin{aligned}
&= \frac{\hat{C}_3}{\sqrt{c^2 + 4\alpha}} \left(-\frac{1}{\mu + \zeta_1} + \frac{1 - e^{(\zeta_2 + \mu)\xi}}{\mu + \zeta_2} + \frac{e^{(\zeta_2 + \mu)\xi}}{\zeta_2 - \mu} \right) \|\Phi - \Psi\|_\mu \\
&\leq \frac{\hat{C}_3}{\sqrt{c^2 + 4\alpha}} \left(\underbrace{-\frac{1}{\mu + \zeta_1} + \frac{1}{\mu + \zeta_2} + \frac{1}{\zeta_2 - \mu}}_{\tilde{C}_{\xi < 0}} \right) \|\Phi - \Psi\|_\mu.
\end{aligned}$$

Now, for the other case $\xi \geq 0$, we obtain

$$\begin{aligned}
&|T_3[\Phi] - T_3[\Psi]| |e^{-\mu|\xi|}| \\
&\leq \frac{\hat{C}_3 e^{-\mu\xi}}{\sqrt{c^2 + 4\alpha}} \left(e^{\zeta_1\xi} \int_{-\infty}^0 e^{-(\zeta_1 + \mu)\eta} d\eta + e^{\zeta_1\xi} \int_0^\xi e^{(\mu - \zeta_1) + \eta} d\eta + e^{\zeta_2\xi} \int_\xi^\infty e^{(\mu - \zeta_2)\eta} d\eta \right) \|\Phi - \Psi\|_\mu \\
&= \frac{\hat{C}_3}{\sqrt{c^2 + 4\alpha}} \left(-\frac{e^{(\zeta_1 - \mu)\xi}}{\mu + \zeta_1} + \frac{1 - e^{(\zeta_1 - \mu)\xi}}{\mu - \zeta_1} + \frac{1}{\zeta_2 - \mu} \right) \|\Phi - \Psi\|_\mu \\
&\leq \frac{\hat{C}_3}{\sqrt{c^2 + 4\alpha}} \left(\underbrace{-\frac{1}{\mu + \zeta_1} + \frac{1}{\mu - \zeta_1} + \frac{1}{\zeta_2 - \mu}}_{\tilde{C}_{\xi > 0}} \right) \|\Phi - \Psi\|_\mu
\end{aligned}$$

Let $C_3 = \frac{\hat{C}_3}{\sqrt{c^2 + 4\alpha}} \max\{\tilde{C}_{\xi > 0}, \tilde{C}_{\xi < 0}\}$, thus $|T_3[\Phi] - T_3[\Psi]| |e^{-\mu|\xi|}| \leq C_3 \|\Phi - \Psi\|_\mu$.

Therefore, there exists a constant $C^* > 0$ such that

$$\|T[\Phi] - T[\Psi]\|_\mu \leq C^* \|\Phi - \Psi\|_\mu,$$

and hence T is continuous with respect to the norm $\|\cdot\|_\mu$. $\square$

Lemma 3.3. *If $c > \bar{c}$, then $T : \Gamma \rightarrow \Gamma$ is compact with respect to the norm $\|\cdot\|_\mu$.*

Proof. Let $\Phi \in \Gamma$, then consider

$$\begin{aligned}
|F_1(\Phi)| &= |(\alpha - 1)\phi_1 + \phi_1^2 - \phi_1\phi_2 - \phi_1\phi_3 + \phi_2 + \phi_3| \\
&= |\phi_1(\alpha - 1 + \phi_1) + (1 - \phi_1)\phi_2 + (1 - \phi_1)\phi_3| \\
&\leq |\alpha + 1 - \phi_1||\phi_1| + |\phi_2||1 - \phi_1| + |\phi_3||1 - \phi_1| \\
&\leq |\alpha + 1 - c_1||\phi_1| + |\phi_2||c_1| + |\phi_3||c_1| \leq M_1,
\end{aligned}$$

since ϕ_1, ϕ_2 and ϕ_3 are bounded in Γ . Now consider

$$\begin{aligned}
|T_1[\Phi]| &= \frac{1}{\sqrt{c^2 + 4D\alpha}} \left| \int_{-\infty}^{\xi} e^{\delta_1(\xi-\eta)} F_1[\Phi] d\eta + \int_{\xi}^{\infty} e^{\delta_2(\xi-\eta)} F_1[\Phi] d\eta \right| \\
&\leq \frac{M_1}{\sqrt{c^2 + 4D\alpha}} \left| \int_{-\infty}^{\xi} e^{\delta_1(\xi-\eta)} d\eta + \int_{\xi}^{\infty} e^{\delta_2(\xi-\eta)} d\eta \right| \\
&= \frac{M_1}{\sqrt{c^2 + 4D\alpha}} \left| \frac{-1}{\delta_1} + \frac{1}{\delta_2} \right| = \frac{M_1}{\sqrt{c^2 + 4D\alpha}} \left| \frac{\sqrt{c^2 + 4D\alpha}}{\alpha} \right| = \frac{M_1}{\alpha},
\end{aligned}$$

where we used the definition of δ_1 and δ_2 from (55).

Now for the other compartment

$$|F_2(\Phi)| = |(\alpha - a)\phi_2 + (1 - \phi_1)\phi_3| \leq |\alpha - a||\phi_2| + |1 - \phi_1||\phi_3| \leq M_2,$$

since ϕ_1 , ϕ_2 and ϕ_3 are bounded in Γ . Now consider

$$\begin{aligned}
|T_2[\Phi]| &= \frac{1}{\sqrt{c^2 + 4D\alpha}} \left| \int_{-\infty}^{\xi} e^{\sigma_1(\xi-\eta)} F_2[\Phi] d\eta + \int_{\xi}^{\infty} e^{\sigma_2(\xi-\eta)} F_2[\Phi] d\eta \right| \\
&\leq \frac{M_2}{\sqrt{c^2 + 4D\alpha}} \left| \int_{-\infty}^{\xi} e^{\sigma_1(\xi-\eta)} d\eta + \int_{\xi}^{\infty} e^{\sigma_2(\xi-\eta)} d\eta \right| = \frac{M_2}{\alpha}.
\end{aligned}$$

Similarly, noting that ϕ_2 and ϕ_3 are bounded, for the third component we have

$$|F_3(\Phi)| = |(\alpha - \gamma)\phi_3 + \theta\phi_2| \leq M_3.$$

Thus,

$$\begin{aligned}
|T_3[\Phi]| &= \frac{1}{\sqrt{c^2 + 4\alpha}} \left| \int_{-\infty}^{\xi} e^{\zeta_1(\xi-\eta)} F_3[\Phi] d\eta + \int_{\xi}^{\infty} e^{\zeta_2(\xi-\eta)} F_3[\Phi] d\eta \right| \\
&\leq \frac{M_3}{\sqrt{c^2 + 4\alpha}} \left| \int_{-\infty}^{\xi} e^{\zeta_1(\xi-\eta)} d\eta + \int_{\xi}^{\infty} e^{\zeta_2(\xi-\eta)} d\eta \right| = \frac{M_3}{\alpha}.
\end{aligned}$$

Now consider

$$\left| \frac{d}{d\xi} T_1[\Phi] \right| = \left| \frac{1}{\sqrt{c^2 + 4D\alpha}} \left(\delta_1 \int_{-\infty}^{\xi} e^{\delta_1(\xi-\eta)} F_1[\Phi] d\eta + \delta_2 \int_{\xi}^{\infty} e^{\delta_2(\xi-\eta)} F_1[\Phi] d\eta \right) \right|$$

$$\begin{aligned} &\leq \frac{M_1}{\sqrt{c^2 + 4D\alpha}} \left(|\delta_1| \int_{-\infty}^{\xi} e^{\delta_1(\xi-\eta)} d\eta + \delta_2 \int_{\xi}^{\infty} e^{\delta_2(\xi-\eta)} d\eta \right) \\ &\leq \frac{M_1}{\sqrt{c^2 + 4D\alpha}} \left(\frac{|\delta_1|}{-\delta_1} + 1 \right) = \frac{2M_1}{\sqrt{c^2 + 4D\alpha}}. \end{aligned}$$

Thus, $\left| \frac{d}{d\xi} T_1[\Phi] \right| \leq \frac{2M_1}{\sqrt{c^2 + 4D\alpha}}$. Similarly, we can find that $\left| \frac{d}{d\xi} T_2[\Phi] \right| \leq \frac{2M_2}{\sqrt{c^2 + 4D\alpha}}$, and $\left| \frac{d}{d\xi} T_3[\Phi] \right| \leq \frac{2M_3}{\sqrt{c^2 + 4\alpha}}$, hence

$$\left\| \frac{d}{d\xi} T[\phi] \right\| \leq 2 \max \left\{ \frac{2M_1}{\sqrt{c^2 + 4D\alpha}}, \frac{2M_2}{\sqrt{c^2 + 4D\alpha}}, \frac{2M_3}{\sqrt{c^2 + 4\alpha}} \right\}.$$

To obtain a convergent subsequence, we introduce a continuous cut-off function as follows. We define the operator

$$T^n[\Phi](\xi) = \begin{cases} T[\Phi](\xi), & \xi \in [-n, n] \\ T[\Phi](-n), & \xi \in [-\infty, -n] \\ T[\Phi](n), & \xi \in [n, \infty] \end{cases}$$

which equals with T on the interval $[-n, n]$ and is constant outside this region.

Let $\{\Phi_i\}_{i \geq 0} \subset \Gamma$. Since T has uniformly bounded derivatives, the family

$$T^n[\Phi_i](\xi)$$

is, for each fixed n , uniformly bounded and equicontinuous with respect to the norm $\|\cdot\|_\mu$. By the Arzelà–Ascoli theorem, the sequence $T^n[\Phi_i](\xi)$ therefore admits a convergent subsequence. Consequently, for each n , the operator T^n is compact with respect to the norm $\|\cdot\|_\mu$.

What remains is to show that, as $n \rightarrow \infty$, $T^n \rightarrow T$ in the operator norm. We have

$$\begin{aligned} \|T^n - T\|_{op} &= \sup_{\Phi \in \Gamma; \|\Phi\|=1} \sup_{\xi \in \mathbb{R}} |T^n[\Phi] - T[\Phi]| e^{-\mu|\xi|} \\ &= \sup_{\Phi \in \Gamma; \|\Phi\|=1} \sup_{\xi \in [-\infty, -n] \cup [n, \infty]} |T^n[\Phi] - T[\Phi]| e^{-\mu|\xi|} \\ &\leq \frac{2}{\alpha} \max\{M_1, M_2, M_3\} e^{-\mu n} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Thus, by Theorem 3.4.2 in [11], T is also a compact operator. $\square$

Combining the results of the previous lemmas, we apply Schauder's Fixed Point Theorem [11] to establish the existence of a travelling wave solution for our system of equations (8).

Theorem 3.4. *If $c \geq \bar{c}$, then system (8) has a non-negative, non-trivial solution $(B(\xi), I(\xi), V(\xi)) \in \Gamma$ with $\lim_{\xi \rightarrow -\infty} (B(\xi), I(\xi), V(\xi)) = (0, 0, 0)$ and the solution stays bounded.*

We chose α , defined in (58), large enough and μ , from the definition of the norm (54), small enough. We will prove the theorem in the following steps.

Proof. Step 1: For $c > \bar{c}$ defined in (20), the previous lemmas satisfy the requirements of Schauder's fixed point theorem; thus, the operator T has at least one fixed point which we will denote as $(B, I, V) \in \Gamma$ and it satisfies the ODE system (8).

Step 2: For $c = \bar{c}$, we employ the limiting argument of Fang and Zhao [8] (Theorem 3.1). Choose a sequence $\{c_n\}_{n \geq 1} \subset (\bar{c}, \infty)$ such that

$$\lim_{n \rightarrow \infty} c_n = \bar{c}.$$

For each c_n , there exists a travelling-wave solution (B^n, I^n, V^n) of the ODE system (8). From the previous lemma, these solutions satisfy the uniform derivative bound

$$\begin{aligned} \left\| \frac{d}{d\xi} (B^n, I^n, V^n) \right\| &= \left\| \frac{d}{d\xi} T[(B^n, I^n, V^n)] \right\| \\ &\leq 2 \max \left\{ \frac{2M_1}{\sqrt{c_n^2 + 4D\alpha}}, \frac{2M_2}{\sqrt{c_n^2 + 4D\alpha}}, \frac{2M_3}{\sqrt{c_n^2 + 4\alpha}} \right\} \\ &\leq 2 \max \left\{ \frac{2M_1}{\sqrt{\bar{c}^2 + 4D\alpha}}, \frac{2M_2}{\sqrt{\bar{c}^2 + 4D\alpha}}, \frac{2M_3}{\sqrt{\bar{c}^2 + 4\alpha}} \right\}. \end{aligned}$$

Moreover, the bounds

$$0 \leq B^n \leq 1, \quad 0 \leq I^n \leq 1, \quad 0 \leq V^n \leq \frac{\theta}{\gamma}$$

imply that the sequence $\{(B^n, I^n, V^n)\}$ is uniformly bounded and equicontinuous. By the Arzelà–Ascoli theorem, there exists a subsequence $\{(B^{n_k}, I^{n_k}, V^{n_k})\}$ that converges pointwise to some $(\tilde{B}, \tilde{I}, \tilde{V}) \in \chi$. Consider

$$\begin{aligned} \lim_{k \rightarrow \infty} B^{n_k} &= \lim_{k \rightarrow \infty} T_1[(B^{n_k}, I^{n_k}, V^{n_k})] \\ &= \lim_{k \rightarrow \infty} \frac{1}{\sqrt{c_{n_k}^2 + 4D\alpha}} \left(\int_{-\infty}^{\xi} e^{\sigma_1^{n_k}(\xi-\eta)} F_1[(B^{n_k}, I^{n_k}, V^{n_k})] d\eta \right. \\ &\quad \left. + \int_{\xi}^{\infty} e^{\sigma_2^{n_k}(\xi-\eta)} F_1[(B^{n_k}, I^{n_k}, V^{n_k})] d\eta \right) \end{aligned} \quad (59)$$

where

$$\sigma_1^{n_k} = \frac{c_{n_k}k - \sqrt{c_{n_k}^2 + 4D\alpha}}{2D}, \quad \sigma_2^{n_k} = \frac{c_{n_k}k + \sqrt{c_{n_k}^2 + 4D\alpha}}{2D}.$$

Applying the Lebesgue Dominated Convergence Theorem to the above limit, we obtain

$$\tilde{B} = T_1[(\tilde{B}, \tilde{I}, \tilde{V})].$$

Analogous arguments yield

$$\tilde{I} = T_2[(\tilde{B}, \tilde{I}, \tilde{V})], \quad \tilde{V} = T_3[(\tilde{B}, \tilde{I}, \tilde{V})].$$

Consequently,

$$(\tilde{B}, \tilde{I}, \tilde{V}) = T[(\tilde{B}, \tilde{I}, \tilde{V})],$$

and hence $(\tilde{B}, \tilde{I}, \tilde{V})$ is a nonnegative solution of the ODE system (8).

Step 3: Note that for every $c > \bar{c}$ (B, I, V) are bounded above by their corresponding upper solutions.

$$\begin{aligned} 0 \leq B(\xi) &\leq \bar{B}(\xi) = e^{\xi\rho} \frac{c\rho + a + 1 - D\rho^2}{(c\rho + a - D\rho^2)(c\rho + 1 - D\rho^2)}, \\ 0 \leq I(\xi) &\leq \bar{I}(\xi) = e^{\xi\rho} \frac{1}{c\rho + a - D\rho^2}, \\ 0 \leq V(\xi) &\leq \bar{V}(\xi) = e^{\xi\rho}. \end{aligned}$$

In particular when $\xi \rightarrow -\infty$ we have

$$\begin{aligned} 0 &\leq \lim_{\xi \rightarrow -\infty} B(\xi) \leq \lim_{\xi \rightarrow -\infty} \bar{B}(\xi) = 0, \\ 0 &\leq \lim_{\xi \rightarrow -\infty} I(\xi) \leq \lim_{\xi \rightarrow -\infty} \bar{I}(\xi) = 0, \\ 0 &\leq \lim_{\xi \rightarrow -\infty} V(\xi) \leq \lim_{\xi \rightarrow -\infty} \bar{V}(\xi) = 0. \end{aligned}$$

Now consider the critical case $c = \bar{c}$. Let $\{c_n\}_{n \geq 1} \subset (\bar{c}, c_N]$ for some fixed N be any decreasing sequence with

$$c_n > \bar{c}, \quad \lim_{n \rightarrow \infty} c_n = \bar{c}.$$

For each c_n , let (B^n, I^n, V^n) denote the corresponding travelling-wave solution. Each component is uniformly bounded above by its upper solution:

$$0 \leq B^n(\xi) \leq \bar{B}^n(\xi), \quad 0 \leq I^n(\xi) \leq \bar{I}^n(\xi), \quad 0 \leq V^n(\xi) \leq \bar{V}^n(\xi).$$

Since $\bar{B}^n(\xi) \rightarrow 0$ as $\xi \rightarrow -\infty$ for each fixed n . Now we consider the pointwise supremum

$$\overline{\bar{B}}(\xi) := \limsup_{n \rightarrow \infty} \bar{B}^n(\xi).$$

If

$$\lim_{\xi \rightarrow -\infty} \overline{\bar{B}}(\xi) > 0,$$

then for each index $n > 0$ there would be a $k > n$ such that

$$\lim_{\xi \rightarrow -\infty} \bar{B}^k(\xi) > 0,$$

contradicting the fact that $e^{\xi\rho} \rightarrow 0$ as $\xi \rightarrow -\infty$. Hence,

$$\lim_{\xi \rightarrow -\infty} \overline{\bar{B}}(\xi) = 0.$$

Therefore,

$$0 \leq \lim_{\xi \rightarrow -\infty} B^n(\xi) = 0.$$

An identical argument applied to I^n and V^n shows that

$$\lim_{\xi \rightarrow -\infty} (B^n(\xi), I^n(\xi), V^n(\xi)) = (0, 0, 0).$$

Thus the limiting travelling wave at $c = \bar{c}$ connects to $(0, 0, 0)$ as $\xi \rightarrow -\infty$.

Step 4: Now we consider the other end of the interval i.e. $\xi \rightarrow \infty$. For all $c > \bar{c}$, the corresponding solutions are uniformly bounded between their upper and lower solutions. To treat the critical case $c = \bar{c}$, let $\{c_n\}_{n \geq 1} \subset (\bar{c}, c_N]$, for some fixed N , be a decreasing sequence such that

$$c_n > \bar{c}, \quad \lim_{n \rightarrow \infty} c_n = \bar{c}.$$

For each c_n , let (B^n, I^n, V^n) denote the associated travelling-wave solution. Each component is uniformly bounded by its corresponding upper and lower solutions, namely

$$\begin{aligned} 0 \leq B^n(\xi) \leq \bar{B}^n(\xi) \leq 1, \quad 0 \leq \underline{I}_n(\xi) \leq I^n(\xi) \leq \bar{I}^n(\xi) \leq 1, \\ 0 \leq \underline{V}_n(\xi) \leq V^n(\xi) \leq \bar{V}^n(\xi) \leq \frac{\theta}{\gamma}. \end{aligned}$$

By the convergence result in (59), the sequence $\{(B^n, I^n, V^n)\}_{n \geq 1}$ admits a limit

$$\lim_{n \rightarrow \infty} (B^n, I^n, V^n) = (\tilde{B}, \tilde{I}, \tilde{V}),$$

and this limit inherits the same bounds. Consequently, $(\tilde{B}, \tilde{I}, \tilde{V})$ is a travelling-wave solution of (8) with wave speed $\bar{c}$.

Step 5: In the last step, we need to show there are no nontrivial solutions with $B \equiv 0$ and $(I, V) \neq (0, 0)$. Assume, for contradiction, that $B(\xi) \equiv 0$ for all ξ while (I, V) is not identically zero. Substituting $B \equiv 0$ into the first equation of (7) yields

$$0 = I(\xi) + V(\xi) \quad \text{for all } \xi.$$

In our biological model I and V are nonnegative functions. Hence $I(\xi) \geq 0$ and $V(\xi) \geq 0$ for all ξ , and the identity $I + V \equiv 0$ forces $I \equiv 0$ and $V \equiv 0$. This contradicts the assumption that (I, V) is not identically zero. Therefore B cannot be identically zero. $\square$

4. Nonexistence for small speed $c \leq c_m$

The objective of this section is to demonstrate that travelling waves cease to exist for speeds $c \in [0, c_m)$, where c_m was defined in (21). Here we adapt the approach of [40,41]. We look at the phase space system of (8) and it is given by

$$\begin{aligned} B' &= W \\ W' &= \frac{c}{D}W + \frac{B}{D} - \frac{B^2}{D} + \frac{BI}{D} + \frac{BV}{D} - \frac{I}{D} - \frac{V}{D} \\ I' &= X \\ X' &= \frac{c}{D}X + a\frac{I}{D} - \frac{V}{D} + \frac{BV}{D} \\ V' &= Z \\ Z' &= cZ - \theta I + \gamma V, \end{aligned} \tag{60}$$

with boundary conditions

$$\begin{aligned} (B(-\infty), W(-\infty), I(-\infty), X(-\infty), V(-\infty), Z(-\infty)) &= ((0, 0, 0, 0, 0, 0) \\ \|(B(\infty), W(\infty), I(\infty), X(\infty), V(\infty), Z(\infty))\| &< \infty. \end{aligned}$$

Setting the right hand side as a vector valued function $g(B, W, I, X, V, Z)$, we can compute the Jacobian matrix Dg at $(0, 0, 0, 0, 0, 0)$ given by

$$\left(\begin{array}{cc|cccc} 0 & 1 & 0 & 0 & 0 & 0 \\ \frac{1}{D} & \frac{c}{D} & -\frac{1}{D} & 0 & -\frac{1}{D} & 0 \\ \hline 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{a}{D} & \frac{c}{D} & -\frac{1}{D} & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & -\theta & 0 & \gamma & c \end{array} \right).$$

It has a $2 + 4$ block structure, where the first block corresponds to the cancer only case. The characteristic equation for an eigenvalue ρ is given by

$$\begin{aligned} 0 = H(\rho) &= \\ \underbrace{\left(\rho^2 - \frac{c}{D}\rho - \frac{1}{D} \right)}_{H_1^*(\rho)} &\underbrace{\left[\frac{c^2}{D}\rho^2 + c \left(-\rho^3 - \frac{1}{D}\rho^3 + \frac{\gamma\rho}{D} + \frac{a\rho}{D} \right) + \rho^4 - \gamma\rho^2 - a\frac{\rho^2}{D} + \frac{a\gamma}{D} - \frac{\theta}{D} \right]}_{H_2^*(\rho)} \end{aligned} \tag{61}$$

Lemma 4.1. Consider $0 < c < c_m$. If $(B(\xi), W(\xi), I(\xi), X(\xi), V(\xi), Z(\xi))$ is a non-negative solution to (60), then there exists a positive constant γ such that

$$\sup_{\xi < 0} \{B(\xi)e^{-\gamma\xi}\} < \infty, \quad \sup_{\xi < 0} \{I(\xi)e^{-\gamma\xi}\} < \infty, \quad \sup_{\xi < 0} \{V(\xi)e^{-\gamma\xi}\} < \infty. \tag{62}$$

Proof. We show that the equilibrium point $P_0 = (0, 0, 0, 0, 0, 0)$ of the system (60) is hyperbolic. Then, by the Stable Manifold Theorem [25] we have exponential approach to the steady state. Suppose that P_0 is non-hyperbolic, meaning that the Jacobian admits an eigenvalue with zero real part. Denoting this eigenvalue by $\rho = ib$ and substituting into (61), we obtain

$$H(ib) = \left(-b^2 - \frac{1}{D} - \frac{icb}{D}\right) \left[i \left(cb^3 + \frac{cb^3}{D} + \frac{c\gamma b}{D} + \frac{cab}{D}\right) + b^4 + \gamma b^2 + \frac{ab^2}{D} + \frac{a\gamma}{D} - \frac{\theta}{D}\right],$$

which is nonzero for all $b \in \mathbb{R}$. Indeed, the first factor has strictly negative real part for all values of b , and hence is never zero. For the second factor, at $b = 0$ it is nonzero by the assumption $\theta > a\gamma$; for $b > 0$ the imaginary part is strictly positive and thus nonzero; and for $b < 0$ the imaginary part is strictly negative and thus nonzero. Since both factors of $H(ib)$ are nonzero for all $b \in \mathbb{R}$, it follows that $\rho = ib$ cannot be a root of $H(\rho) = 0$. Therefore, the equilibrium P_0 is hyperbolic. The Stable Manifold Theorem then implies that there exists a positive constant γ such that

$$\sup_{\xi < 0} \{B(\xi)e^{-\gamma\xi}\} < \infty, \quad \sup_{\xi < 0} \{I(\xi)e^{-\gamma\xi}\} < \infty, \quad \sup_{\xi < 0} \{V(\xi)e^{-\gamma\xi}\} < \infty. \quad \square$$

Lemma 4.2. Consider $H_1^*(\rho)$ and $H_2^*(\rho)$ as defined in (61). If $0 < c < c_m$, then $H_2^*(\rho) < 0$ for all $\rho \geq 0$. The first factor $H_1^*(\rho)$ has a unique positive root at

$$\rho_1^* = \frac{c}{2D} + \frac{1}{2} \sqrt{\frac{c^2}{D^2} + \frac{4}{D}}.$$

Proof. Since H_2^* is quadratic in c with a positive leading coefficient for all $\rho > 0$, any two real roots would necessarily produce an interval on which $H_2^*(\rho)$ is negative. Setting $H_2^*(\rho) = 0$ and solving for c yields

$$\begin{aligned} c_{\pm}(\rho) &= \frac{-\left(-\rho^3 - \frac{1}{D}\rho^3 + \frac{\gamma\rho}{D} + \frac{a\rho}{D}\right) \pm \sqrt{\left(-\rho^3 - \frac{1}{D}\rho^3 + \frac{\gamma\rho}{D} + \frac{a\rho}{D}\right)^2 - 4\frac{\rho^2}{D}\left(-\gamma\rho^2 - a\frac{\rho^2}{D} + \frac{a\gamma}{D} - \frac{\theta}{D}\right)}}{2\frac{\rho^2}{D}} \\ &= \frac{(\rho^2 + D\rho^2 - \gamma - a) \pm \sqrt{(\rho^2 - D\rho^2 - \gamma + a)^2 + 4\theta}}{2\rho}. \end{aligned}$$

The value c_m was defined as the infimum of the positive branch of these:

$$c_m = \inf_{\rho \geq 0} \frac{(\rho^2 + D\rho^2 - \gamma - a) + \sqrt{(\rho^2 - D\rho^2 - \gamma + a)^2 + 4\theta}}{2\rho}$$

Hence, an ordering exists such that $c_- < 0 < c_m \leq c_+$ for all $\rho > 0$. Evaluating at $\rho = 0$ gives

$$H_2^*(0) = \frac{a\gamma}{D} - \frac{\theta}{D},$$

which is strictly negative under the assumption $\theta > a\gamma$. Therefore, for all $0 < c < c_m$ and $\rho \geq 0$, we have $H_2^*(\rho) < 0$. $\square$

We adopt the approach of [32], originally developed in [40], and make use of the negative one-sided Laplace transform, defined as

$$\mathcal{L}[\phi(\cdot)](s) = \hat{\phi}(s) = \int_{-\infty}^0 e^{-s\xi} \phi(\xi) d\xi, \quad s \geq 0.$$

Here s serves as a measure of the rate at which $\phi(\xi)$ decays to zero as $\xi \rightarrow -\infty$. This interpretation is motivated by two key properties of the negative one-sided Laplace transform: its monotonicity and its behaviour under differentiation, both of which make it particularly well-suited for studying the decay rates of solutions to ODEs.

Suppose now that non-negative solutions to (8) (9) exist for $0 < c < c_m$. We apply the negative one-sided Laplace transform to examine their decay behaviour. Should the transform become unbounded for some $s > 0$, this would contradict the bounds established in (62) (see Lemma 4.1).

Moreover, if $\phi(\xi)$ is non-negative, then $\hat{\phi}(s)$ is non-decreasing in s , since by definition $\frac{d\hat{\phi}}{ds} \geq 0$ for $\xi < 0$. Consequently, $\hat{\phi}(s)$ is well-defined on an interval of the form $[0, s^*)$, where either $s^* < \infty$ and

$$\lim_{s \rightarrow s^*} \hat{\phi}(s) = \infty,$$

or $s^* = \infty$. To establish the non-existence of non-negative solutions for $0 < c < c_m$, we prove the following theorem.

Theorem 4.3. *No nontrivial solutions of system (8) (9) exist for $0 < c < c_m$.*

Proof. The proof proceeds by contradiction. Assume that solutions of (8) and (9) exist for $0 < c < c_m$. Applying the negative one-sided Laplace transform we get

$$\begin{aligned} \hat{B}(s) = \mathcal{L}[B](s) &= \int_{-\infty}^0 e^{-s\xi} B(\xi) d\xi, & \hat{I}(s) = \mathcal{L}[I](s) &= \int_{-\infty}^0 e^{-s\xi} I(\xi) d\xi, \\ \hat{V}(s) = \mathcal{L}[V](s) &= \int_{-\infty}^0 e^{-s\xi} V(\xi) d\xi. \end{aligned}$$

Since $\hat{B}$, $\hat{I}$, and $\hat{V}$ are non-decreasing in s each transform is either finite for all $s \geq 0$, or there exists a finite blow-up points s_B^* , s_I^* , and s_V^* such that

$$\lim_{s \rightarrow s_B^*} \hat{B} = \infty, \quad \lim_{s \rightarrow s_I^*} \hat{I} = \infty, \quad \text{and} \quad \lim_{s \rightarrow s_V^*} \hat{V} = \infty.$$

Furthermore, by Lemma 4.1, these values s_B^*, s_I^*, s_V^* are strictly positive, since for any non-negative solution of (60) there exists a constant $\gamma > 0$ such that

$$\hat{B} = \int_{-\infty}^0 e^{-s\xi} B(\xi) d\xi \leq \sup_{\xi < 0} \{B(\xi)e^{-\gamma\xi}\} \int_{-\infty}^0 e^{(\gamma-s)\xi} d\xi$$

which is finite for all $s \in [0, \gamma)$. An analogous observation holds for $\hat{I}$ and $\hat{V}$.

Hence if solutions to (8) (9) exist, their Laplace transform exist either in a non-empty interval $[0, s^*)$, or in $[0, \infty)$.

Applying the negative one-sided Laplace transform to the travelling wave system (8) then yields

$$\begin{aligned} (s^2 D - sc)\hat{B} + B'(0) + (sD - c)B(0) &= \mathcal{L}[B(-B + I + V)] - \mathcal{L}[-B + I + V] \\ (s^2 D - sc - a)\hat{I} + I'(0) + (sD - c)I(0) &= \mathcal{L}[BV] - \hat{V} \\ (s^2 - sc - \gamma)\hat{V} + V'(0) + (s - c)V(0) &= -\theta\hat{I} \end{aligned} \quad (63)$$

Step 1: We begin with the assumption $s_B^* = s_I^* = s_V^* = \infty$. Adding the equations in (63) yields

$$\begin{aligned} (s^2 D - sc)\hat{B} + (s^2 D - sc - a)\hat{I} + (s^2 - sc - \gamma)\hat{V} + (sD - c)[B(0) + I(0) + V(0)] \\ + B'(0) + I'(0) + V'(0) + \mathcal{L}[(1 - B)(B - I - 2V)] + \theta\hat{I} = 0. \end{aligned}$$

As (63) is assumed to be true for all $s \in [0, \infty)$ we need to have

$$\begin{aligned} 0 = \lim_{s \rightarrow \infty} (s^2 D - sc)\hat{B} + (s^2 D - sc - a)\hat{I} + (s^2 - sc - \gamma)\hat{V} + (sD - c)[B(0) + I(0) + V(0)] \\ + B'(0) + I'(0) + V'(0) + \mathcal{L}[(1 - B)(B - I - 2V)] + \theta\hat{I} \end{aligned}$$

For the nonlinear part, since $0 < B < 1$, $0 < I$ and $0 < V$, we estimate

$$\begin{aligned} |\mathcal{L}[(1 - B)(B - I - 2V)]| &= \left| \int_{-\infty}^0 e^{-s\xi} (1 - B(\xi))(B(\xi) - I(\xi) - 2V(\xi)) d\xi \right| \\ &\leq \int_{-\infty}^0 e^{-s\xi} |(1 - B(\xi))| |B(\xi) - I(\xi) - 2V(\xi)| d\xi \leq \int_{-\infty}^0 e^{-s\xi} |B(\xi) - I(\xi) - 2V(\xi)| d\xi \\ &\leq \int_{-\infty}^0 e^{-s\xi} |B(\xi)| + |I(\xi)| + 2|V(\xi)| d\xi = \hat{B} + \hat{I} + 2\hat{V}. \end{aligned}$$

Thus, we obtain

$$\begin{aligned}
0 &= \lim_{s \rightarrow \infty} (s^2 D - sc) \hat{B} + (s^2 D - sc - a) \hat{I} + (s^2 - sc - \gamma) \hat{V} + (sD - c)[B(0) + I(0) + V(0)] \\
&\quad + B'(0) + I'(0) + V'(0) + \mathcal{L}[(1 - B)(B - I - 2V)] + \theta \hat{I} \\
&\geq \lim_{s \rightarrow \infty} (s^2 D - sc) \hat{B} + (s^2 D - sc - a) \hat{I} + (s^2 - sc - \gamma) \hat{V} + (sD - c)[B(0) + I(0) + V(0)] \\
&\quad + B'(0) + I'(0) + V'(0) - |\mathcal{L}[(1 - B)(B - I - 2V)]| + \theta \hat{I} \\
&\geq \lim_{s \rightarrow \infty} (s^2 D - sc) \hat{B} + (s^2 D - sc - a) \hat{I} + (s^2 - sc - \gamma) \hat{V} + (sD - c)[B(0) + I(0) + V(0)] \\
&\quad + B'(0) + I'(0) + V'(0) - \hat{B} - \hat{I} - 2\hat{V} + \theta \hat{I} \\
&= \lim_{s \rightarrow \infty} (s^2 D - sc - 1) \hat{B} + (s^2 D - sc - a - 1 + \theta) \hat{I} + (s^2 - sc - \gamma - 2) \hat{V} \\
&\quad + (sD - c)[B(0) + I(0) + V(0)] + B'(0) + I'(0) + V'(0)
\end{aligned}$$

We have that

$$\begin{aligned}
0 &\geq \underbrace{\lim_{s \rightarrow \infty} (s^2 D - sc - 1) \hat{B} + (s^2 D - sc - a - 1 + \theta) \hat{I} + (s^2 - sc - \gamma - 2) \hat{V}}_{\rightarrow \infty} \\
&\quad + \underbrace{(sD - c)[B(0) + I(0) + V(0)]}_{\rightarrow \infty} + \underbrace{B'(0) + I'(0) + V'(0)}_{\text{finite}} \rightarrow \infty,
\end{aligned}$$

which yields a contradiction, and hence at least one of the s_B^* , s_I^* , s_V^* must be finite.

Step 2: Here we show that if one of the s_B^* , s_I^* , s_V^* is finite, then all three are finite and are equal. We again argue by contradiction. Suppose that $s_I^* \neq s_V^*$, and assume $s_I^* < s_V^*$. Taking the limit $s \rightarrow s_I^*$ in the third equation of (63), we obtain

$$\lim_{s \rightarrow s_I^*} \underbrace{(s^2 - sc - \gamma) \hat{V} + V'(0) + (s - c)V(0)}_{\text{finite}} + \underbrace{\theta \hat{I}}_{\rightarrow \infty} \neq 0$$

which contradicts the third equation of (63). Now, if we consider $s_I^* > s_V^*$, the second equation gives

$$\lim_{s \rightarrow s_V^*} \underbrace{(s^2 D - sc - a) \hat{I} + I'(0) + (sD - c)I(0) - \mathcal{L}[BV]}_{\text{finite}} + \underbrace{\hat{V}}_{\infty} \neq 0$$

which is again a contradiction to equation two of (63), and therefore $s_I^* = s_V^*$. Now suppose that $s_B^* \neq s_I^* = s_V^*$. In the case $s_I^* < s_B^*$, the first equation of (63) gives

$$\lim_{s \rightarrow s_I^*} \underbrace{(s^2 D - sc - 1) \hat{B} + B'(0) + (sD - c)B(0) - \mathcal{L}[B(-B + I + V)]}_{\text{finite}} + \underbrace{\hat{I} + \hat{V}}_{\rightarrow \infty} \neq 0$$

which yields a similar contradiction to the first equation of (63). Finally, suppose that $s_I^* = s_V^* > s_B^*$. Returning to the same equation, we write

$$\lim_{s \rightarrow s_B^*} \underbrace{(s^2 D - sc) \hat{B} + B'(0) + (sD - c)B(0) - \mathcal{L}[B(-B + I + V)] + \hat{I} + \hat{V}}_{\text{finite}} + \underbrace{\hat{B}}_{\rightarrow \infty} \neq 0$$

This final contradiction establishes that all thresholds are the same and we call it $s^* := s_B^* = s_I^* = s_V^*$.

Step 3: We now show that for $s \in [0, s^*]$ the nonlinear terms remain bounded. Indeed, by Lemma 4.1 we have

$$\begin{aligned} 0 < \mathcal{L}[B^2] &= \int_{-\infty}^0 e^{-\xi s} B^2(\xi) d\xi \leq \sup_{\xi < 0} \{B e^{-\gamma \xi}\} \int_{-\infty}^0 e^{(\gamma-s)\xi} B(\xi) d\xi \\ &= \sup_{\xi < 0} \{B e^{-\gamma \xi}\} \hat{B}(s - \gamma) < \infty. \end{aligned}$$

Taking the limit s approaching s^* , we obtain

$$\lim_{s \rightarrow s^*} \mathcal{L}[B^2] = \lim_{s \rightarrow s^*} \sup_{\xi < 0} \{B e^{-\gamma \xi}\} \hat{B}(s - \gamma) = \sup_{\xi < 0} \{B e^{-\gamma \xi}\} \hat{B}(s^* - \gamma) < \infty.$$

By an analogous argument, $\mathcal{L}[BI]$ and $\mathcal{L}[BV]$ are likewise bounded.

Step 4: Multiplying equations two and three of system (63) yields

$$\begin{aligned} (s^2 D - sc - a)(s^2 - sc - \gamma) \hat{I} \hat{V} &= \left(\underbrace{\mathcal{L}[BV] - I'(0) - (sD - c)I(0)}_{k_1} - \hat{V} \right) \\ &\quad \times \left(\underbrace{-V'(0) - (s - c)V(0)}_{k_2} - \theta \hat{I} \right). \end{aligned} \quad (64)$$

We now multiply the two sides of (64) with $(s^2 D - sc - 1) \hat{B}$ and we obtain

$$\begin{aligned} (s^2 D - sc - 1)(s^2 D - sc - a)(s^2 - sc - \gamma) \hat{B} \hat{I} \hat{V} &= (k_1 - \hat{V}) (k_2 - \theta \hat{I}) (s^2 D - sc - 1) \hat{B} \\ &= (k_1 k_2 - k_1 \theta \hat{I} - k_2 \hat{V} + \hat{I} \hat{V}) (s^2 D - sc - 1) \hat{B}, \end{aligned}$$

which implies

$$\begin{aligned} &\underbrace{(s^2 D - sc - 1) \left((s^2 D - sc - a)(s^2 - sc - \gamma) - \theta \right)}_{DH_2^*(s)} \hat{B} \hat{I} \hat{V} \\ &= (k_1 k_2 - k_1 \theta \hat{I} - k_2 \hat{V}) (s^2 D - sc - 1) \hat{B}. \end{aligned}$$

We rewrite the above equation in terms of the characteristic equation (61), and we have

$$H_2^*(s) = \frac{(k_1 k_2 - \theta l_1 \hat{I} - k_2 \hat{V}) (s^2 D - sc - 1)}{D \hat{I} \hat{V}}$$

Table 1

Comparison of theoretical and numerical wave speeds for $a > 1$, and $a < 1$.

Case	θ	Numerical c	minimum of $S(\rho)$ c_m	theoretical speed $\bar{c}$
$a < 1$	$\theta = 25$	$c_N = 0.705$	$c_m = 0.72$	$\bar{c} = 0.9687$
	$\theta = 150$	$c_N = 3.362$	$c_m = 3.2457$	$\bar{c} = 3.225$
	$\theta = 250$	$c_N = 4.4422$	$c_m = 4.2324$	$\bar{c} = 4.2324$
$a > 1$	$\theta = 150$	$c_N = 2.851$	$c_m = 2.8472$	$\bar{c} = 2.853$
	$\theta = 350$	$c_N = 4.982$	$c_m = 4.6961$	$\bar{c} = 4.6961$
	$\theta = 500$	$c_N = 4.4422$	$c_m = 5.5403$	$\bar{c} = 5.5403$

Taking the limit $s \rightarrow s^*$ with $s^* > 0$, we have $\lim_{s \rightarrow s^*} \hat{V} = \lim_{s \rightarrow s^*} \hat{I} = \infty$, and thus we obtain $H_2^*(s) \rightarrow 0$. Since k_1 and k_2 are finite, this contradicts Lemma 10, where we had shown $H_2^*(s) < 0$. Hence, no value $s^* > 0$ can be admitted without producing a contradiction with either (63) or the condition $H_2^*(s) < 0$. If $s^* = 0$, then $\hat{B}(s)$, $\hat{I}(s)$, and $\hat{V}(s)$ are unbounded for all $s > 0$, which contradicts the bounds established in Lemma 4.1. Therefore, no non-negative solution (B, I, V) to (8) exists for $0 < c < c_m$. $\square$

5. Numerics

In this section we show how our theoretical minimum wave speeds compare to the numerically observed wave speeds. To address this, in Fig. 4, we present travelling-wave simulations of the model, carried out in MATLAB using the pdepe solver. On the left column of Fig. 4, we run the simulations using the parameter values used previously by [2]. In that case the parameter a was greater than one and for γ they used the value $\frac{40}{3}$. We compare the difference in wave formation for smaller a value in the right column of Fig. 4.

We compare the numerical wave speeds c_N with the theoretical one $\bar{c}$. As shown in Table 1, the numerical speed agrees more closely with the minimum c_m of the theoretical characteristic curve $S(\rho)$ in most cases, for both $a > 1$ and $a < 1$. In some parameter regimes, the agreement is less precise, but this is expected: the assumptions required for the theorem and for constructing the upper and lower solutions prevent us from fully resolving the interval between c_m and $\bar{c}$ and a foggy regime exists between the two. This uncertainty contributes to the discrepancy between the numerically observed wave speeds and the theoretical minimum speed $\bar{c}$.

6. Conclusion

In this paper, we study a reaction–diffusion model designed to capture the spatiotemporal dynamics of oncolytic virotherapy (OVT) and tumor cell populations. The model, originally proposed by Baabdulla et al. [2], describes the interactions between uninfected cancer cells, infected cancer cells, and free virions. A central open question in OVT is whether the virus, when administered as a monotherapy, can spread through and eradicate the tumor before the tumor itself invades the surrounding tissue. Mathematical modelling provides a natural framework to explore this question by comparing the invasion speed of the virus with that of the cancer cells, and by identifying which viral parameters could be enhanced to tip the competition in favour of successful tumor clearance.

In this work, we rigorously establish the existence of travelling wave solutions for this non-cooperative three-component system, proving that such waves exist for all wave speeds $c \geq \bar{c}$ and do not exist for speeds $0 < c < c_m$, where $c_m \leq \bar{c}$. In cases where $c_m < \bar{c}$ we find an




Fig. 4. Solutions of the full model (2) for two different values of the parameter a . We are showing the solutions as (B, I, V) where $B = 1 - C$. The left column displays simulations for $a = \frac{10}{3} > 1$ at times $t = 8$ (dotted lines) and $t = 34$ (solid lines). Panel (a) ($\theta = 150$) exhibits no oscillations, while Panel (e) ($\theta = 350$) shows oscillatory behaviour has started. Panel (c) ($\theta = 500$) illustrates the dynamics in an oscillatory regime with longer oscillations. The right column shows numerical solutions for the case $a < 1$ at times $t = 8$ (dotted lines) and $t = 54$ (solid lines), illustrating the formation of oscillatory dynamics behind the travelling wave. For visual clarity, the variables B and I have been multiplied by a factor of 10. Panel (b) ($\theta = 25$) exhibits no oscillations, whereas Panel (d) ($\theta = 150$) shows oscillations converging toward the coexistence equilibrium. Panel (f) ($\theta = 250$) corresponds to a regime beyond the Hopf bifurcation, where oscillatory patterns arise.

undecided region $c \in (c_m, \bar{c})$, which leaves room for future research. While travelling waves have been extensively studied for both cooperative and non-cooperative systems, the present model poses additional challenges due to its three interacting variables and non-monotone structure, which impose technical restrictions in the construction of upper and lower solutions. As a result, the proof naturally leads to parameter regimes where existence can be established and others where the status of travelling waves remains unclear. These “foggy” regions highlight interesting mathematical questions for future investigation, particularly regarding whether the remaining gaps can be closed by refining the analytical framework. Finally, to complement the theoretical

analysis, we conduct numerical simulations that illustrate the full PDE dynamics, confirm the formation of travelling invasion fronts, and demonstrate how wave propagation depends on the viral parameters. Together, the analytical and numerical results provide a deeper understanding of the mechanisms that influence the success of OVT as a monotherapy and point toward avenues for optimizing viral design.

Acknowledgments

TH is supported through a Discovery Grant of the Natural Sciences and Engineering Research Council of Canada (NSERC), RGPIN-2023-04269.

Appendix A

Consider the cancer-only model

$$C_t = DC_{xx} + C(1 - C).$$

The travelling wave analysis for this model is well established [24,5] and invasion fronts of the form $C(x + ct)$ exist for $c > 2\sqrt{D}$. (Note that here we use the form $x + ct$, i.e. waves moving to the left for $c > 0$.) The wave is constructed through a heteroclinic orbit in phase space connecting a stable node at $(0, 0)$ to a saddle at $(1, 0)$. The saddle at $(1, 0)$ has the characteristic equation

$$H_1^*(\rho) = \rho^2 - \frac{c}{D}\rho - \frac{1}{D} = 0,$$

where ρ denotes the eigenvalue and, at the same time, the exponential decay or growth rate near the steady state. This saddle has a stable and an unstable manifold related to the negative and positive solutions of $H_1^*(\rho) = 0$. The stable direction corresponds to the heteroclinic connection mentioned above, while the unstable direction does not lead to a travelling wave connection, as shown many times in the classical literature [24,5]. (Note that due to the formulation in $x + ct$, as opposed to $x - ct$, the roles of stable and unstable manifolds are switched as compared to the more classical case of $x - ct$.)

The above expression for $H_1^*(\rho)$ is exactly the same term as in our model in (61). Hence this term corresponds to the cancer-only case and it does not lead to a valid invasion front.

Data availability

No data was used for the research described in the article.

References

- [1] S.M. Al-Tuwairqi, N.O. Al-Johani, E.A. Simbawa, Modeling dynamics of cancer virotherapy with immune response, *Adv. Differ. Equ.* (ISSN 1687-1839) (26) (2020) 438, <https://doi.org/10.1186/s13662-020-02893-6>.
- [2] A.A. Baabdulla, T. Hillen, Oscillations in a spatial oncolytic virus model, *Bull. Math. Biol.* (ISSN 0092-8240) 86 (8) (2023).
- [3] T. Bansod, T. Hillen, Spiral waves and turbulence in mathematical models of oncolytic virotherapy induced by Hopf bifurcation, *Front. Appl. Math. Stat.* (ISSN 2297-4687) 12 (2026).
- [4] D.K. Bhatt, T. Janzen, T. Daemen, F.J. Weissing, Modelling the spatial dynamics of oncolytic virotherapy in the presence of virus-resistant tumour cells, *PLoS Comput. Biol.* (ISSN 1553-7358) 18 (12) (2022) e1010076.

- [5] N.F. Britton, *Essential Mathematical Biology*, 1st ed. 2003. edition, Springer Undergraduate Mathematics Series, Springer London, London, ISBN 9781447100492, 2003.
- [6] R. Eftimie, J. Dushoff, B.W. Bridle, J.L. Bramson, D.J.D. Earn, Multi-stability and multi-instability phenomena in a mathematical model of tumor-immune-virus interactions, *Bull. Math. Biol.* (ISSN 0092-8240) 73 (12) (2011) 2932–2961, <https://doi.org/10.1007/s11538-011-9653-5>.
- [7] M. El-Hachem, S.W. McCue, M.J. Simpson, Travelling wave analysis of cellular invasion into surrounding tissues, *Physica D* (ISSN 0167-2789) 428 (2021) 133026.
- [8] J. Fang, X.-Q. Zhao, Monotone wavefronts for partially degenerate reaction-diffusion systems, *J. Dyn. Differ. Equ.* (ISSN 1040-7294) 21 (4) (2009) 663–680.
- [9] M. Feinberg, D. Terman, Traveling composition waves on isothermal catalyst surfaces, *Arch. Ration. Mech. Anal.* (ISSN 0003-9527) 116 (1) (1991) 35–69.
- [10] A. Friedman, J.P. Tian, G. Fulci, E.A. Chiocca, J. Wang, Glioma virotherapy: effects of innate immune suppression and increased viral replication capacity, *Cancer Res. (Chic. Ill.)* (ISSN 0008-5472) 66 (4) (2006) 2314–2319.
- [11] T. Hillen, *Elements of Applied Functional Analysis*, AMS Open Math Notes, American Mathematical Society, 2023, <https://www.ams.org/open-math-notes/omn-view-lisztng?listingId=111364>.
- [12] A.L. Jenner, P.S. Kim, F. Frascoli, Oncolytic virotherapy for tumours following a Gompertz growth law, *J. Theor. Biol.* 480 (2019) 129–140.
- [13] N.L. Komarova, D. Wodarz, ODE models for oncolytic virus dynamics, *J. Theor. Biol.* (ISSN 0022-5193) 263 (4) (2010) 530–543.
- [14] Y. Kuramoto, *Chemical Oscillations, Waves, and Turbulence*, Springer-Verlag, 1984.
- [15] J.H. Lagergren, J.T. Nardini, R.E. Baker, M.J. Simpson, K.B. Flores, Biologically-informed neural networks guide mechanistic modeling from sparse experimental data, *PLoS Comput. Biol.* (ISSN 1553-7358) 16 (12) (2020) e1008462.
- [16] S.A. Levin, H.C. Muller-Landau, R. Nathan, J. Chave, The ecology and evolution of seed dispersal: a theoretical perspective, *Annu. Rev. Ecol. Evol. Syst.* (ISSN 1543-592X) 34 (1) (2003) 575–604.
- [17] X. Liang, X.-Q. Zhao, Asymptotic speeds of spread and traveling waves for monotone semiflows with applications, *Commun. Pure Appl. Math.* (ISSN 0010-3640) 60 (1) (2007) 1–40.
- [18] D. Lin, Y. Shen, T. Liang, Oncolytic virotherapy: basic principles, recent advances and future directions, *Signal Transduct. Targeted Ther.* (ISSN 2059-3635) 8 (1) (2023) 156.
- [19] S. Ma, Traveling wavefronts for delayed reaction-diffusion systems via a fixed point theorem, *J. Differ. Equ.* (ISSN 0022-0396) 171 (2) (2001) 294–314.
- [20] R.L. Martuza, A. Malick, J.M. Markert, K.L. Ruffner, D.M. Coen, Experimental therapy of human glioma by means of a genetically engineered virus mutant, *Science* (ISSN 0036-8075) 252 (5007) (1991) 854–856.
- [21] N. Mohammadnejad, T. Hillen, Stacked waves and qualitative analysis of an oncolytic virus-immune model, *bioRxiv*, <https://doi.org/10.1101/2025.09.04.674336>, 2025.
- [22] N. Mohammadnejad, T. Hillen, Qualitative optimization of oncolytic virotherapy and immune therapy combination treatments, *Math. Biosci. Eng.* (ISSN 1551-0018) 23 (2) (2026) 388–420.
- [23] D. Morselli, M.E. Delitala, F. Frascoli, Agent-based and continuum models for spatial dynamics of infection by oncolytic viruses, *Bull. Math. Biol.* (ISSN 0092-8240) 85 (10) (2023) 92.
- [24] J.D. Murray, *Mathematical Biology, Interdisciplinary Applied Mathematics*, vol. 17–18, 3rd ed. edition, Springer, New York, ISBN 0387952233, 2002–2003.
- [25] L. Perko, *Differential Equations and Dynamical Systems*, 3rd ed. 2001. edition, Texts in Applied Mathematics, vol. 7, Springer New York, New York, ISBN 9781461300038, 2001.
- [26] P. Pooladvand, C.-O. Yun, A.-R. Yoon, P.S. Kim, F. Frascoli, The role of viral infectivity in oncolytic virotherapy outcomes: a mathematical study, *Math. Biosci.* (ISSN 0025-5564) 334 (2021) 108520, <https://doi.org/10.1016/j.mbs.2020.108520>.
- [27] B. Sandstede, A. Scheel, *Spiral Waves: Linear and Nonlinear Theory*, American Math. Soc., 2023.
- [28] J.A. Sherratt, J.D. Murray, Epidermal wound healing: the clinical implications of a simple mathematical model, *Cell Transplant.* (ISSN 0963-6897) 1 (5) (1992) 365.
- [29] J.G. Skellam, Random dispersal in theoretical populations, *Bull. Math. Biol.* (ISSN 0092-8240) 53 (1) (1991) 135–165.
- [30] J. Smoller, *Shock Waves and Reaction—Diffusion Equations*, 1st ed. 1983. edition, Grundlehren der mathematischen Wissenschaften, A Series of Comprehensive Studies in Mathematics, vol. 258, Springer New York, New York, NY, ISBN 1-4684-0152-1, 1983.
- [31] K.M. Storey, S.E. Lawler, T.L. Jackson, Modeling oncolytic viral therapy, immune checkpoint inhibition, and the complex dynamics of innate and adaptive immunity in glioblastoma treatment, *Front. Physiol.* (ISSN 1664-042X) 11 (2020) 151.

- [32] R. Thiessen, T. Hillen, Travelling wave solutions to a microtube-driven glioma invasion model, *J. Differ. Equ.* (ISSN 0022-0396) 452 (2026) 113759.
- [33] J.P. Tian, The replicability of oncolytic virus: defining conditions in tumor virotherapy, *Math. Biosci. Eng.* (ISSN 1551-0018) 8 (3) (2011) 841–860, <https://doi.org/10.3934/mbe.2011.8.841>.
- [34] A. Volpert, V.A. Volpert, V.A. Volpert, Traveling Wave Solutions of Parabolic Systems, *Translations of Mathematical Monographs*, vol. 140, American Mathematical Society, Providence, RI, ISBN 0-8218-4609-4, 1994.
- [35] H. Wang, Spreading speeds and traveling waves for non-cooperative reaction–diffusion systems, *J. Nonlinear Sci.* (ISSN 0938-8974) 21 (5) (2011) 747–783.
- [36] H.F. Weinberger, M.A. Lewis, B. Li, Analysis of linear determinacy for spread in cooperative models, *J. Math. Biol.* (ISSN 0303-6812) 45 (3) (2002) 183–218.
- [37] D. Wodarz, Viruses as antitumor weapons: defining conditions for tumor remission, *Cancer Res.* (ISSN 0008-5472) 61 (8) (04 2001) 3501–3507.
- [38] D. Wodarz, N.L. Komarova, Towards predictive computational models of oncolytic virus therapy: basis for experimental validation and model selection, *PLoS ONE* (ISSN 1932-6203) 4 (1) (2009) e4271.
- [39] J. Wu, X. Zou, Traveling wave fronts of reaction-diffusion systems with delay, *J. Dyn. Differ. Equ.* (ISSN 1040-7294) 13 (3) (2001) 651–687.
- [40] T. Zhang, W. Wang, Existence of traveling wave solutions for influenza model with treatment, *J. Math. Anal. Appl.* (ISSN 0022-247X) 419 (1) (2014) 469–495.
- [41] T. Zhang, W. Wang, K. Wang, Minimal wave speed for a class of non-cooperative diffusion–reaction system, *J. Differ. Equ.* (ISSN 0022-0396) 260 (3) (2016) 2763–2791.