



MAT1320-Linear Algebra

Lecture Notes

Special Types of Square Matrices

Mehmet E. KÖROĞLU

Fall 2024

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS

mkoroglu@yildiz.edu.tr

Table of contents

1. Special Types of Square Matrices

2. Complex Matrices

Special Types of Square Matrices

Periodic Matrix

A square matrix A such that $A^{k+1} = A$, for some positive integer k , is called a **periodic** matrix.

Periodic Matrix

A square matrix A such that $A^{k+1} = A$, for some positive integer k , is called a **periodic** matrix. For example,

$$A = \begin{pmatrix} 1 & -2 & -6 \\ -3 & 2 & 9 \\ 2 & 0 & -3 \end{pmatrix} \Rightarrow A^3 = A$$

is a periodic matrix of period 2.

Idempotent Matrix

A square matrix A such that $A^2 = A$ is called an **idempotent** matrix.

Idempotent Matrix

A square matrix A such that $A^2 = A$ is called an **idempotent** matrix. For example,

$$A = \begin{pmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{pmatrix} \Rightarrow A^2 = A$$

is an idempotent matrix.

Nilpotent Matrix

A square matrix A such that $A^k = 0$, for some positive integer k , is called a **nilpotent** matrix

Nilpotent Matrix

A square matrix A such that $A^k = 0$, for some positive integer k , is called a **nilpotent** matrix and k is called the nilpotency index of A .

Nilpotent Matrix

A square matrix A such that $A^k = 0$, for some positive integer k , is called a **nilpotent** matrix and k is called the nilpotency index of A . For example,

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \\ -1 & -2 & -3 \end{pmatrix} \Rightarrow A^2 = 0$$

is a nilpotent matrix of nilpotency index 2.

Involute or Involutary Matrix

A square matrix A such that $A^2 = I$ is called an **involute or involutory** matrix,

Involute or Involutary Matrix

A square matrix A such that $A^2 = I$ is called an **involute or involutory** matrix, where I is the identity matrix of the appropriate size.

Involute or Involutary Matrix

A square matrix A such that $A^2 = I$ is called an **involute or involutory** matrix, where I is the identity matrix of the appropriate size. For example,

$$A = \begin{pmatrix} -5 & -8 & 0 \\ 3 & 5 & 0 \\ 1 & 2 & -1 \end{pmatrix} \Rightarrow A^2 = I_3$$

is an involute matrix.

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero.

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero. Such a matrix is sometimes denoted by

$$D = \text{diag} (d_{11}, d_{22}, \dots, d_{nn})$$

where some or all the d_{ii} may be zero.

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero. Such a matrix is sometimes denoted by

$$D = \text{diag} (d_{11}, d_{22}, \dots, d_{nn})$$

where some or all the d_{ii} may be zero. For example,

$$\begin{pmatrix} 3 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 2 \end{pmatrix},$$

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero. Such a matrix is sometimes denoted by

$$D = \text{diag}(d_{11}, d_{22}, \dots, d_{nn})$$

where some or all the d_{ii} may be zero. For example,

$$\begin{pmatrix} 3 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \quad \begin{pmatrix} 4 & 0 \\ 0 & -5 \end{pmatrix},$$

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero. Such a matrix is sometimes denoted by

$$D = \text{diag}(d_{11}, d_{22}, \dots, d_{nn})$$

where some or all the d_{ii} may be zero. For example,

$$\begin{pmatrix} 3 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \quad \begin{pmatrix} 4 & 0 \\ 0 & -5 \end{pmatrix}, \quad \begin{pmatrix} 6 & & \\ & 0 & \\ & & -9 \\ & & & 8 \end{pmatrix}$$

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero. Such a matrix is sometimes denoted by

$$D = \text{diag}(d_{11}, d_{22}, \dots, d_{nn})$$

where some or all the d_{ii} may be zero. For example,

$$\begin{pmatrix} 3 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \quad \begin{pmatrix} 4 & 0 \\ 0 & -5 \end{pmatrix}, \quad \begin{pmatrix} 6 & & \\ & 0 & \\ & & -9 \\ & & & 8 \end{pmatrix}$$

are diagonal matrices, which may be represented, respectively, by

$$\text{diag}(3, -7, 2),$$

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero. Such a matrix is sometimes denoted by

$$D = \text{diag}(d_{11}, d_{22}, \dots, d_{nn})$$

where some or all the d_{ii} may be zero. For example,

$$\begin{pmatrix} 3 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \quad \begin{pmatrix} 4 & 0 \\ 0 & -5 \end{pmatrix}, \quad \begin{pmatrix} 6 & & \\ & 0 & \\ & & -9 \\ & & & 8 \end{pmatrix}$$

are diagonal matrices, which may be represented, respectively, by

$$\text{diag}(3, -7, 2), \quad \text{diag}(4, -5),$$

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero. Such a matrix is sometimes denoted by

$$D = \text{diag}(d_{11}, d_{22}, \dots, d_{nn})$$

where some or all the d_{ii} may be zero. For example,

$$\begin{pmatrix} 3 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \quad \begin{pmatrix} 4 & 0 \\ 0 & -5 \end{pmatrix}, \quad \begin{pmatrix} 6 & & \\ & 0 & \\ & & -9 \\ & & & 8 \end{pmatrix}$$

are diagonal matrices, which may be represented, respectively, by

$$\text{diag}(3, -7, 2), \quad \text{diag}(4, -5), \quad \text{diag}(6, 0, -9, 8)$$

Diagonal Matrices

A square matrix $D = [d_{ij}]$ is **diagonal** if its nondiagonal entries are all zero. Such a matrix is sometimes denoted by

$$D = \text{diag}(d_{11}, d_{22}, \dots, d_{nn})$$

where some or all the d_{ii} may be zero. For example,

$$\begin{pmatrix} 3 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \quad \begin{pmatrix} 4 & 0 \\ 0 & -5 \end{pmatrix}, \quad \begin{pmatrix} 6 & & \\ & 0 & \\ & & -9 \\ & & & 8 \end{pmatrix}$$

are diagonal matrices, which may be represented, respectively, by

$$\text{diag}(3, -7, 2), \quad \text{diag}(4, -5), \quad \text{diag}(6, 0, -9, 8)$$

Observe that patterns of 0's in the third matrix have been omitted.

Triangular Matrices

A square matrix $A = [a_{ij}]$ is **upper triangular or simply triangular** if all entries below the (main) diagonal are equal to 0 that is, if $a_{ij} = 0$ for $i > j$.

Triangular Matrices

A square matrix $A = [a_{ij}]$ is **upper triangular or simply triangular** if all entries below the (main) diagonal are equal to 0 that is, if $a_{ij} = 0$ for $i > j$. Generic upper triangular matrices of orders 2, 3, 4 are as follows:

$$\begin{pmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{pmatrix},$$

Triangular Matrices

A square matrix $A = [a_{ij}]$ is **upper triangular or simply triangular** if all entries below the (main) diagonal are equal to 0 that is, if $a_{ij} = 0$ for $i > j$. Generic upper triangular matrices of orders 2, 3, 4 are as follows:

$$\begin{pmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{pmatrix}, \quad \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix},$$

Triangular Matrices

A square matrix $A = [a_{ij}]$ is **upper triangular or simply triangular** if all entries below the (main) diagonal are equal to 0 that is, if $a_{ij} = 0$ for $i > j$. Generic upper triangular matrices of orders 2, 3, 4 are as follows:

$$\begin{pmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{pmatrix}, \quad \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix}, \quad \begin{pmatrix} c_{11} & c_{12} & c_{13} & c_{14} \\ & c_{22} & c_{23} & c_{24} \\ & & c_{33} & c_{34} \\ & & & c_{44} \end{pmatrix}$$

Triangular Matrices

A square matrix $A = [a_{ij}]$ is **upper triangular or simply triangular** if all entries below the (main) diagonal are equal to 0 that is, if $a_{ij} = 0$ for $i > j$. Generic upper triangular matrices of orders 2, 3, 4 are as follows:

$$\begin{pmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{pmatrix}, \quad \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix}, \quad \begin{pmatrix} c_{11} & c_{12} & c_{13} & c_{14} \\ & c_{22} & c_{23} & c_{24} \\ & & c_{33} & c_{34} \\ & & & c_{44} \end{pmatrix}$$

As with diagonal matrices, it is common practice to omit patterns of 0's.

Triangular Matrices

A square matrix $A = [a_{ij}]$ is **upper triangular or simply triangular** if all entries below the (main) diagonal are equal to 0 that is, if $a_{ij} = 0$ for $i > j$. Generic upper triangular matrices of orders 2, 3, 4 are as follows:

$$\begin{pmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{pmatrix}, \quad \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix}, \quad \begin{pmatrix} c_{11} & c_{12} & c_{13} & c_{14} \\ & c_{22} & c_{23} & c_{24} \\ & & c_{33} & c_{34} \\ & & & c_{44} \end{pmatrix}$$

As with diagonal matrices, it is common practice to omit patterns of 0's.

Note: A lower triangular matrix is a square matrix whose entries above the diagonal are all zero.

Symmetric and Skew-Symmetric Matrices

- A matrix A is **symmetric** if $A^T = A$.

Symmetric and Skew-Symmetric Matrices

- A matrix A is **symmetric** if $A^T = A$. Equivalently, $A = [a_{ij}]$ is symmetric if symmetric elements (mirror elements with respect to the diagonal) are equal-that is, if each $a_{ij} = a_{ji}$.

Symmetric and Skew-Symmetric Matrices

- A matrix A is **symmetric** if $A^T = A$. Equivalently, $A = [a_{ij}]$ is symmetric if symmetric elements (mirror elements with respect to the diagonal) are equal-that is, if each $a_{ij} = a_{ji}$.
- A matrix A is **skew-symmetric** if $A^T = -A$

Symmetric and Skew-Symmetric Matrices

- A matrix A is **symmetric** if $A^T = A$. Equivalently, $A = [a_{ij}]$ is symmetric if symmetric elements (mirror elements with respect to the diagonal) are equal-that is, if each $a_{ij} = a_{ji}$.
- A matrix A is **skew-symmetric** if $A^T = -A$ or, equivalently, if each $a_{ij} = -a_{ji}$.

Symmetric and Skew-Symmetric Matrices

- A matrix A is **symmetric** if $A^T = A$. Equivalently, $A = [a_{ij}]$ is symmetric if symmetric elements (mirror elements with respect to the diagonal) are equal—that is, if each $a_{ij} = a_{ji}$.
- A matrix A is **skew-symmetric** if $A^T = -A$ or, equivalently, if each $a_{ij} = -a_{ji}$. Clearly, the diagonal elements of such a matrix must be zero, because $a_{ii} = -a_{ii}$ implies $a_{ii} = 0$.

Symmetric and Skew-Symmetric Matrices

- A matrix A is **symmetric** if $A^T = A$. Equivalently, $A = [a_{ij}]$ is symmetric if symmetric elements (mirror elements with respect to the diagonal) are equal-that is, if each $a_{ij} = a_{ji}$.
- A matrix A is **skew-symmetric** if $A^T = -A$ or, equivalently, if each $a_{ij} = -a_{ji}$. Clearly, the diagonal elements of such a matrix must be zero, because $a_{ii} = -a_{ii}$ implies $a_{ii} = 0$.
- Note that a matrix A must be square if $A^T = A$ or $A^T = -A$.

Symmetric and Skew-Symmetric Matrices

Example

$$\text{Let } A = \begin{pmatrix} 2 & -3 & 5 \\ -3 & 6 & 7 \\ 5 & 7 & -8 \end{pmatrix}, B = \begin{pmatrix} 0 & 3 & -4 \\ -3 & 0 & 5 \\ 4 & -5 & 0 \end{pmatrix},$$
$$C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Symmetric and Skew-Symmetric Matrices

Example

$$\text{Let } A = \begin{pmatrix} 2 & -3 & 5 \\ -3 & 6 & 7 \\ 5 & 7 & -8 \end{pmatrix}, B = \begin{pmatrix} 0 & 3 & -4 \\ -3 & 0 & 5 \\ 4 & -5 & 0 \end{pmatrix},$$
$$C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- i. By inspection, the symmetric elements in A are equal, or $A^T = A$. Thus, A is symmetric.

Symmetric and Skew-Symmetric Matrices

Example

$$\text{Let } A = \begin{pmatrix} 2 & -3 & 5 \\ -3 & 6 & 7 \\ 5 & 7 & -8 \end{pmatrix}, B = \begin{pmatrix} 0 & 3 & -4 \\ -3 & 0 & 5 \\ 4 & -5 & 0 \end{pmatrix},$$
$$C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- i. By inspection, the symmetric elements in A are equal, or $A^T = A$. Thus, A is symmetric.
- ii. The diagonal elements of B are 0 and symmetric elements are negatives of each other, or $B^T = -B$. Thus, B is skew-symmetric.

Symmetric and Skew-Symmetric Matrices

Example

$$\text{Let } A = \begin{pmatrix} 2 & -3 & 5 \\ -3 & 6 & 7 \\ 5 & 7 & -8 \end{pmatrix}, B = \begin{pmatrix} 0 & 3 & -4 \\ -3 & 0 & 5 \\ 4 & -5 & 0 \end{pmatrix},$$
$$C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- i. By inspection, the symmetric elements in A are equal, or $A^T = A$. Thus, A is symmetric.
- ii. The diagonal elements of B are 0 and symmetric elements are negatives of each other, or $B^T = -B$. Thus, B is skew-symmetric.
- iii. Because C is not square, C is neither symmetric nor skew-symmetric.

Orthogonal Matrices

A real matrix A is **orthogonal** if $A^T = A^{-1}$ — that is, if $AA^T = A^T A = I$. Thus, A must necessarily be square and invertible.

Orthogonal Matrices

A real matrix A is **orthogonal** if $A^T = A^{-1}$ — that is, if $AA^T = A^T A = I$. Thus, A must necessarily be square and invertible.

Example

Let $A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$.

Orthogonal Matrices

A real matrix A is **orthogonal** if $A^T = A^{-1}$ — that is, if $AA^T = A^T A = I$. Thus, A must necessarily be square and invertible.

Example

Let $A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$. Multiplying A by A^T yields I ; that is, $AA^T = I$.

Orthogonal Matrices

A real matrix A is **orthogonal** if $A^T = A^{-1}$ — that is, if $AA^T = A^T A = I$. Thus, A must necessarily be square and invertible.

Example

Let $A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$. Multiplying A by A^T yields I ; that is, $AA^T = I$. This means $A^T A = I$, as well. Thus, $A^T = A^{-1}$; that is, A is orthogonal.

Complex Matrices

Let A be a matrix with complex entries.

Complex Matrices

Let A be a matrix with complex entries. Recall that if $z = a + bi$ is a complex number, then $\bar{z} = a - bi$ is its conjugate.

Complex Matrices

Let A be a matrix with complex entries. Recall that if $z = a + bi$ is a complex number, then $\bar{z} = a - bi$ is its conjugate.

The conjugate of a complex matrix A , written \bar{A} , is the matrix obtained from A by taking the conjugate of each entry in A .

Complex Matrices

Let A be a matrix with complex entries. Recall that if $z = a + bi$ is a complex number, then $\bar{z} = a - bi$ is its conjugate.

The conjugate of a complex matrix A , written \bar{A} , is the matrix obtained from A by taking the conjugate of each entry in A .

That is, if $A = [a_{ij}]$, then $\bar{A} = [b_{ij}]$ where $b_{ij} = \bar{a}_{ij}$.

Complex Matrices

Let A be a matrix with complex entries. Recall that if $z = a + bi$ is a complex number, then $\bar{z} = a - bi$ is its conjugate.

The conjugate of a complex matrix A , written \bar{A} , is the matrix obtained from A by taking the conjugate of each entry in A .

That is, if $A = [a_{ij}]$, then $\bar{A} = [\bar{a}_{ij}]$ where $\bar{a}_{ij} = \overline{a_{ij}}$. We denote this fact by writing $\bar{A} = [\bar{a}_{ij}]$.

Complex Matrices

Let A be a matrix with complex entries. Recall that if $z = a + bi$ is a complex number, then $\bar{z} = a - bi$ is its conjugate.

The conjugate of a complex matrix A , written \bar{A} , is the matrix obtained from A by taking the conjugate of each entry in A .

That is, if $A = [a_{ij}]$, then $\bar{A} = [\bar{a}_{ij}]$ where $\bar{a}_{ij} = \overline{a_{ij}}$. We denote this fact by writing $\bar{A} = [\bar{a}_{ij}]$.

The two operations of transpose and conjugation commute for any complex matrix A , and the special notation A^H is used for the **conjugate transpose** of A .

Complex Matrices

Let A be a matrix with complex entries. Recall that if $z = a + bi$ is a complex number, then $\bar{z} = a - bi$ is its conjugate.

The conjugate of a complex matrix A , written \bar{A} , is the matrix obtained from A by taking the conjugate of each entry in A .

That is, if $A = [a_{ij}]$, then $\bar{A} = [\bar{a}_{ij}]$ where $\bar{a}_{ij} = \overline{a_{ij}}$. We denote this fact by writing $\bar{A} = [\bar{a}_{ij}]$.

The two operations of transpose and conjugation commute for any complex matrix A , and the special notation A^H is used for the **conjugate transpose** of A . That is, $A^H = (\bar{A})^T = \overline{(A^T)}$

Complex Matrices

Let A be a matrix with complex entries. Recall that if $z = a + bi$ is a complex number, then $\bar{z} = a - bi$ is its conjugate.

The conjugate of a complex matrix A , written \bar{A} , is the matrix obtained from A by taking the conjugate of each entry in A .

That is, if $A = [a_{ij}]$, then $\bar{A} = [\bar{a}_{ij}]$ where $\bar{a}_{ij} = \overline{a_{ij}}$. We denote this fact by writing $\bar{A} = [\bar{a}_{ij}]$.

The two operations of transpose and conjugation commute for any complex matrix A , and the special notation A^H is used for the **conjugate transpose** of A . That is, $A^H = (\bar{A})^T = \overline{(A^T)}$

Note that if A is real, then $A^H = A^T$.

Complex Matrices

Let A be a matrix with complex entries. Recall that if $z = a + bi$ is a complex number, then $\bar{z} = a - bi$ is its conjugate.

The conjugate of a complex matrix A , written \bar{A} , is the matrix obtained from A by taking the conjugate of each entry in A .

That is, if $A = [a_{ij}]$, then $\bar{A} = [\bar{a}_{ij}]$ where $\bar{a}_{ij} = \overline{a_{ij}}$. We denote this fact by writing $\bar{A} = [\bar{a}_{ij}]$.

The two operations of transpose and conjugation commute for any complex matrix A , and the special notation A^H is used for the **conjugate transpose** of A . That is, $A^H = (\bar{A})^T = \overline{(A^T)}$

Note that if A is real, then $A^H = A^T$. Some texts use A^* instead of A^H .

Example

Let

$$A = \begin{pmatrix} 2 + 8i & 5 - 3i & 4 - 7i \\ 6i & 1 - 4i & 3 + 2i \end{pmatrix}.$$

Example

Let

$$A = \begin{pmatrix} 2 + 8i & 5 - 3i & 4 - 7i \\ 6i & 1 - 4i & 3 + 2i \end{pmatrix}.$$

Then

$$A^H = \begin{pmatrix} 2 - 8i & -6i \\ 5 + 3i & 1 + 4i \\ 4 + 7i & 3 - 2i \end{pmatrix}.$$

Hermitian and Unitary Matrices

A complex matrix A is said to be **Hermitian** if $A^H = A$.

Hermitian and Unitary Matrices

A complex matrix A is said to be **Hermitian** if $A^H = A$. Clearly, $A = [a_{ij}]$ is Hermitian if and only if symmetric elements are conjugate;

Hermitian and Unitary Matrices

A complex matrix A is said to be **Hermitian** if $A^H = A$. Clearly, $A = [a_{ij}]$ is Hermitian if and only if symmetric elements are conjugate; that is, if each $a_{ij} = \bar{a}_{ji}$, in which case each diagonal element a_{ii} must be real.

Hermitian and Unitary Matrices

A complex matrix A is said to be **Hermitian** if $A^H = A$. Clearly, $A = [a_{ij}]$ is Hermitian if and only if symmetric elements are conjugate; that is, if each $a_{ij} = \bar{a}_{ji}$, in which case each diagonal element a_{ii} must be real.

A complex matrix A is said to be **skew-Hermitian** if $A^H = -A$.

Hermitian and Unitary Matrices

A complex matrix A is said to be **Hermitian** if $A^H = A$. Clearly, $A = [a_{ij}]$ is Hermitian if and only if symmetric elements are conjugate; that is, if each $a_{ij} = \bar{a}_{ji}$, in which case each diagonal element a_{ii} must be real.

A complex matrix A is said to be **skew-Hermitian** if $A^H = -A$. Similarly, if A is skew-symmetric, then each diagonal element $a_{ii} = 0$ or pure complex.

Hermitian and Unitary Matrices

A complex matrix A is said to be **Hermitian** if $A^H = A$. Clearly, $A = [a_{ij}]$ is Hermitian if and only if symmetric elements are conjugate; that is, if each $a_{ij} = \bar{a}_{ji}$, in which case each diagonal element a_{ii} must be real.

A complex matrix A is said to be **skew-Hermitian** if $A^H = -A$. Similarly, if A is skew-symmetric, then each diagonal element $a_{ii} = 0$ or pure complex.

Note that A must be square if $A^H = A$ or $A^H = -A$.

Hermitian and Unitary Matrices

A complex matrix A is said to be **Hermitian** if $A^H = A$. Clearly, $A = [a_{ij}]$ is Hermitian if and only if symmetric elements are conjugate; that is, if each $a_{ij} = \bar{a}_{ji}$, in which case each diagonal element a_{ii} must be real.

A complex matrix A is said to be **skew-Hermitian** if $A^H = -A$. Similarly, if A is skew-symmetric, then each diagonal element $a_{ii} = 0$ or pure complex.

Note that A must be square if $A^H = A$ or $A^H = -A$.

A complex matrix A is **unitary** if $A^H A = A A^H = I$; that is, if $A^H = A^{-1}$.

Hermitian and Unitary Matrices

Example

Consider the following complex matrices:

$$A = \begin{pmatrix} 3 & 1 - 2i & 4 + 7i \\ 1 + 2i & -4 & -2i \\ 4 - 7i & 2i & 5 \end{pmatrix}, \quad B = \frac{1}{2} \begin{pmatrix} 1 & -i & -1 + i \\ i & 1 & 1 + i \\ 1 + i & -1 + i & 0 \end{pmatrix}$$

Hermitian and Unitary Matrices

Example

Consider the following complex matrices:

$$A = \begin{pmatrix} 3 & 1 - 2i & 4 + 7i \\ 1 + 2i & -4 & -2i \\ 4 - 7i & 2i & 5 \end{pmatrix}, \quad B = \frac{1}{2} \begin{pmatrix} 1 & -i & -1 + i \\ i & 1 & 1 + i \\ 1 + i & -1 + i & 0 \end{pmatrix}$$

- i. By inspection, the diagonal elements of A are real, and the symmetric elements $1 - 2i$ and $1 + 2i$ are conjugate, $4 + 7i$ and $4 - 7i$ are conjugate, and $-2i$ and $2i$ are conjugate. Thus, A is Hermitian.

Hermitian and Unitary Matrices

Example

Consider the following complex matrices:

$$A = \begin{pmatrix} 3 & 1-2i & 4+7i \\ 1+2i & -4 & -2i \\ 4-7i & 2i & 5 \end{pmatrix}, \quad B = \frac{1}{2} \begin{pmatrix} 1 & -i & -1+i \\ i & 1 & 1+i \\ 1+i & -1+i & 0 \end{pmatrix}$$

- i. By inspection, the diagonal elements of A are real, and the symmetric elements $1-2i$ and $1+2i$ are conjugate, $4+7i$ and $4-7i$ are conjugate, and $-2i$ and $2i$ are conjugate. Thus, A is Hermitian.
- ii. Multiplying B by B^H yields I ; that is, $BB^H = I$. This implies $B^HB = I$, as well. Thus, $B^H = B^{-1}$ which means B is unitary.

?