



MAT1320-Linear Algebra

Lecture Notes

Echelon Form of a Matrix

Mehmet E. KÖROĞLU
Fall 2024

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS
mkoroglu@yildiz.edu.tr

Table of contents

1. Echelon Form of a Matrix
2. Elementary Row (Column) Operations
3. Finding an Inverse using Elementary Row Operations

Echelon Form of a Matrix

Echelon Form of a Matrix

An $m \times n$ matrix A is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

Echelon Form of a Matrix

An $m \times n$ matrix A is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.

Echelon Form of a Matrix

An $m \times n$ matrix A is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.
- b) The first nonzero entry from the left of a nonzero row is a 1.

Echelon Form of a Matrix

An $m \times n$ matrix A is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.
- b) The first nonzero entry from the left of a nonzero row is a 1. This entry is called a **leading one** of its row.

Echelon Form of a Matrix

An $m \times n$ matrix A is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.
- b) The first nonzero entry from the left of a nonzero row is a 1. This entry is called a **leading one** of its row.
- c) For each nonzero row, the leading one appears to the right and below any leading ones in preceding rows.

Echelon Form of a Matrix

An $m \times n$ matrix A is said to be in **reduced row echelon form (RREF)** if it satisfies the following properties:

- a) All zero rows, if there are any, appear at the bottom of the matrix.
- b) The first nonzero entry from the left of a nonzero row is a 1. This entry is called a **leading one** of its row.
- c) For each nonzero row, the leading one appears to the right and below any leading ones in preceding rows.
- d) If a column contains a leading one, then all other entries in that column are zero.

Echelon Form of a Matrix

- A matrix in reduced row echelon form appears as a **staircase** ("echelon") pattern of leading ones descending from the upper left corner of the matrix.

Echelon Form of a Matrix

- A matrix in reduced row echelon form appears as a **staircase ("echelon")** pattern of leading ones descending from the upper left corner of the matrix.
- An $m \times n$ matrix satisfying properties a), b), and c) is said to be in **row echelon form (REF)**. There may be no zero rows.

Echelon Form of a Matrix

- A matrix in reduced row echelon form appears as a staircase ("echelon") pattern of leading ones descending from the upper left corner of the matrix.
- An $m \times n$ matrix satisfying properties a), b), and c) is said to be in row echelon form (REF). There may be no zero rows.
- A similar definition can be formulated in the obvious manner for reduced column echelon form and column echelon form.

Echelon Form of a Matrix

Example

The following are matrices in reduced row echelon form, since they satisfy properties *a)*, *b)*, and *d)*:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

Echelon Form of a Matrix

Example

The following are matrices in reduced row echelon form, since they satisfy properties *a)*, *b)*, and *d)*:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 0 & 0 & -2 & 4 \\ 0 & 1 & 0 & 0 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Echelon Form of a Matrix

Example

The following are matrices in reduced row echelon form, since they satisfy properties *a)*, *b)*, and *d)*:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 0 & 0 & -2 & 4 \\ 0 & 1 & 0 & 0 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

and

$$C = \begin{pmatrix} 1 & 2 & 0 & 0 & 1 \\ 0 & 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Echelon Form of a Matrix

Example

The matrices that follow are not in reduced row echelon form.

(Why not?)

$$D = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -3 \end{pmatrix},$$

Echelon Form of a Matrix

Example

The matrices that follow are not in reduced row echelon form.

(Why not?)

$$D = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -3 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 2 & -2 & 5 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Echelon Form of a Matrix

Example

The matrices that follow are not in reduced row echelon form.
(Why not?)

$$D = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -3 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 2 & -2 & 5 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

$$F = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 1 & -2 & 5 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

Echelon Form of a Matrix

Example

The matrices that follow are not in reduced row echelon form.
(Why not?)

$$D = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -3 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 2 & -2 & 5 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

$$F = \begin{pmatrix} 1 & 0 & 3 & 4 \\ 0 & 1 & -2 & 5 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad G = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & -2 & 5 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Echelon Form of a Matrix

Example

The following are matrices in row echelon form.

$$H = \begin{pmatrix} 1 & 5 & 0 & 2 & -2 & 4 \\ 0 & 1 & 0 & 3 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

Echelon Form of a Matrix

Example

The following are matrices in row echelon form.

$$H = \begin{pmatrix} 1 & 5 & 0 & 2 & -2 & 4 \\ 0 & 1 & 0 & 3 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Echelon Form of a Matrix

Example

The following are matrices in row echelon form.

$$H = \begin{pmatrix} 1 & 5 & 0 & 2 & -2 & 4 \\ 0 & 1 & 0 & 3 & 4 & 8 \\ 0 & 0 & 0 & 1 & 7 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Note: We shall now show that every matrix can be put into row (column) echelon form, or into reduced row (column) echelon form, by means of certain row (column) operations.

Elementary Row (Column) Operations

Elementary Row (Column) Operations

An elementary row (column) operation on a matrix A is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).

Elementary Row (Column) Operations

An elementary row (column) operation on a matrix A is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.

Elementary Row (Column) Operations

An elementary row (column) operation on a matrix A is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

Elementary Row (Column) Operations

An elementary row (column) operation on a matrix A is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

We now introduce the following notation for elementary row and elementary column operations on matrices:

Elementary Row (Column) Operations

An elementary row (column) operation on a matrix A is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

We now introduce the following notation for elementary row and elementary column operations on matrices:

- Interchange rows (columns) i and j , **Type I:** $\mathbf{r}_i \leftrightarrow \mathbf{r}_j$ ($\mathbf{c}_i \leftrightarrow \mathbf{c}_j$)

Elementary Row (Column) Operations

An elementary row (column) operation on a matrix A is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

We now introduce the following notation for elementary row and elementary column operations on matrices:

- Interchange rows (columns) i and j , **Type I:** $\mathbf{r}_i \leftrightarrow \mathbf{r}_j$ ($\mathbf{c}_i \leftrightarrow \mathbf{c}_j$)
- Replace row (column) i by k times row (column) i , **Type II:**
 $k\mathbf{r}_i \rightarrow \mathbf{r}_i$ ($k\mathbf{c}_i \rightarrow \mathbf{c}_i$)

Elementary Row (Column) Operations

An elementary row (column) operation on a matrix A is any one of the following operations:

1. **Type I:** Interchange any two rows (columns).
2. **Type II:** Multiply a row (column) by a nonzero number.
3. **Type III:** Add a multiple of one row (column) to another.

We now introduce the following notation for elementary row and elementary column operations on matrices:

- Interchange rows (columns) i and j , **Type I:** $\mathbf{r}_i \leftrightarrow \mathbf{r}_j$ ($\mathbf{c}_i \leftrightarrow \mathbf{c}_j$)
- Replace row (column) i by k times row (column) i , **Type II:**
 $k\mathbf{r}_i \rightarrow \mathbf{r}_i$ ($k\mathbf{c}_i \rightarrow \mathbf{c}_i$)
- Replace row (column) j by k times row (column) i + row (column) j , **Type III:** $k\mathbf{r}_i + \mathbf{r}_j \rightarrow \mathbf{r}_j$ ($k\mathbf{c}_i + \mathbf{c}_j \rightarrow \mathbf{c}_j$)

Echelon Form of a Matrix

Example

$$\text{Let } A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}.$$

Echelon Form of a Matrix

Example

$$\text{Let } A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}.$$

Interchanging rows 1 and 3 of A , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Echelon Form of a Matrix

Example

$$\text{Let } A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}.$$

Interchanging rows 1 and 3 of A , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Multiplying the third row of A by $\frac{1}{3}$, we obtain

$$C = A_{\frac{1}{3}r_3 \rightarrow r_3} = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 1 & 1 & 2 & -3 \end{pmatrix}$$

Echelon Form of a Matrix

Adding (-2) times row 2 of A to row 3 of A , we obtain

$$D = A_{-2r_2+r_3 \rightarrow r_3} = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ -1 & -3 & 6 & -5 \end{pmatrix}$$

Echelon Form of a Matrix

Adding (-2) times row 2 of A to row 3 of A , we obtain

$$D = A_{-2r_2+r_3 \rightarrow r_3} = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ -1 & -3 & 6 & -5 \end{pmatrix}$$

Observe that in obtaining D from A , row 2 of A did not change.

Echelon Form of a Matrix

Adding (-2) times row 2 of A to row 3 of A , we obtain

$$D = A_{-2r_2+r_3 \rightarrow r_3} = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ -1 & -3 & 6 & -5 \end{pmatrix}$$

Observe that in obtaining D from A , row 2 of A did not change.

Note: An $m \times n$ matrix B is said to be row (column) equivalent to an $m \times n$ matrix A if B can be produced by applying a finite sequence of elementary row (column) operations to A .

Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of A to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of A to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so B is row equivalent to A .

Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of A to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so B is row equivalent to A . Multiplying row 1 of B by 2, we obtain

$$C = B_{2r_1 \rightarrow r_1} = \begin{pmatrix} 2 & 4 & 8 & 6 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of A to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so B is row equivalent to A . Multiplying row 1 of B by 2, we obtain

$$C = B_{2r_1 \rightarrow r_1} = \begin{pmatrix} 2 & 4 & 8 & 6 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so C is row equivalent to B .

Example

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 2 & 1 & 3 & 2 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

If we add 2 times row 3 of A to its second row, we obtain

$$B = A_{2r_3+r_2 \rightarrow r_2} = \begin{pmatrix} 1 & 2 & 4 & 3 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so B is row equivalent to A . Multiplying row 1 of B by 2, we obtain

$$C = B_{2r_1 \rightarrow r_1} = \begin{pmatrix} 2 & 4 & 8 & 6 \\ 4 & -3 & 7 & 8 \\ 1 & -2 & 2 & 3 \end{pmatrix}$$

so C is row equivalent to B . It then follows that C is row equivalent to A , since we obtained C by applying two successive elementary row operations to A .

Echelon Form of a Matrix

Theorem

Every nonzero $m \times n$ matrix $\mathbf{A} = [a_{ij}]$ is row (column) equivalent to a matrix in row (column) echelon form.

Echelon Form of a Matrix

Theorem

Every nonzero $m \times n$ matrix $\mathbf{A} = [a_{ij}]$ is row (column) equivalent to a matrix in row (column) echelon form.

Theorem

The RREF of a matrix is unique.

Echelon Form of a Matrix

Theorem

Every nonzero $m \times n$ matrix $\mathbf{A} = [a_{ij}]$ is row (column) equivalent to a matrix in row (column) echelon form.

Theorem

The RREF of a matrix is unique.

Note: In any matrix, the first column with a nonzero entry is called the **pivot column**; the first nonzero entry in the pivot column is called the **pivot**.

Echelon Form of a Matrix

Theorem

Every nonzero $m \times n$ matrix $\mathbf{A} = [a_{ij}]$ is row (column) equivalent to a matrix in row (column) echelon form.

Theorem

The RREF of a matrix is unique.

Note: In any matrix, the first column with a nonzero entry is called the **pivot column**; the first nonzero entry in the pivot column is called the **pivot**.

Example

Let $A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}$. Find REF and RREF of A .

Echelon Form of a Matrix

Theorem

Every nonzero $m \times n$ matrix $\mathbf{A} = [a_{ij}]$ is row (column) equivalent to a matrix in row (column) echelon form.

Theorem

The RREF of a matrix is unique.

Note: In any matrix, the first column with a nonzero entry is called the **pivot column**; the first nonzero entry in the pivot column is called the **pivot**.

Example

Let $A = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 2 & 3 & 0 & -2 \\ 3 & 3 & 6 & -9 \end{pmatrix}$. Find REF and RREF of A .

Example (cont.)

Interchanging rows 1 and 3 of A , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Example (cont.)

Interchanging rows 1 and 3 of A , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Multiplying the first row of B by $\frac{1}{3}$, we obtain

$$C = B_{\frac{1}{3}r_1 \rightarrow r_1} = \begin{pmatrix} 1 & 1 & 2 & -3 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Example (cont.)

Interchanging rows 1 and 3 of A , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Multiplying the first row of B by $\frac{1}{3}$, we obtain

$$C = B_{\frac{1}{3}r_1 \rightarrow r_1} = \begin{pmatrix} 1 & 1 & 2 & -3 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding (-2) times row 1 of C to row 2 of C , we obtain

$$D = C_{-2r_1 + r_2 \rightarrow r_2} = \underbrace{\begin{pmatrix} 1 & 1 & 2 & -3 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{REF}}$$

Example (cont.)

Interchanging rows 1 and 3 of A , we obtain

$$B = A_{r_1 \leftrightarrow r_3} = \begin{pmatrix} 3 & 3 & 6 & -9 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Multiplying the first row of B by $\frac{1}{3}$, we obtain

$$C = B_{\frac{1}{3}r_1 \rightarrow r_1} = \begin{pmatrix} 1 & 1 & 2 & -3 \\ 2 & 3 & 0 & -2 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding (-2) times row 1 of C to row 2 of C , we obtain

$$D = C_{-2r_1 + r_2 \rightarrow r_2} = \underbrace{\begin{pmatrix} 1 & 1 & 2 & -3 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{REF}}$$

Example (cont.)

Adding (-2) times row 3 of D to row 1 of D and 4 times row 3 of D to row 2 of D , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \begin{pmatrix} 1 & 1 & 0 & -7 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Example (cont.)

Adding (-2) times row 3 of D to row 1 of D and 4 times row 3 of D to row 2 of D , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \begin{pmatrix} 1 & 1 & 0 & -7 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding (-1) times row 2 of E to row 1 of E , we obtain

$$F = E_{-r_2+r_1 \rightarrow r_1} = \underbrace{\begin{pmatrix} 1 & 0 & 0 & -19 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{RREF}}$$

Example (cont.)

Adding (-2) times row 3 of D to row 1 of D and 4 times row 3 of D to row 2 of D , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \begin{pmatrix} 1 & 1 & 0 & -7 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding (-1) times row 2 of E to row 1 of E , we obtain

$$F = E_{-r_2+r_1 \rightarrow r_1} = \underbrace{\begin{pmatrix} 1 & 0 & 0 & -19 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{RREF}}$$

Note: The number of nonzero rows in the REF or RREF of a matrix \mathbf{A} is called the **rank** of \mathbf{A} and denoted by $\text{rank}(\mathbf{A})$.

Example (cont.)

Adding (-2) times row 3 of D to row 1 of D and 4 times row 3 of D to row 2 of D , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \begin{pmatrix} 1 & 1 & 0 & -7 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

Adding (-1) times row 2 of E to row 1 of E , we obtain

$$F = E_{-r_2+r_1 \rightarrow r_1} = \underbrace{\begin{pmatrix} 1 & 0 & 0 & -19 \\ 0 & 1 & 0 & 12 \\ 0 & 0 & 1 & 2 \end{pmatrix}}_{\text{RREF}}$$

Note: The number of nonzero rows in the REF or RREF of a matrix \mathbf{A} is called the **rank** of \mathbf{A} and denoted by **rank** (\mathbf{A}). The

rank of matrix \mathbf{A} given in the previous example is 3.

Finding an Inverse using Elementary Row Operations

Finding an Inverse using Elementary Row Operations

- If the inverse of a matrix \mathbf{A} exists, then there is a series of elementary row operations that reduces \mathbf{A} to the identity matrix.

Finding an Inverse using Elementary Row Operations

- If the inverse of a matrix \mathbf{A} exists, then there is a series of elementary row operations that reduces \mathbf{A} to the identity matrix.
- Suppose the required elementary row operations are (in order) $\mathbf{E}_1, \mathbf{E}_2, \dots, \mathbf{E}_n$, then

$$\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} = \mathbf{I}$$

which means that $\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 = \mathbf{A}^{-1}$.

Finding an Inverse using Elementary Row Operations

- If the inverse of a matrix \mathbf{A} exists, then there is a series of elementary row operations that reduces \mathbf{A} to the identity matrix.
- Suppose the required elementary row operations are (in order) $\mathbf{E}_1, \mathbf{E}_2, \dots, \mathbf{E}_n$, then

$$\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} = \mathbf{I}$$

which means that $\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 = \mathbf{A}^{-1}$.

- Furthermore, because

$$\mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{I} = \mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1$$

we can use the following technique:

Finding an Inverse using Elementary Row Operations

- Write \mathbf{A} and \mathbf{I} side-by-side.

Finding an Inverse using Elementary Row Operations

- Write **A** and **I** side-by-side.
- Then apply the same row operations to both **A** and **I** until **A** is reduced to the identity matrix.

Finding an Inverse using Elementary Row Operations

- Write **A** and **I** side-by-side.
- Then apply the same row operations to both **A** and **I** until **A** is reduced to the identity matrix.
- At the same time, the identity matrix will be "reduced" to the inverse matrix.

$$\begin{array}{c|c} \mathbf{A} & \mathbf{I} \\ \mathbf{E}_1\mathbf{A} & \mathbf{E}_1\mathbf{I} \\ \mathbf{E}_2\mathbf{E}_1\mathbf{A} & \mathbf{E}_2\mathbf{E}_1\mathbf{I} \\ \underbrace{\mathbf{E}_n \dots \mathbf{E}_2\mathbf{E}_1\mathbf{A}}_{\text{reduced to I}} & \underbrace{\mathbf{E}_n \dots \mathbf{E}_2\mathbf{E}_1\mathbf{I}}_{\mathbf{A}^{-1}} \end{array}$$

Finding an Inverse using Elementary Row Operations

- Write **A** and **I** side-by-side.
- Then apply the same row operations to both **A** and **I** until **A** is reduced to the identity matrix.
- At the same time, the identity matrix will be "reduced" to the inverse matrix.

$$\begin{array}{c|c} \mathbf{A} & \mathbf{I} \\ \hline \mathbf{E}_1 \mathbf{A} & \mathbf{E}_1 \mathbf{I} \\ \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} & \mathbf{E}_2 \mathbf{E}_1 \mathbf{I} \\ \mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} & \mathbf{E}_n \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{I} \\ \hline \underbrace{\hspace{10em}}_{\text{reduced to } \mathbf{I}} & \underbrace{\hspace{10em}}_{\mathbf{A}^{-1}} \end{array}$$

Here is the fully worked out example:

Example

Let $A = \begin{pmatrix} 3 & 3 & 6 \\ 2 & 3 & 0 \\ 0 & 0 & 1 \end{pmatrix}$. By using elementary row operations, find the inverse of the A .

Example

Let $A = \begin{pmatrix} 3 & 3 & 6 \\ 2 & 3 & 0 \\ 0 & 0 & 1 \end{pmatrix}$. By using elementary row operations, find the inverse of the A .

$$B = [A|I_3] = \left(\begin{array}{ccc|ccc} 3 & 3 & 6 & 1 & 0 & 0 \\ 2 & 3 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Example

Let $A = \begin{pmatrix} 3 & 3 & 6 \\ 2 & 3 & 0 \\ 0 & 0 & 1 \end{pmatrix}$. By using elementary row operations, find the inverse of the A .

$$B = [A|I_3] = \left(\begin{array}{ccc|ccc} 3 & 3 & 6 & 1 & 0 & 0 \\ 2 & 3 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Multiplying the first row of B by $\frac{1}{3}$, we obtain

$$C = B_{\frac{1}{3}r_1 \rightarrow r_1} = \left(\begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 2 & 3 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Example (cont.)

Adding (-2) times row 1 of C to row 2 of C , we obtain

$$D = C_{-2r_1+r_2 \rightarrow r_2} \left(\begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 0 & 1 & -4 & \frac{-2}{3} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Example (cont.)

Adding (-2) times row 1 of C to row 2 of C , we obtain

$$D = C_{-2r_1+r_2 \rightarrow r_2} \left(\begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 0 & 1 & -4 & \frac{-2}{3} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding (-2) times row 3 of D to row 1 of D and 4 times row 3 of D to row 2 of D , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \left(\begin{array}{ccc|ccc} 1 & 1 & 0 & \frac{1}{3} & 0 & -2 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Example (cont.)

Adding (-2) times row 1 of C to row 2 of C , we obtain

$$D = C_{-2r_1+r_2 \rightarrow r_2} \left(\begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 0 & 1 & -4 & \frac{-2}{3} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding (-2) times row 3 of D to row 1 of D and 4 times row 3 of D to row 2 of D , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \left(\begin{array}{ccc|ccc} 1 & 1 & 0 & \frac{1}{3} & 0 & -2 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding (-1) times row 2 of E to row 1 of E , we obtain

$$E = \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -1 & -6 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Example (cont.)

Adding (-2) times row 1 of C to row 2 of C , we obtain

$$D = C_{-2r_1+r_2 \rightarrow r_2} \left(\begin{array}{ccc|ccc} 1 & 1 & 2 & \frac{1}{3} & 0 & 0 \\ 0 & 1 & -4 & \frac{-2}{3} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding (-2) times row 3 of D to row 1 of D and 4 times row 3 of D to row 2 of D , we obtain

$$E = D_{\substack{4r_3+r_2 \rightarrow r_2 \\ -2r_3+r_1 \rightarrow r_1}} = \left(\begin{array}{ccc|ccc} 1 & 1 & 0 & \frac{1}{3} & 0 & -2 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

Adding (-1) times row 2 of E to row 1 of E , we obtain

$$E = \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -1 & -6 \\ 0 & 1 & 0 & \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right), \quad A^{-1} = \left(\begin{array}{ccc} 1 & -1 & -6 \\ \frac{-2}{3} & 1 & 4 \\ 0 & 0 & 1 \end{array} \right).$$

?