



MAT1320-Linear Algebra

Lecture Notes

Eigenvalues and Eigenvectors, Characteristic Polynomial,
Diagonalization, Cayley-Hamilton Theorem

Mehmet E. KÖROĞLU

Fall 2024

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS

mkoroglu@yildiz.edu.tr

Table of contents

1. Eigenvalues and Eigenvectors
2. The Characteristic Equation
3. Finding Eigenvalues and Eigenvectors
4. Similar Matrices
5. Diagonalization
6. Cayley-Hamilton Theorem

Eigenvalues and Eigenvectors

Eigenvalues and Eigenvectors

Definition

A scalar λ is called an **eigenvalue** of the $n \times n$ square matrix A if there is a nontrivial solution \vec{x} of $A\vec{x} = \lambda\vec{x}$.

Eigenvalues and Eigenvectors

Definition

A scalar λ is called an **eigenvalue** of the $n \times n$ square matrix A if there is a nontrivial solution \vec{x} of $A\vec{x} = \lambda\vec{x}$. Such an \vec{x} is called an **eigenvector** corresponding to the eigenvalue λ .

Eigenvalues and Eigenvectors

Definition

A scalar λ is called an **eigenvalue** of the $n \times n$ square matrix A if there is a nontrivial solution \vec{x} of $A\vec{x} = \lambda\vec{x}$. Such an \vec{x} is called an **eigenvector** corresponding to the eigenvalue λ .

Note: Note that an eigenvector cannot be $\vec{0}$, but an eigenvalue can be $0 \in \mathbb{R}$. If 0 is an eigenvalue of A , then there must be some nontrivial vector \vec{x} for which $A\vec{x} = 0\vec{x} = \vec{0}$ which implies that A is not invertible.

Eigenvalues and Eigenvectors

Definition

A scalar λ is called an **eigenvalue** of the $n \times n$ square matrix A if there is a nontrivial solution \vec{x} of $A\vec{x} = \lambda\vec{x}$. Such an \vec{x} is called an **eigenvector** corresponding to the eigenvalue λ .

Note: Note that an eigenvector cannot be $\vec{0}$, but an eigenvalue can be $0 \in \mathbb{R}$. If 0 is an eigenvalue of A , then there must be some nontrivial vector \vec{x} for which $A\vec{x} = 0\vec{x} = \vec{0}$ which implies that A is not invertible.

Theorem

The $n \times n$ square matrix A is invertible if and only if 0 is not an eigenvalue of A .

Eigenvalues and Eigenvectors

Definition

A scalar λ is called an **eigenvalue** of the $n \times n$ square matrix A if there is a nontrivial solution \vec{x} of $A\vec{x} = \lambda\vec{x}$. Such an \vec{x} is called an **eigenvector** corresponding to the eigenvalue λ .

Note: Note that an eigenvector cannot be $\vec{0}$, but an eigenvalue can be $0 \in \mathbb{R}$. If 0 is an eigenvalue of A , then there must be some nontrivial vector \vec{x} for which $A\vec{x} = 0\vec{x} = \vec{0}$ which implies that A is not invertible.

Theorem

The $n \times n$ square matrix A is invertible if and only if 0 is not an eigenvalue of A .

Note: The **eigenspace** of the $n \times n$ matrix A corresponding to the eigenvalue λ of A is the set of all eigenvectors of A corresponding to λ .

The Characteristic Equation

The Characteristic Equation

Let A be an $n \times n$ square matrix. Then we have

$$A\vec{x} = \lambda\vec{x}$$

The Characteristic Equation

Let A be an $n \times n$ square matrix. Then we have

$$A\vec{x} = \lambda\vec{x} \Rightarrow \lambda\vec{x} - A\vec{x} = \vec{0}$$

The Characteristic Equation

Let A be an $n \times n$ square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow \lambda\vec{x} - A\vec{x} = \vec{0} \\ \lambda I_n \vec{x} - A\vec{x} &= \vec{0} \end{aligned}$$

The Characteristic Equation

Let A be an $n \times n$ square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow \lambda\vec{x} - A\vec{x} = \vec{0} \\ \lambda I_n \vec{x} - A\vec{x} &= \vec{0} \Rightarrow (\lambda I_n - A) \vec{x} = \vec{0} \end{aligned}$$

The Characteristic Equation

Let A be an $n \times n$ square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow \lambda\vec{x} - A\vec{x} = \vec{0} \\ \lambda I_n \vec{x} - A\vec{x} &= \vec{0} \Rightarrow (\lambda I_n - A) \vec{x} = \vec{0} \end{aligned}$$

Note: A scalar λ is an eigenvalue of an $n \times n$ matrix A if and only if λ satisfies the equation $\det(\lambda I - A) = 0$.

The Characteristic Equation

Let A be an $n \times n$ square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow \lambda\vec{x} - A\vec{x} = \vec{0} \\ \lambda I_n \vec{x} - A\vec{x} &= \vec{0} \Rightarrow (\lambda I_n - A) \vec{x} = \vec{0} \end{aligned}$$

Note: A scalar λ is an eigenvalue of an $n \times n$ matrix A if and only if λ satisfies the equation $\det(\lambda I - A) = 0$.

Definition

Let A be an $n \times n$ square matrix.

The Characteristic Equation

Let A be an $n \times n$ square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow \lambda\vec{x} - A\vec{x} = \vec{0} \\ \lambda I_n \vec{x} - A\vec{x} &= \vec{0} \Rightarrow (\lambda I_n - A) \vec{x} = \vec{0} \end{aligned}$$

Note: A scalar λ is an eigenvalue of an $n \times n$ matrix A if and only if λ satisfies the equation $\det(\lambda I - A) = 0$.

Definition

Let A be an $n \times n$ square matrix. Then the equation $\det(\lambda I - A) = 0$ is called the **characteristic equation** of the matrix A and the result of the determinant $\det(\lambda I - A)$ is polynomial of the form $P_A(\lambda) = \lambda^n + a_{n-1}\lambda^{n-1} + \dots + a_1\lambda + a_0$ is called **characteristic polynomial** of A .

The Characteristic Equation

Let A be an $n \times n$ square matrix. Then we have

$$\begin{aligned} A\vec{x} &= \lambda\vec{x} \Rightarrow \lambda\vec{x} - A\vec{x} = \vec{0} \\ \lambda I_n \vec{x} - A\vec{x} &= \vec{0} \Rightarrow (\lambda I_n - A) \vec{x} = \vec{0} \end{aligned}$$

Note: A scalar λ is an eigenvalue of an $n \times n$ matrix A if and only if λ satisfies the equation $\det(\lambda I - A) = 0$.

Definition

Let A be an $n \times n$ square matrix. Then the equation $\det(\lambda I - A) = 0$ is called the **characteristic equation** of the matrix A and the result of the determinant $\det(\lambda I - A)$ is polynomial of the form $P_A(\lambda) = \lambda^n + a_{n-1}\lambda^{n-1} + \dots + a_1\lambda + a_0$ is called **characteristic polynomial** of A .

Note: $P_A(\lambda) = \lambda^n + (-1)^{n-1} \text{Tr}(A) \lambda^{n-1} + \dots + \det(A)$. For example, for a 2×2 square matrix A ,

$$P_A(\lambda) = \lambda^2 - \text{Tr}(A) \lambda + \det(A).$$

Finding Eigenvalues and Eigenvectors

Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial $|\lambda I_n - A| = P_A(\lambda)$ of A .

Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial $|\lambda I_n - A| = P_A(\lambda)$ of A .
2. Find the roots of $P_A(\lambda)$ to find the eigenvalues of A .

Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial $|\lambda I_n - A| = P_A(\lambda)$ of A .
2. Find the roots of $P_A(\lambda)$ to find the eigenvalues of A .
3. Repeat (a) and (b) for each eigenvalue λ of A .

Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial $|\lambda I_n - A| = P_A(\lambda)$ of A .
2. Find the roots of $P_A(\lambda)$ to find the eigenvalues of A .
3. Repeat (a) and (b) for each eigenvalue λ of A .
 - (a) Form the matrix $M = \lambda I - A$ by subtracting λ down the diagonal of A .

Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial $|\lambda I_n - A| = P_A(\lambda)$ of A .
2. Find the roots of $P_A(\lambda)$ to find the eigenvalues of A .
3. Repeat (a) and (b) for each eigenvalue λ of A .
 - (a) Form the matrix $M = \lambda I - A$ by subtracting λ down the diagonal of A .
 - (b) Find a basis for the solution space of the homogeneous system $M\mathbf{X} = 0$.

Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial $|\lambda I_n - A| = P_A(\lambda)$ of A .
2. Find the roots of $P_A(\lambda)$ to find the eigenvalues of A .
3. Repeat (a) and (b) for each eigenvalue λ of A .
 - (a) Form the matrix $M = \lambda I - A$ by subtracting λ down the diagonal of A .
 - (b) Find a basis for the solution space of the homogeneous system $M\mathbf{X} = 0$.

Note 1: The eigenvalues of a triangular matrix are the entries on its main diagonal.

Finding Eigenvalues and Eigenvectors

1. Find characteristic polynomial $|\lambda I_n - A| = P_A(\lambda)$ of A .
2. Find the roots of $P_A(\lambda)$ to find the eigenvalues of A .
3. Repeat (a) and (b) for each eigenvalue λ of A .
 - (a) Form the matrix $M = \lambda I - A$ by subtracting λ down the diagonal of A .
 - (b) Find a basis for the solution space of the homogeneous system $M\mathbf{X} = 0$.

Note 1: The eigenvalues of a triangular matrix are the entries on its main diagonal.

Note 2: If $P_A(\lambda)$ has multiple roots, then there exists multiple eigenvalues.

Finding Eigenvalues and Eigenvectors

Example

Let $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$. Find

1. characteristic polynomial,

Finding Eigenvalues and Eigenvectors

Example

Let $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$. Find

1. characteristic polynomial,
2. eigenvalues,

Finding Eigenvalues and Eigenvectors

Example

Let $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$. Find

1. characteristic polynomial,
2. eigenvalues,
3. eigenvectors.

Finding Eigenvalues and Eigenvectors

Solution (1)

$$P_A(\lambda) = \det(\lambda I_3 - A)$$

$$\lambda I_3 - A = \lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$$

Finding Eigenvalues and Eigenvectors

Solution (1)

$$P_A(\lambda) = \det(\lambda I_3 - A)$$

$$\begin{aligned}\lambda I_3 - A &= \lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} \\ &= \begin{pmatrix} \lambda - 3 & -6 & 8 \\ 0 & \lambda & -6 \\ 0 & 0 & \lambda - 2 \end{pmatrix}\end{aligned}$$

Finding Eigenvalues and Eigenvectors

Solution (1)

$$P_A(\lambda) = \det(\lambda I_3 - A)$$

$$\begin{aligned}\lambda I_3 - A &= \lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix} \\ &= \begin{pmatrix} \lambda - 3 & -6 & 8 \\ 0 & \lambda & -6 \\ 0 & 0 & \lambda - 2 \end{pmatrix} \\ \Rightarrow P_A(\lambda) &= \lambda(\lambda - 2)(\lambda - 3).\end{aligned}$$

Finding Eigenvalues and Eigenvectors

Solution (2)

Since, eigenvalues are the roots of $P_A(\lambda)$, we have

$$P_A(\lambda) = \lambda(\lambda - 2)(\lambda - 3) = 0$$

Finding Eigenvalues and Eigenvectors

Solution (2)

Since, eigenvalues are the roots of $P_A(\lambda)$, we have

$$\begin{aligned}P_A(\lambda) &= \lambda(\lambda - 2)(\lambda - 3) = 0 \\ \Rightarrow \lambda_1 &= 0, \lambda_2 = 2, \lambda_3 = 3.\end{aligned}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_1 = 0$,

$$(0I_3 - A) = -A = \begin{pmatrix} -3 & -6 & 8 \\ 0 & 0 & -6 \\ 0 & 0 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_1 = 0$,

$$(0I_3 - A) = -A = \begin{pmatrix} -3 & -6 & 8 \\ 0 & 0 & -6 \\ 0 & 0 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow A\vec{x} = \vec{0}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_1 = 0$,

$$(0I_3 - A) = -A = \begin{pmatrix} -3 & -6 & 8 \\ 0 & 0 & -6 \\ 0 & 0 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow A\vec{x} = \vec{0} \Rightarrow \begin{cases} x + 2y = 0 \\ z = 0 \end{cases}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_1 = 0$,

$$(0I_3 - A) = -A = \begin{pmatrix} -3 & -6 & 8 \\ 0 & 0 & -6 \\ 0 & 0 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow A\vec{x} = \vec{0} \Rightarrow \begin{cases} x + 2y = 0 \\ z = 0 \end{cases}$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = t \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, t \in \mathbb{R} \Rightarrow \vec{v}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}.$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_2 = 2$,

$$(2I_3 - A) = \begin{pmatrix} -1 & -6 & 8 \\ 0 & 2 & -6 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 10 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{pmatrix}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_2 = 2$,

$$(2I_3 - A) = \begin{pmatrix} -1 & -6 & 8 \\ 0 & 2 & -6 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 10 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (2I_n - A) \vec{x} = \vec{0}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_2 = 2$,

$$(2I_3 - A) = \begin{pmatrix} -1 & -6 & 8 \\ 0 & 2 & -6 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 10 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (2I_n - A) \vec{x} = \vec{0} \Rightarrow \begin{cases} x + 10z = 0 \\ y - 3z = 0 \end{cases}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_2 = 2$,

$$(2I_3 - A) = \begin{pmatrix} -1 & -6 & 8 \\ 0 & 2 & -6 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 10 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (2I_n - A) \vec{x} = \vec{0} \Rightarrow \begin{cases} x + 10z = 0 \\ y - 3z = 0 \end{cases}$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = t \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}, t \in \mathbb{R} \Rightarrow \vec{v}_2 = \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}.$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_3 = 3$,

$$(3I_3 - A) = \begin{pmatrix} 0 & -6 & 8 \\ 0 & 3 & -6 \\ 0 & 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_3 = 3$,

$$(3I_3 - A) = \begin{pmatrix} 0 & -6 & 8 \\ 0 & 3 & -6 \\ 0 & 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (3I_n - A) \vec{x} = \vec{0}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_3 = 3$,

$$(3I_3 - A) = \begin{pmatrix} 0 & -6 & 8 \\ 0 & 3 & -6 \\ 0 & 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (3I_n - A) \vec{x} = \vec{0} \Rightarrow \begin{cases} y = 0 \\ z = 0 \end{cases}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_3 = 3$,

$$(3I_3 - A) = \begin{pmatrix} 0 & -6 & 8 \\ 0 & 3 & -6 \\ 0 & 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \Rightarrow (3I_n - A) \vec{x} = \vec{0} \Rightarrow \begin{cases} y = 0 \\ z = 0 \end{cases}$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \end{pmatrix} = t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, t \in \mathbb{R} \Rightarrow \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

Similar Matrices

Definition

The $n \times n$ matrices A and B are said to be **similar** if there is an invertible $n \times n$ matrix P such that $A = PBP^{-1}$.

Definition

The $n \times n$ matrices A and B are said to be **similar** if there is an invertible $n \times n$ matrix P such that $A = PBP^{-1}$.

Note: If $n \times n$ matrices are similar, then they have the same characteristic polynomial and hence the same eigenvalues (with the same multiplicities).

Similar Matrices

Definition

The $n \times n$ matrices A and B are said to be **similar** if there is an invertible $n \times n$ matrix P such that $A = PBP^{-1}$.

Note: If $n \times n$ matrices are similar, then they have the same characteristic polynomial and hence the same eigenvalues (with the same multiplicities).

Definition

A square matrix A is said to be **diagonalizable** if it is similar to a diagonal matrix. In other words, a diagonal matrix A has the property that there exists an invertible matrix P and a diagonal matrix D such that $A = PDP^{-1}$.

Similar Matrices

Definition

The $n \times n$ matrices A and B are said to be **similar** if there is an invertible $n \times n$ matrix P such that $A = PBP^{-1}$.

Note: If $n \times n$ matrices are similar, then they have the same characteristic polynomial and hence the same eigenvalues (with the same multiplicities).

Definition

A square matrix A is said to be **diagonalizable** if it is similar to a diagonal matrix. In other words, a diagonal matrix A has the property that there exists an invertible matrix P and a diagonal matrix D such that $A = PDP^{-1}$.

Note: $A = PDP^{-1} \Rightarrow A^k = PD^kP^{-1}$

Diagonalization

Theorem (The Diagonalization Theorem)

1. *An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.*

Diagonalization

Theorem (The Diagonalization Theorem)

1. An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.
2. If $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are linearly independent eigenvectors of A and $\lambda_1, \lambda_2, \dots, \lambda_n$ are their corresponding eigenvalues, then $A = PDP^{-1}$, where

$$P = \begin{pmatrix} \vec{v}_1 & \dots & \vec{v}_n \end{pmatrix} \text{ and } D = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}.$$

Diagonalization

Example

We have found the eigenvalues and corresponding eigenvectors of

the matrix $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$ as $\lambda_1 = 0, \lambda_2 = 2, \lambda_3 = 3$ and

$$\vec{v}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}, \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}. \text{ Then}$$

1. show that $\vec{v}_1, \vec{v}_2, \vec{v}_3$ are linear independent,

Diagonalization

Example

We have found the eigenvalues and corresponding eigenvectors of

the matrix $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$ as $\lambda_1 = 0, \lambda_2 = 2, \lambda_3 = 3$ and

$$\vec{v}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}, \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}. \text{ Then}$$

1. show that $\vec{v}_1, \vec{v}_2, \vec{v}_3$ are linear independent,
2. show that there exists matrices P and D such that $A = PDP^{-1}$,

Diagonalization

Example

We have found the eigenvalues and corresponding eigenvectors of

the matrix $A = \begin{pmatrix} 3 & 6 & -8 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{pmatrix}$ as $\lambda_1 = 0, \lambda_2 = 2, \lambda_3 = 3$ and

$$\vec{v}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} -10 \\ 3 \\ 1 \end{pmatrix}, \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}. \text{ Then}$$

1. show that $\vec{v}_1, \vec{v}_2, \vec{v}_3$ are linear independent,
2. show that there exists matrices P and D such that $A = PDP^{-1}$,
3. and find A^{125} .

Solution

$$1. P = \left(\begin{array}{ccc} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \end{array} \right) = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix} \text{ and since}$$
$$\det(P) = 1 \neq 0, \vec{v}_1, \vec{v}_2, \vec{v}_3 \text{ are linearly independent.}$$

Solution

$$1. P = \begin{pmatrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \end{pmatrix} = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix} \text{ and since}$$
$$\det(P) = 1 \neq 0, \vec{v}_1, \vec{v}_2, \vec{v}_3 \text{ are linearly independent.}$$

$$2. P = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix}, P^{-1} = \begin{pmatrix} 0 & 1 & -3 \\ 0 & 0 & 1 \\ 1 & 2 & 4 \end{pmatrix} \text{ and}$$

$$D = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} \Rightarrow A = PDP^{-1}.$$

Solution

$$1. P = \begin{pmatrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \end{pmatrix} = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix} \text{ and since}$$

$\det(P) = 1 \neq 0$, $\vec{v}_1, \vec{v}_2, \vec{v}_3$ are linearly independent.

$$2. P = \begin{pmatrix} -2 & -10 & 1 \\ 1 & 3 & 0 \\ 0 & 1 & 0 \end{pmatrix}, P^{-1} = \begin{pmatrix} 0 & 1 & -3 \\ 0 & 0 & 1 \\ 1 & 2 & 4 \end{pmatrix} \text{ and}$$

$$D = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} \Rightarrow A = PDP^{-1}.$$

$$3. A^{125} = PD^{125}P^{-1} = P \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2^{125} & 0 \\ 0 & 0 & 3^{125} \end{pmatrix} P^{-1}.$$

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem

Theorem (Cayley-Hamilton Theorem)

Every square matrix A is a root of its characteristic polynomial, i.e., $P_A(A) = O_{n \times n}$.

Cayley-Hamilton Theorem

Theorem (Cayley-Hamilton Theorem)

Every square matrix A is a root of its characteristic polynomial, i.e., $P_A(A) = O_{n \times n}$.

Note: The inverse or any power of a square matrix can be computed by using Cayley-Hamilton Theorem.

Cayley-Hamilton Theorem

Theorem (Cayley-Hamilton Theorem)

Every square matrix A is a root of its characteristic polynomial, i.e., $P_A(A) = O_{n \times n}$.

Note: The inverse or any power of a square matrix can be computed by using Cayley-Hamilton Theorem.

$$P_A(A) = A^n + a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n = 0$$

Cayley-Hamilton Theorem

Theorem (Cayley-Hamilton Theorem)

Every square matrix A is a root of its characteristic polynomial, i.e., $P_A(A) = O_{n \times n}$.

Note: The inverse or any power of a square matrix can be computed by using Cayley-Hamilton Theorem.

$$\begin{aligned}P_A(A) &= A^n + a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n = 0 \\ \Rightarrow A^n &= -\left(a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n\right)\end{aligned}$$

Cayley-Hamilton Theorem

Theorem (Cayley-Hamilton Theorem)

Every square matrix A is a root of its characteristic polynomial, i.e., $P_A(A) = O_{n \times n}$.

Note: The inverse or any power of a square matrix can be computed by using Cayley-Hamilton Theorem.

$$\begin{aligned}P_A(A) &= A^n + a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n = 0 \\ \Rightarrow A^n &= -\left(a_{n-1}A^{n-1} + \dots + a_1A + a_0I_n\right)\end{aligned}$$

If $a_0 \neq 0$, then

$$I_n = A \underbrace{\frac{-1}{a_0} \left(A^{n-1} + a_{n-1}A^{n-2} + \dots + a_1I_n \right)}_{A^{-1}}$$

Example

Example

Let $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$. Find

1. characteristic polynomial,

Example

Example

Let $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$. Find

1. characteristic polynomial,
2. eigenvalues,

Example

Example

Let $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$. Find

1. characteristic polynomial,
2. eigenvalues,
3. eigenvectors,

Example

Example

Let $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$. Find

1. characteristic polynomial,
2. eigenvalues,
3. eigenvectors,
4. matrices P and D such that $A = PDP^{-1}$, if any,

Example

Example

Let $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$. Find

1. characteristic polynomial,
2. eigenvalues,
3. eigenvectors,
4. matrices P and D such that $A = PDP^{-1}$, if any,
5. A^{-1} and A^5 (by using Cayley-Hamilton Theorem).

Finding Eigenvalues and Eigenvectors

Solution (1)

$$P_A(\lambda) = \det(\lambda I_n - A)$$

$$\lambda I_2 - A = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$$

Finding Eigenvalues and Eigenvectors

Solution (1)

$$P_A(\lambda) = \det(\lambda I_n - A)$$

$$\begin{aligned}\lambda I_2 - A &= \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \\ &= \begin{pmatrix} \lambda - 1 & -4 \\ -2 & \lambda - 3 \end{pmatrix}\end{aligned}$$

Finding Eigenvalues and Eigenvectors

Solution (1)

$$P_A(\lambda) = \det(\lambda I_n - A)$$

$$\begin{aligned}\lambda I_2 - A &= \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \\ &= \begin{pmatrix} \lambda - 1 & -4 \\ -2 & \lambda - 3 \end{pmatrix} \\ \Rightarrow P_A(\lambda) &= (\lambda - 1)(\lambda - 3) - 8 \\ &= \lambda^2 - 4\lambda - 5.\end{aligned}$$

Finding Eigenvalues and Eigenvectors

Solution (2)

The eigenvalues are the roots of $P_A(\lambda)$ such that

$$P_A(\lambda) = \lambda^2 - 4\lambda - 5 = 0$$

Finding Eigenvalues and Eigenvectors

Solution (2)

The eigenvalues are the roots of $P_A(\lambda)$ such that

$$P_A(\lambda) = \lambda^2 - 4\lambda - 5 = 0$$

$$\Rightarrow \lambda_1 = -1, \lambda_2 = 5.$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_1 = -1$,

$$(I_2 + A) = \begin{pmatrix} 2 & 4 \\ 2 & 4 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_1 = -1$,

$$(I_2 + A) = \begin{pmatrix} 2 & 4 \\ 2 & 4 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (I_2 + A) \vec{x} = \vec{0}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_1 = -1$,

$$(I_2 + A) = \begin{pmatrix} 2 & 4 \\ 2 & 4 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (I_2 + A) \vec{x} = \vec{0} \Rightarrow x + 2y = 0$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_1 = -1$,

$$(I_2 + A) = \begin{pmatrix} 2 & 4 \\ 2 & 4 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (I_2 + A) \vec{x} = \vec{0} \Rightarrow x + 2y = 0$$

$$\Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = t \begin{pmatrix} -2 \\ 1 \end{pmatrix}, \quad t \in \mathbb{R} \Rightarrow \vec{v}_1 = \begin{pmatrix} -2 \\ 1 \end{pmatrix}.$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_2 = 5$,

$$(5I_2 - A) = \begin{pmatrix} 4 & -4 \\ -2 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_2 = 5$,

$$(5I_2 - A) = \begin{pmatrix} 4 & -4 \\ -2 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$
$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (5I_2 - A) \vec{x} = \vec{0}$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_2 = 5$,

$$(5I_2 - A) = \begin{pmatrix} 4 & -4 \\ -2 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (5I_2 - A) \vec{x} = \vec{0} \Rightarrow x - y = 0$$

Finding Eigenvalues and Eigenvectors

Solution (3)

For $\lambda_2 = 5$,

$$(5I_2 - A) = \begin{pmatrix} 4 & -4 \\ -2 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$$

$$\vec{x} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow (5I_2 - A) \vec{x} = \vec{0} \Rightarrow x - y = 0$$

$$\Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = t \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad t \in \mathbb{R} \Rightarrow \vec{v}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

Finding Eigenvalues and Eigenvectors

Solution (4)

$$P = \begin{pmatrix} -2 & 1 \\ 1 & 1 \end{pmatrix}, P^{-1} = \begin{pmatrix} -\frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} \end{pmatrix} \text{ and } D = \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix} \Rightarrow$$
$$A = PDP^{-1}.$$

Cayley-Hamilton Theorem

Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

Cayley-Hamilton Theorem

Solution (5)

$$\begin{aligned}P_A(A) &= A^2 - 4A - 5I_2 = 0 \\ \Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} &= I_2\end{aligned}$$

Cayley-Hamilton Theorem

Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

$$\Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} = I_2$$

$$A^{-1} = \frac{1}{5} \left(\begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right) = \frac{1}{5} \begin{pmatrix} -3 & 4 \\ 2 & -1 \end{pmatrix}$$

Cayley-Hamilton Theorem

Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

$$\Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} = I_2$$

$$A^{-1} = \frac{1}{5} \left(\begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right) = \frac{1}{5} \begin{pmatrix} -3 & 4 \\ 2 & -1 \end{pmatrix}$$

$$A^2 = 4A + 5I_2$$

Cayley-Hamilton Theorem

Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

$$\Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} = I_2$$

$$A^{-1} = \frac{1}{5} \left(\begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right) = \frac{1}{5} \begin{pmatrix} -3 & 4 \\ 2 & -1 \end{pmatrix}$$

$$A^2 = 4A + 5I_2$$

$$A^4 = (4A + 5I_2)(4A + 5I_2)$$

Cayley-Hamilton Theorem

Solution (5)

$$P_A(A) = A^2 - 4A - 5I_2 = 0$$

$$\Rightarrow \underbrace{A \frac{1}{5} (A - 4I_2)}_{A^{-1}} = I_2$$

$$A^{-1} = \frac{1}{5} \left(\begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right) = \frac{1}{5} \begin{pmatrix} -3 & 4 \\ 2 & -1 \end{pmatrix}$$

$$A^2 = 4A + 5I_2$$

$$A^4 = (4A + 5I_2)(4A + 5I_2)$$

$$A^5 = A(4A + 5I_2)(4A + 5I_2)$$

?