

**DEFINITIONS** A **power series about  $x = 0$**  is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \cdots + c_n x^n + \cdots \quad (1)$$

A **power series about  $x = a$**  is a series of the form

$$\sum_{n=0}^{\infty} c_n (x - a)^n = c_0 + c_1 (x - a) + c_2 (x - a)^2 + \cdots + c_n (x - a)^n + \cdots \quad (2)$$

in which the **center**  $a$  and the **coefficients**  $c_0, c_1, c_2, \dots, c_n, \dots$  are constants.

## The Radius of Convergence of a Power Series

**3 Theorem** For a given power series  $\sum_{n=0}^{\infty} c_n (x - a)^n$  there are only three possibilities:

- (i) The series converges only when  $x = a$ .  $\Rightarrow R = 0$
- (ii) The series converges for all  $x$ .  $\Rightarrow R = \infty$
- (iii) There is a positive number  $R$  such that the series converges if  $|x - a| < R$  and diverges if  $|x - a| > R$ .

endpoints. Thus in case (iii) there are four possibilities for the interval of convergence:

$$(a - R, a + R) \quad (a - R, a + R] \quad [a - R, a + R) \quad [a - R, a + R]$$

### How to Test a Power Series for Convergence

1. Use the *Ratio Test* (or *Root Test*) to find the interval where the series converges absolutely. Ordinarily, this is an open interval

$$|x - a| < R \quad \text{or} \quad a - R < x < a + R.$$

2. If the interval of absolute convergence is finite, test for convergence or divergence at each endpoint, as in Examples 3a and b. Use a Comparison Test, the Integral Test, or the Alternating Series Test.
3. If the interval of absolute convergence is  $a - R < x < a + R$ , the series diverges for  $|x - a| > R$  (it does not even converge conditionally) because the  $n$ th term does not approach zero for those values of  $x$ .

## Operations on Power Series

**THEOREM 19—The Series Multiplication Theorem for Power Series** If

$A(x) = \sum_{n=0}^{\infty} a_n x^n$  and  $B(x) = \sum_{n=0}^{\infty} b_n x^n$  converge absolutely for  $|x| < R$ , and

$$c_n = a_0 b_n + a_1 b_{n-1} + a_2 b_{n-2} + \cdots + a_{n-1} b_1 + a_n b_0 = \sum_{k=0}^n a_k b_{n-k},$$

then  $\sum_{n=0}^{\infty} c_n x^n$  converges absolutely to  $A(x)B(x)$  for  $|x| < R$ :

$$\left( \sum_{n=0}^{\infty} a_n x^n \right) \cdot \left( \sum_{n=0}^{\infty} b_n x^n \right) = \sum_{n=0}^{\infty} c_n x^n.$$

**THEOREM 20** If  $\sum_{n=0}^{\infty} a_n x^n$  converges absolutely for  $|x| < R$ , then  $\sum_{n=0}^{\infty} a_n (f(x))^n$  converges absolutely for any continuous function  $f$  on  $|f(x)| < R$ .

## Differentiation and Integration of Power Series

**2 Theorem** If the power series  $\sum c_n (x - a)^n$  has radius of convergence  $R > 0$ , then the function  $f$  defined by

$$f(x) = c_0 + c_1(x - a) + c_2(x - a)^2 + \cdots = \sum_{n=0}^{\infty} c_n (x - a)^n$$

is differentiable (and therefore continuous) on the interval  $(a - R, a + R)$  and

$$(i) \quad f'(x) = c_1 + 2c_2(x - a) + 3c_3(x - a)^2 + \cdots = \sum_{n=1}^{\infty} n c_n (x - a)^{n-1}$$

$$(ii) \quad \int f(x) dx = C + c_0(x - a) + c_1 \frac{(x - a)^2}{2} + c_2 \frac{(x - a)^3}{3} + \cdots \\ = C + \sum_{n=0}^{\infty} c_n \frac{(x - a)^{n+1}}{n + 1}$$

The radii of convergence of the power series in Equations (i) and (ii) are both  $R$ .

This is called **term-by-term differentiation and integration**.