



MAT1320-Linear Algebra

Lecture Notes

System of Linear Equations, Gauss-Jordan Elimination Method

Mehmet E. KÖROĞLU
Fall 2024

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS
mkoroglu@yildiz.edu.tr

Table of contents

1. System of Linear Equations
2. The Rank of A Matrix
3. Gauss-Jordan Elimination Method
4. Homogenous System of Linear Equations

System of Linear Equations

System of Linear Equations

$$\left\{ \begin{array}{l} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{array} \right. \quad (1)$$

is a **system of linear equations** of m equations in n unknowns.

System of Linear Equations

$$\left\{ \begin{array}{l} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{array} \right. \quad (1)$$

is a **system of linear equations** of m equations in n unknowns. The matrix

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

is called the **coefficient matrix** of the system.

System of Linear Equations

Then for $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}$ the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}$$

System of Linear Equations

Then for $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}$ the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}$$

where \mathbf{X} is called the **vector of unknowns** and \mathbf{b} is called the **vector of constants**.

System of Linear Equations

Then for $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}$ the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}$$

where \mathbf{X} is called the **vector of unknowns** and \mathbf{b} is called the **vector of constants**.

Definition

The **augmented matrix** of the system is denoted by $[A|\mathbf{b}]$ and obtained by appending the vector of constants \mathbf{b} as the $(n+1)$ -th column to the coefficient matrix A .

The Rank of A Matrix

The Rank of A Matrix

Definition

Let REF or RREF of the matrix A be the C . The number of nonzero rows of C is called **rank** of A and denoted by $\text{rank}(A)$.

The Rank of A Matrix

Definition

Let REF or RREF of the matrix A be the C . The number of nonzero rows of C is called **rank** of A and denoted by $\text{rank}(A)$.

Example

The matrix A and its RREF C is given below.

$$A = \begin{pmatrix} 1 & 0 & 1 & 0 & 1 \\ 2 & 0 & 2 & 0 & 2 \\ 1 & -1 & 0 & 1 & 0 \\ 2 & -2 & 0 & 2 & 0 \end{pmatrix} \sim C = \begin{pmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Then $\text{rank}(A) = 2$.

Gauss-Jordan Elimination Method

Gauss-Jordan Elimination Method

Let $A\mathbf{X} = \mathbf{b}$ be the system of linear equation with m equations in n unknowns given in (1).

Gauss-Jordan Elimination Method

Let $A\mathbf{X} = \mathbf{b}$ be the system of linear equation with m equations in n unknowns given in (1). Let $[A|\mathbf{b}]$ be the augmented matrix of the system.

Gauss-Jordan Elimination Method

Let $A\mathbf{X} = \mathbf{b}$ be the system of linear equation with m equations in n unknowns given in (1). Let $[A|\mathbf{b}]$ be the augmented matrix of the system. Examination of the solutions of a system of linear equations by transforming augmented matrix of the system $[A|\mathbf{b}]$ into RREF is called **Gauss-Jordan Elimination Method**.

Gauss-Jordan Elimination Method

Let $A\mathbf{X} = \mathbf{b}$ be the system of linear equation with m equations in n unknowns given in (1). Let $[A|\mathbf{b}]$ be the augmented matrix of the system. Examination of the solutions of a system of linear equations by transforming augmented matrix of the system $[A|\mathbf{b}]$ into RREF is called **Gauss-Jordan Elimination Method**. In the Gauss-Jordan Elimination Method we have the following cases:

1. If $\text{rank}(A) \neq \text{rank}([A|\mathbf{b}])$, then the system has no solution. In this case we called system as **inconsistent**.

Gauss-Jordan Elimination Method

Let $A\mathbf{X} = \mathbf{b}$ be the system of linear equation with m equations in n unknowns given in (1). Let $[A|\mathbf{b}]$ be the augmented matrix of the system. Examination of the solutions of a system of linear equations by transforming augmented matrix of the system $[A|\mathbf{b}]$ into RREF is called **Gauss-Jordan Elimination Method**. In the Gauss-Jordan Elimination Method we have the following cases:

1. If $\text{rank}(A) \neq \text{rank}([A|\mathbf{b}])$, then the system has no solution. In this case we called system as **inconsistent**.
2. If $\text{rank}(A) = \text{rank}([A|\mathbf{b}])$, then the system has solution.

Gauss-Jordan Elimination Method

Let $A\mathbf{X} = \mathbf{b}$ be the system of linear equation with m equations in n unknowns given in (1). Let $[A|\mathbf{b}]$ be the augmented matrix of the system. Examination of the solutions of a system of linear equations by transforming augmented matrix of the system $[A|\mathbf{b}]$ into RREF is called **Gauss-Jordan Elimination Method**. In the Gauss-Jordan Elimination Method we have the following cases:

1. If $\text{rank}(A) \neq \text{rank}([A|\mathbf{b}])$, then the system has no solution. In this case we called system as **inconsistent**.
2. If $\text{rank}(A) = \text{rank}([A|\mathbf{b}])$, then the system has solution.
 - 2.1 If $\text{rank}(A) = n$, i.e., the number of unknowns and the number of equations are equal, then the system has a unique solution.

Gauss-Jordan Elimination Method

Let $A\mathbf{X} = \mathbf{b}$ be the system of linear equation with m equations in n unknowns given in (1). Let $[A|\mathbf{b}]$ be the augmented matrix of the system. Examination of the solutions of a system of linear equations by transforming augmented matrix of the system $[A|\mathbf{b}]$ into RREF is called **Gauss-Jordan Elimination Method**. In the Gauss-Jordan Elimination Method we have the following cases:

1. If $\text{rank}(A) \neq \text{rank}([A|\mathbf{b}])$, then the system has no solution. In this case we called system as **inconsistent**.
2. If $\text{rank}(A) = \text{rank}([A|\mathbf{b}])$, then the system has solution.
 - 2.1 If $\text{rank}(A) = n$, i.e., the number of unknowns and the number of equations are equal, then the system has a unique solution.
 - 2.2 If $\text{rank}(A) = r < n$, i.e., the number of unknowns is greater than the number of equations, then the system has infinitely many solutions depending on $n - r$ parameters.

Gauss-Jordan Elimination Method

Example (Gauss-Jordan Elimination)

Let the system of linear equations $x - y + 3z = 1$, $y = -2x + 5$, $9z - x - 5y + 7 = 0$ be given.

Gauss-Jordan Elimination Method

Example (Gauss-Jordan Elimination)

Let the system of linear equations $x - y + 3z = 1$, $y = -2x + 5$, $9z - x - 5y + 7 = 0$ be given.

1. For the vector of unknowns $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, state the system as $\mathbf{Ax} = \mathbf{b}$.

Gauss-Jordan Elimination Method

Example (Gauss-Jordan Elimination)

Let the system of linear equations $x - y + 3z = 1$, $y = -2x + 5$, $9z - x - 5y + 7 = 0$ be given.

1. For the vector of unknowns $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, state the system as

$$\mathbf{Ax} = \mathbf{b}.$$

2. Write the augmented matrix of the given system and put it into RREF.

Gauss-Jordan Elimination Method

Example (Gauss-Jordan Elimination)

Let the system of linear equations $x - y + 3z = 1$, $y = -2x + 5$, $9z - x - 5y + 7 = 0$ be given.

1. For the vector of unknowns $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, state the system as

$$\mathbf{Ax} = \mathbf{b}.$$

2. Write the augmented matrix of the given system and put it into RREF.
3. Find $\text{rank}(\mathbf{A})$ and $\text{rank}([\mathbf{A}|\mathbf{b}])$.

Gauss-Jordan Elimination Method

Example (Gauss-Jordan Elimination)

Let the system of linear equations $x - y + 3z = 1$, $y = -2x + 5$, $9z - x - 5y + 7 = 0$ be given.

1. For the vector of unknowns $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, state the system as

$$\mathbf{Ax} = \mathbf{b}.$$

2. Write the augmented matrix of the given system and put it into RREF.
3. Find $\text{rank}(\mathbf{A})$ and $\text{rank}([\mathbf{A}|\mathbf{b}])$.
4. If any, find all solutions of the system.

1. The given system can be arranged as follows.

$$\begin{array}{rclcl} x & -y & +3z & = & 1 \\ 2x & +y & & = & 5 \\ -x & -5y & +9z & = & -7 \end{array}$$

Çözüm:1

1. The given system can be arranged as follows.

$$\begin{array}{rrcr} x & -y & +3z & = & 1 \\ 2x & +y & & = & 5 \\ -x & -5y & +9z & = & -7 \end{array}$$

Then

$$\begin{bmatrix} 1 & -1 & 3 \\ 2 & 1 & 0 \\ -1 & -5 & 9 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \\ -7 \end{bmatrix}$$

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right]$.

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array}$$

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 3 & -6 & 3 \\ 0 & -6 & 12 & -6 \end{array} \right] \begin{array}{l} H_2\left(\frac{1}{3}\right) \\ H_3\left(\frac{-1}{6}\right) \\ \sim \end{array}$$

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 3 & -6 & 3 \\ 0 & -6 & 12 & -6 \end{array} \right] \begin{array}{l} H_2\left(\frac{1}{3}\right) \\ H_3\left(\frac{-1}{6}\right) \\ \sim \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 1 & -2 & 1 \\ 0 & 1 & -2 & 1 \end{array} \right] \begin{array}{l} H_{32}(-1) \\ \sim \end{array}$$

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 3 & -6 & 3 \\ 0 & -6 & 12 & -6 \end{array} \right] \begin{array}{l} H_2\left(\frac{1}{3}\right) \\ H_3\left(\frac{-1}{6}\right) \\ \sim \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 1 & -2 & 1 \\ 0 & 1 & -2 & 1 \end{array} \right] \begin{array}{l} H_{32}(-1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{l} H_{12}(1) \\ \sim \end{array}$$

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 2 & 1 & 0 & 5 \\ -1 & -5 & 9 & -7 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 3 & -6 & 3 \\ 0 & -6 & 12 & -6 \end{array} \right] \begin{array}{l} H_2\left(\frac{1}{3}\right) \\ H_3\left(\frac{-1}{6}\right) \\ \sim \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 1 & -2 & 1 \\ 0 & 1 & -2 & 1 \end{array} \right] \begin{array}{l} H_{32}(-1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 1 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{l} H_{12}(1) \\ \sim \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 1 & 2 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

3. $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{b}]) = 2.$

Çözüm:

3. $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{b}]) = 2.$

4. Since $[\mathbf{A}|\mathbf{b}] \sim \left[\begin{array}{ccc|c} 1 & 0 & 1 & 2 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$

Çözüm:

3. $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{b}]) = 2.$

4. Since $[\mathbf{A}|\mathbf{b}] \sim \left[\begin{array}{ccc|c} 1 & 0 & 1 & 2 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$ we have
$$\begin{aligned} x &= 2 - z \\ y &= 1 + 2z \end{aligned}.$$

Çözüm:

3. $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{b}]) = 2.$

4. Since $[\mathbf{A}|\mathbf{b}] \sim \left[\begin{array}{ccc|c} 1 & 0 & 1 & 2 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$ we have $\begin{matrix} x = 2 - z \\ y = 1 + 2z \end{matrix}$. Then

for arbitrary parameter $z = t \in \mathbb{R}$ we have $\begin{matrix} x = 2 - t \\ y = 1 + 2t \\ z = t \end{matrix}$. Then

$$\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ 2 \\ 1 \end{bmatrix} \text{ is the solution of the system.}$$

Homogenous System of Linear Equations

Homogenous System of Linear Equations

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0 \end{cases} \quad (2)$$

is a **homogeneous linear system** of m equations in n unknowns.

Homogenous System of Linear Equations

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0 \end{cases} \quad (2)$$

is a **homogeneous linear system** of m equations in n unknowns.

A homogenous system always has solution, because $\text{rank}(A) = \text{rank}([A|\mathbf{b}])$.

Homogenous System of Linear Equations

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0 \end{cases} \quad (2)$$

is a **homogeneous linear system** of m equations in n unknowns.

A homogenous system always has solution, because $\text{rank}(A) = \text{rank}([A|\mathbf{b}])$.

1. If $\text{rank}(A) = n$, i.e., the number of unknowns and the number of equations are equal, then the system has zero solution as the unique solution.

Homogenous System of Linear Equations

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0 \end{cases} \quad (2)$$

is a **homogeneous linear system** of m equations in n unknowns.

A homogenous system always has solution, because $\text{rank}(A) = \text{rank}([A|\mathbf{b}])$.

1. If $\text{rank}(A) = n$, i.e., the number of unknowns and the number of equations are equal, then the system has zero solution as the unique solution.
2. If $\text{rank}(A) = r < n$, i.e., the number of unknowns is greater than the number of equations, then the system has infinitely

Mehmet E. KÖROĞLU many solutions depending on $n - r$ parameters.

Homogenous System of Linear Equations

Example

Let the homogenous system $\begin{cases} x - y + 3z = 0 \\ 2x + y = 0 \\ -x - 5y + 9z = 0 \end{cases}$ be given.

Homogenous System of Linear Equations

Example

Let the homogenous system $\begin{cases} x - y + 3z = 0 \\ 2x + y = 0 \\ -x - 5y + 9z = 0 \end{cases}$ be given.

1. For the vector of unknowns $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, state the system as

$$\mathbf{Ax} = \mathbf{b}.$$

Homogenous System of Linear Equations

Example

Let the homogenous system $\begin{cases} x - y + 3z = 0 \\ 2x + y = 0 \\ -x - 5y + 9z = 0 \end{cases}$ be given.

1. For the vector of unknowns $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, state the system as

$$\mathbf{Ax} = \mathbf{b}.$$

2. Write the augmented matrix of the given system and put it into RREF.

Homogenous System of Linear Equations

Example

Let the homogenous system $\begin{cases} x - y + 3z = 0 \\ 2x + y = 0 \\ -x - 5y + 9z = 0 \end{cases}$ be given.

1. For the vector of unknowns $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, state the system as

$$\mathbf{Ax} = \mathbf{b}.$$

2. Write the augmented matrix of the given system and put it into RREF.
3. Find $\text{rank}(\mathbf{A})$ and $\text{rank}([\mathbf{A}|\mathbf{b}])$.

Homogenous System of Linear Equations

Example

Let the homogenous system $\begin{cases} x - y + 3z = 0 \\ 2x + y = 0 \\ -x - 5y + 9z = 0 \end{cases}$ be given.

1. For the vector of unknowns $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, state the system as $\mathbf{Ax} = \mathbf{b}$.
2. Write the augmented matrix of the given system and put it into RREF.
3. Find $\text{rank}(\mathbf{A})$ and $\text{rank}([\mathbf{A}|\mathbf{b}])$.
4. Find all the solutions of the system.

1. We rearrange the system as follows:

$$\begin{array}{rrcr} x & -y & +3z & = 0 \\ 2x & +y & & = 0 \\ -x & -5y & +9z & = 0 \end{array}$$

1. We rearrange the system as follows:

$$\begin{array}{rrcr} x & -y & +3z & = 0 \\ 2x & +y & & = 0 \\ -x & -5y & +9z & = 0 \end{array}$$

Then

$$\begin{bmatrix} 1 & -1 & 3 \\ 2 & 1 & 0 \\ -1 & -5 & 9 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right]$.

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array}$$

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 3 & -6 & 0 \\ 0 & -6 & 12 & 0 \end{array} \right] \begin{array}{l} H_2\left(\frac{1}{3}\right) \\ H_3\left(\frac{-1}{6}\right) \\ \sim \end{array}$$

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 3 & -6 & 0 \\ 0 & -6 & 12 & 0 \end{array} \right] \begin{array}{l} H_2\left(\frac{1}{3}\right) \\ H_3\left(\frac{-1}{6}\right) \\ \sim \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 1 & -2 & 0 \end{array} \right] \begin{array}{l} H_{32}(-1) \\ \sim \end{array}$$

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 3 & -6 & 0 \\ 0 & -6 & 12 & 0 \end{array} \right] \begin{array}{l} H_2\left(\frac{1}{3}\right) \\ H_3\left(\frac{-1}{6}\right) \\ \sim \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 1 & -2 & 0 \end{array} \right] \begin{array}{l} H_{32}(-1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{l} H_{12}(1) \\ \sim \end{array}$$

Çözüm:2

2. The augmented matrix of the system is $\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right]$.

Apply elementary row operations to the augmented matrix to get REF and RREF.

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 2 & 1 & 0 & 0 \\ -1 & -5 & 9 & 0 \end{array} \right] \begin{array}{l} H_{21}(-2) \\ H_{31}(1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 3 & -6 & 0 \\ 0 & -6 & 12 & 0 \end{array} \right] \begin{array}{l} H_2\left(\frac{1}{3}\right) \\ H_3\left(\frac{-1}{6}\right) \\ \sim \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 1 & -2 & 0 \end{array} \right] \begin{array}{l} H_{32}(-1) \\ \sim \end{array} \left[\begin{array}{ccc|c} 1 & -1 & 3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{l} H_{12}(1) \\ \sim \end{array}$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

3. $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{b}]) = 2.$

Çözüm:

3. $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{b}]) = 2.$

4. Since $[\mathbf{A}|\mathbf{b}] \sim \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$ we have

Çözüm:

3. $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{b}]) = 2.$

4. Since $[\mathbf{A}|\mathbf{b}] \sim \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$ we have
$$\begin{aligned} x &= -z \\ y &= 2z \end{aligned}$$

Çözüm:

3. $\text{rank}(\mathbf{A}) = \text{rank}([\mathbf{A}|\mathbf{b}]) = 2.$

4. Since $[\mathbf{A}|\mathbf{b}] \sim \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$ we have $\begin{array}{l} x = -z \\ y = 2z \end{array}$ Then for

arbitrary parameter $z = t \in \mathbb{R}$ we get $\begin{array}{l} x = -t \\ y = 2t \end{array}$. Thus, all
 $z = t$

solutions are of the form $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} = t \begin{bmatrix} -1 \\ 2 \\ 1 \end{bmatrix}.$

?