

# Asymptotic center–manifold for systems with parameter perturbations

Prabal S. Negi

Okinawa Institute of Science and Technology Graduate University, Onna, Okinawa  
904-0495, Japan.

Corresponding author(s). E-mail(s): [prabal.negi@oist.jp](mailto:prabal.negi@oist.jp);

## Abstract

Asymptotic center-manifold approximations are constructed for systems that have been perturbed slightly away from the bifurcation point in the parameter space. This is done by extending the existing system to include parameter perturbations as dynamic variables, leading to an extended critical subspace and dimensionality of the center-manifold. The extension is done in a very general setting and it is shown that the consequences of such a system extension are non-trivial. The extended critical subspace has a certain structure associated with it, which may be deduced from the original system. The application of the center-manifold theorem to the extended system leads to a graph equation which rigorously valid in the vicinity of the bifurcation point (even in parameter space). The graph equation is approximated as an asymptotic power series in the amplitudes of the critical eigenmodes leading to general expressions for restricted resolvent solutions for the various terms in the power series at each order. A Ginzburg-Landau model near bifurcation is used to demonstrate the validity of the asymptotic center-manifold in the extended space.

## 1 Problem Definition

### 1.1 The Standard System

Consider a system with its state variable denoted by  $\mathbf{U}$ , consisting of  $l$  parameters given by the vector  $\boldsymbol{\mu}$ , and an evolution law given by

$$\frac{\partial \mathbf{U}}{\partial t} = \mathbf{f}(\mathbf{U}, \boldsymbol{\mu}). \quad (1)$$

The system state  $\mathbf{U}$  is  $N$ -dimensional where,  $N$  is assumed to be large. The evolution law may arise from a finite set of ordinary differential equations or, may be a finite representation of an infinite dimensional partial differential equation as often arises from the numerical discretization

of the space-time evolution equations of physical systems. Typically, physical systems will have real parameters  $\boldsymbol{\mu}$  however, allowing for complex parameters presents no additional difficulty and we assume that  $\boldsymbol{\mu} \in \mathbb{C}^l$ . For a certain set of parameter values  $\boldsymbol{\mu} = \boldsymbol{\mu}^b$ , the system is assumed to have a fixed point for a state vector  $\mathbf{U}^b$  satisfying

$$0 = \mathbf{f}(\mathbf{U}^b, \boldsymbol{\mu}^b). \quad (2)$$

The system may be rewritten in terms of the deviation from the fixed point solution  $\mathbf{U} = \mathbf{U}^b + \mathbf{u}$  such that

$$\begin{aligned} \frac{\partial \mathbf{u}}{\partial t} = & \mathbf{L}\mathbf{u} + \\ & [\mathbf{f}(\mathbf{U}^b + \mathbf{u}, \boldsymbol{\mu}^b) - \mathbf{L}\mathbf{u}], \end{aligned} \quad (3)$$

where,  $\mathbf{L}$  is the linearized operator at the fixed point, acting on  $\mathbf{u}$ , defined by

$$\mathbf{L}\mathbf{u} = \left( \frac{\partial \mathbf{f}}{\partial \mathbf{U}} \bigg|_{(\mathbf{U}^b, \boldsymbol{\mu}^b)} \right) \cdot \mathbf{u}. \quad (4)$$

The right hand side in equation (3) has been divided into the linear and non-linear contributions with  $\mathbf{L}(\mathbf{u})$  representing the linear part and the remaining terms in the square brackets representing the non-linear contributions.

Furthermore, we assume that  $\boldsymbol{\mu}^b$  represents the parameters of the system at a bifurcation point. It is generically assumed that the system undergoes a multiple co-dimension bifurcation at  $\boldsymbol{\mu}^b$  and that the linearized operator  $\mathbf{L}$  has a discrete set of  $n$  critical eigenvalues  $\lambda^C$ , with zero growth rate ( $\Re(\lambda^C) = 0$ ). This includes both purely real critical eigenvalues with  $\Re(\lambda^C) = 0$ ,  $\Im(\lambda^C) = 0$ , and purely imaginary critical eigenvalues with  $\Re(\lambda^C) = 0$ ,  $\Im(\lambda^C) \neq 0$ . For real systems complex eigenvalues occur in conjugate pairs. Accordingly, the system has  $n$  critical eigenpairs  $(\lambda_i^C, \mathbf{v}_i^C)$ ,  $i \in 1 \dots n$ , where  $\mathbf{v}_i^C$  represents the eigenvector corresponding to  $\lambda_i^C$ .

The adjoint of the direct linearized operator  $\mathbf{L}$  is denoted as  $\mathbf{L}^\dagger$  which also has  $n$  critical eigenpairs  $(\lambda_i^{C*}, \mathbf{w}_i^C)$ . The scaling of the eigenvectors of the direct and adjoint operators may be defined such that they satisfy the biorthogonality condition

$$\langle \mathbf{w}_i^C, \mathbf{v}_j^C \rangle = \mathbf{w}_i^{C\dagger} \mathbf{v}_j^C = \delta_{i,j}, \quad (5)$$

where  $\langle \cdot, \cdot \rangle$  is the standard inner-product and  $\delta_{i,j}$  is the Kronecker-delta function. The superscript  $^\dagger$  is used here to indicate a complex-conjugated transpose of vectors and matrix operators. For scalars complex-conjugation is indicated by a  $*$  superscript. The superscript  $^H$  which is often used for the same purpose is reserved for objects referring to harmonic forcings. One may define a matrix  $\mathbf{V}$ , whose columns comprise of the  $n$  critical eigenvectors of  $\mathbf{L}$  and the matrix spans the invariant center subspace of the linear operator  $\mathbf{L}$ . The corresponding matrix for the adjoint eigenvectors is denoted  $\mathbf{W}$ . Due to the chosen scaling denoted by equation (5) the matrices satisfy,

$$\mathbf{W}^\dagger \mathbf{V} = \mathbf{I}_n, \quad (6)$$

where,  $\mathbf{I}_n$  represents an identity matrix of order  $n \times n$ . The projection operator into the critical subspace (also the center subspace) is defined by  $\mathbf{P} = \mathbf{V}\mathbf{W}^\dagger$ , while the projection operator into its complement (stable subspace of  $\mathbf{L}$ ) is defined by,  $\mathbf{Q} = \mathbf{I} - \mathbf{V}\mathbf{W}^\dagger$ .

Additionally,  $\boldsymbol{\Lambda}_C$  is defined as an  $n \times n$  diagonal matrix consisting of the critical eigenvalues of  $\mathbf{L}$ , with the implicit assumption of diagonalizability of  $\mathbf{L}$  within this subspace. The ordering of the eigenvalues in  $\boldsymbol{\Lambda}_C$  and eigenvectors in  $\mathbf{V}$  is consistent with each other such that  $\mathbf{L}\mathbf{V} = \mathbf{V}\boldsymbol{\Lambda}_C$ .

## 1.2 The Extended System

The same system is now considered for the case of parameter perturbation so that the evolution dynamics are now evaluated at parameter values of  $\boldsymbol{\mu} = \boldsymbol{\mu}^b + \boldsymbol{\nu}$ , with  $\boldsymbol{\nu}$  being the vector of parameter perturbations,  $\boldsymbol{\nu} = [\nu_1, \nu_2, \dots, \nu_l]$ . In addition, the system is forced with small-amplitude harmonic forcing of the type

$$\mathbf{f}_H(t) = \sum_{i=1}^h \mathbf{f}_i \theta_i(t); \quad \theta_i = \epsilon_i e^{\lambda_i^H t}, \quad (7)$$

where, the various  $\theta_i$  are treated as the dynamic variable,  $\lambda_i^H$  is purely imaginary, with the imaginary part representing the angular frequency of the forcing.  $\mathbf{f}_i$  represents the shape of the  $i^{th}$  harmonic forcing and  $\epsilon_i$  is a scale factor. Constant forcing terms are readily incorporated when a particular (or multiple)  $\lambda_i^H = 0$ . The vector of the various  $\theta_i$  is denoted as  $\boldsymbol{\theta} = [\theta_1; \theta_2; \dots; \theta_h]$ , and the vector of angular frequencies is denoted as  $\boldsymbol{\lambda}^H = [\lambda_1^H; \lambda_2^H; \dots; \lambda_h^H]$ . The resulting evolution equation is

$$\frac{\partial \mathbf{u}}{\partial t} = \mathbf{f}(\mathbf{U}^b + \mathbf{u}, \boldsymbol{\mu}^b + \boldsymbol{\nu}) + \sum_{i=1}^h \mathbf{f}_i \theta_i.$$

The diagonal matrix for the harmonic angular frequencies is represented by  $\boldsymbol{\Lambda}_H = \text{diagm}(\boldsymbol{\lambda}^H)$ . Similarly,  $\boldsymbol{\lambda}^P = [\lambda_1^P; \lambda_2^P; \dots; \lambda_l^P]$  is defined as a vector of length  $l$ , with all  $\lambda_i^P = 0$  identically, for  $i \in 1 \dots l$ . These may be thought of parameter eigenvalues of parameter angular frequencies depending on the point of view. In either case, these vanish identically.  $\boldsymbol{\Lambda}_P = \text{diagm}(\boldsymbol{\lambda}^P)$  is defined as an  $l \times l$  null matrix, to represent

the trivial linear evolution matrix for the parameter variables  $\boldsymbol{\nu}$ . The extended system including the evolution equations for the newly introduced variables  $\boldsymbol{\nu}$  and  $\boldsymbol{\theta}$ , henceforth referred to as the *extended variables*, is given by,

$$\frac{\partial \mathbf{u}}{\partial t} = \mathbf{L}\mathbf{u} + \mathbf{L}_\mu \boldsymbol{\nu} + \mathbf{L}_\theta \boldsymbol{\theta} + \mathbf{f}_{NL}(\mathbf{u}, \boldsymbol{\nu}) \quad (8a)$$

$$\frac{d\boldsymbol{\nu}}{dt} = \boldsymbol{\Lambda}_P \boldsymbol{\nu} \quad (8b)$$

$$\frac{d\boldsymbol{\theta}}{dt} = \boldsymbol{\Lambda}_H \boldsymbol{\theta}, \quad (8c)$$

Here, as done earlier, the evolution equations are split into the linear and non-linear contributions.  $\mathbf{L}$  is the same linearized operator as defined earlier in equation 4.  $\mathbf{L}_\mu$  is the linearized operator acting on the newly introduced extended variable  $\boldsymbol{\nu}$ , defined as

$$\mathbf{L}_\mu \boldsymbol{\nu} = \left( \frac{\partial \mathbf{f}}{\partial \boldsymbol{\mu}} \Big|_{(U^b, \boldsymbol{\mu}^b)} \right) \cdot \boldsymbol{\nu}. \quad (9)$$

Similarly,  $\mathbf{L}_\theta$  is the linearized operator acting on  $\boldsymbol{\theta}$ , defined as,

$$\mathbf{L}_\theta \boldsymbol{\theta} = \left( \frac{\partial \mathbf{f}_H}{\partial \boldsymbol{\theta}} \right) \cdot \boldsymbol{\theta}. \quad (10)$$

This is the same expression as in equation (7), just written in a manner consistent with the other terms. Finally,  $\mathbf{f}_{NL}(\mathbf{u}, \boldsymbol{\nu})$  represents the non-linear terms of the modified system given by

$$\begin{aligned} \mathbf{f}_{NL}(\mathbf{u}, \boldsymbol{\nu}) = & \mathbf{f}(U^b + \mathbf{u}, \boldsymbol{\mu}^b + \boldsymbol{\nu}) \\ & - \mathbf{L}\mathbf{u} - \mathbf{L}_\mu \boldsymbol{\nu}, \end{aligned} \quad (11)$$

where, the dependence on  $U^b$  and  $\boldsymbol{\mu}^b$  has been suppressed for notational convenience.

An overhead hat notation,  $\hat{\cdot}$ , is used to refer to variables and operators associated with the extended space, so that the new state vector, non-linear term and the linearized direct and adjoint operators are denoted by  $\hat{\mathbf{u}}$ ,  $\hat{\mathbf{f}}_{NL}$ ,  $\hat{\mathcal{L}}$  and  $\hat{\mathcal{L}}^\dagger$ , that are defined as,

$$\begin{aligned} \hat{\mathbf{u}} &= [\mathbf{u}; \boldsymbol{\nu}; \boldsymbol{\theta}], \\ \hat{\mathbf{f}}_{NL} &= [\mathbf{f}_{NL}; \mathbf{0}; \mathbf{0}], \end{aligned}$$

$$\hat{\mathbf{L}} = \begin{bmatrix} \mathbf{L} & \mathbf{L}_\mu & \mathbf{L}_\theta \\ \mathbf{0} & \boldsymbol{\Lambda}_P & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \boldsymbol{\Lambda}_H \end{bmatrix}, \quad \hat{\mathbf{L}}^\dagger = \begin{bmatrix} \mathbf{L}^\dagger & \mathbf{0} & \mathbf{0} \\ \mathbf{L}_\mu^\dagger & \boldsymbol{\Lambda}_P^\dagger & \mathbf{0} \\ \mathbf{L}_\theta^\dagger & \mathbf{0} & \boldsymbol{\Lambda}_H^\dagger \end{bmatrix}, \quad (12)$$

where, the superscript  $T$  refers to vector transpose. The full system may then be represented compactly as

$$\frac{\partial \hat{\mathbf{u}}}{\partial t} = \hat{\mathbf{L}}\hat{\mathbf{u}} + \hat{\mathbf{f}}_{NL}. \quad (13)$$

The system defined by equation (8), or its compact form equation (13), with additional dynamical equations for the extended variables will be referred to as the *extended system*.

### 1.3 Extended Subspace Structure

The extended system has a larger critical subspace of dimension  $m = n + l + h$ , as compared to the original system due to the  $l + h$  additional variables introduced. While the extension of the system seems almost trivial, the eigenvector structure in the additional critical dimensions however is far from trivial. Nevertheless, the extended eigenspace structure can be deduced directly from the properties of the original system. It turns out to be incredibly useful to analyze the structure of the extended subspace since the the adjoint extended critical subspace does have a special structure which allows the asymptotic terms of the center-manifold to be evaluated using the original linear system  $\mathbf{L}$ .

There are however some subtleties involved when the system extension introduces new eigenvalues that are resonant with the original linear system  $\mathbf{L}$ . In such a scenario the system extension leads to a generalized eigenspace for the resonant eigenvalues. The subspace structure is now evaluated in detail for both the resonant and non-resonant cases. This is first done for the extended operator  $\hat{\mathbf{L}}$  and then for the adjoint operator  $\hat{\mathbf{L}}^\dagger$ .

#### 1.3.1 Subspace structure for $\hat{\mathbf{L}}$

Observe that if  $(\lambda_i^C, \mathbf{v}_i^C)$  is a critical eigenpair of the original linearized operator  $\mathbf{L}$ , then its extension to  $\hat{\mathbf{L}}$  satisfies,

$$(\lambda_i^C \mathbf{I} - \hat{\mathbf{L}})\hat{\mathbf{v}}_i^C = \mathbf{0}, \quad (14a)$$

$$\hat{\mathbf{v}}_i^C = [\mathbf{v}_i^C; \mathbf{0}; \mathbf{0}], \quad (14b)$$

$\forall i \in 1, \dots, n$ , which may be verified via simple substitution. Hence all critical eigenmodes of  $\mathbf{L}$  can be trivially extended to  $\hat{\mathbf{L}}$ . This definition for the extension of the eigenvectors in fact applies even to modes that are not critical eigenmodes.

The additional critical eigenpairs introduced due to system extension need to be deduced. Two sets of canonical unit vectors are defined such that,

$$\mathbf{e}_i^P = \underbrace{[0; 0; \dots; 0; 1; 0; 0; \dots; 0]}_{i-1}, \quad (15a)$$

$$\mathbf{e}_i^H = \underbrace{[0; 0; \dots; 0; 1; 0; 0; \dots; 0]}_{h-i}. \quad (15b)$$

If the original linear system  $\mathbf{L}$  is non-singular then, the  $l$  eigenpairs generated by the system extension due to  $\nu$  have the structure of the type  $(\lambda_i^P, [\mathbf{v}_i^P; \mathbf{e}_i^P; \mathbf{0}])$ , which satisfy

$$(\lambda_i^P \mathbf{I} - \mathbf{L})\mathbf{v}_i^P = \mathbf{L}_\mu(\mathbf{e}_i^P), \quad (\lambda_i^P = 0) \quad (16a)$$

$$\hat{\mathbf{v}}_i^P = [\mathbf{v}_i^P; \mathbf{e}_i^P; \mathbf{0}], \quad \forall i \in 1, \dots, l. \quad (16b)$$

This can be verified that  $(\lambda_i^P - \hat{\mathbf{L}})\hat{\mathbf{v}}_i^P = \mathbf{0}$ .

However, if  $\mathbf{L}$  is singular then equation (16) is indeterminate and the definition needs to be modified. For the  $i^{th}$  parameter mode, define a vector  $\gamma_i^P \in \mathbb{C}^n$ , where the  $j^{th}$  element of the vector is given by,

$$(\gamma_i^P)_j = \begin{cases} \mathbf{w}_j^{C\dagger} \mathbf{L}_\mu(\mathbf{e}_i^P), & \text{if } \lambda_i^P = \lambda_j^C \\ 0, & \text{otherwise.} \end{cases} \quad (17)$$

$$(\lambda_i^P \mathbf{I} - \mathbf{L})\mathbf{v}_i^P = \mathbf{L}_\mu(\mathbf{e}_i^P) - \mathbf{V}\gamma_i^P, \quad (\lambda_i^P = 0) \quad (18a)$$

$$\mathbf{w}_j^{C\dagger} \mathbf{v}_i^P = 0, \quad \text{if } \lambda_i^P = \lambda_j^C, \quad (18b)$$

$$\hat{\mathbf{v}}_i^P = [\mathbf{v}_i^P; \mathbf{e}_i^P; \mathbf{0}], \quad \forall i \in 1, \dots, l. \quad (18c)$$

In the singular case,  $\hat{\mathbf{v}}_i^P$  is no longer an eigenvector of  $\hat{\mathbf{L}}$ , however, it is a generalized eigenvector satisfying  $(\lambda_i^P \mathbf{I} - \hat{\mathbf{L}})^2 \hat{\mathbf{v}}_i^P = \mathbf{0}$ . Note that  $\gamma_i^P$  is identically zero for the non-singular case and equation (18) reduces to equation (16). Henceforth these modes shall be referred to as *parameter*

*modes* and will be defined by equation (18), since it is the more general definition. The matrix of all the vectors  $\gamma_i^P$  is denoted  $\mathbf{\Gamma}_P$ .

Similarly, when the harmonic forcing is non-resonant, the  $h$  eigenpairs generated by system extension due to  $\theta$  have the following structure,

$$(\lambda_i^H \mathbf{I} - \mathbf{L})\mathbf{v}_i^H = \mathbf{L}_\theta(\mathbf{e}_i^H), \quad (19a)$$

$$\hat{\mathbf{v}}_i^H = [\mathbf{v}_i^H; \mathbf{0}; \mathbf{e}_i^H], \quad \forall i \in 1, \dots, h. \quad (19b)$$

On the other hand, when the harmonic forcing is resonant with some of the angular frequencies of  $\mathbf{L}$  then, analogous to the case of parameter modes, one defines,

$$(\gamma_i^H)_j = \begin{cases} \mathbf{w}_j^{C\dagger} \mathbf{L}_\theta(\mathbf{e}_i^H), & \text{if } \lambda_i^H = \lambda_j^C \\ 0, & \text{otherwise,} \end{cases} \quad (20)$$

where,  $(\gamma_i^H)_j$  is the  $j^{th}$  component of the vector  $\gamma_i^H$ . The extended modes are then given by

$$(\lambda_i^H \mathbf{I} - \mathbf{L})\mathbf{v}_i^H = \mathbf{L}_\theta(\mathbf{e}_i^H) - \mathbf{V}\gamma_i^H, \quad (21a)$$

$$\mathbf{w}_j^{C\dagger} \mathbf{v}_i^H = 0, \quad \text{if } \lambda_i^H = \lambda_j^C, \quad (21b)$$

$$\hat{\mathbf{v}}_i^H = [\mathbf{v}_i^H; \mathbf{0}; \mathbf{e}_i^H], \quad \forall i \in 1, \dots, h. \quad (21c)$$

In the resonant case, this satisfies the generalized eigenvalue equation  $(\lambda_i^H \mathbf{I} - \hat{\mathbf{L}})^2 \hat{\mathbf{v}}_i^H = \mathbf{0}$ . Again, equation (21) reduces to equation (19) when the forcing is non-resonant. Henceforth these modes will be referred to as *forcing modes* and are defined by the more general definition given by equation (21). The matrix of all the vectors  $\gamma_i^H$  is denoted  $\mathbf{\Gamma}_H$ . The critical eigenspace of  $\hat{\mathbf{L}}$  can be represented by three matrices

$$\hat{\mathbf{V}}_C = \begin{bmatrix} \mathbf{V} \\ \mathbf{0} \\ \mathbf{0} \end{bmatrix}, \quad \hat{\mathbf{V}}_P = \begin{bmatrix} \mathbf{V}_P \\ \mathbf{I} \\ \mathbf{0} \end{bmatrix}, \quad \hat{\mathbf{V}}_H = \begin{bmatrix} \mathbf{V}_H \\ \mathbf{0} \\ \mathbf{I} \end{bmatrix}, \quad (22)$$

where,  $\mathbf{V}$  is the matrix of the critical eigenvectors of  $\mathbf{L}$ . The matrix  $\mathbf{V}_P$  comprises of the  $l$  vectors  $\mathbf{v}_i^P$  of the parameter modes, and,  $\mathbf{V}_H$  comprises of the  $h$  vectors  $\mathbf{v}_i^H$  of the forcing modes. The matrix of vectors of the extended eigenspace is denoted by,  $\hat{\mathbf{V}} = [\hat{\mathbf{V}}_C, \hat{\mathbf{V}}_P, \hat{\mathbf{V}}_H]$ .

### 1.3.2 Subspace structure for $\widehat{\mathbf{L}}^\dagger$

The eigenspace structure for the adjoint problem is important to consider since these vectors define the projection operators into the extended critical subspace. The critical eigenmodes of the  $\widehat{\mathbf{L}}^\dagger$  can be extended via the following definition,

$$(\lambda_i^{C*} \mathbf{I} - \mathbf{L}^\dagger) \mathbf{w}_i^C = 0, \quad \forall i \in 1, \dots, n, \quad (23a)$$

$$\widehat{\mathbf{w}}_i^C = [\mathbf{w}_i^C; \boldsymbol{\zeta}_i^P; \boldsymbol{\zeta}_i^H], \quad (23b)$$

with vectors  $\boldsymbol{\zeta}_i^P$  and  $\boldsymbol{\zeta}_i^H$  defined such that the  $j^{th}$  component of  $\boldsymbol{\zeta}_i^P$  given by,

$$(\boldsymbol{\zeta}_i^P)_j = \begin{cases} (\mathbf{L}_\mu^\dagger \mathbf{w}_i^C)_j / (\lambda_i^{C*} - \lambda_j^{P*}); & \text{if } \lambda_i^{C*} \neq \lambda_j^{P*}, \\ 0; & \text{if } \lambda_i^{C*} = \lambda_j^{P*}, \end{cases} \quad (24)$$

$\forall j \in 1, \dots, l$ , and, the  $j^{th}$  component of  $\boldsymbol{\zeta}_i^H$  is given by,

$$(\boldsymbol{\zeta}_i^H)_j = \begin{cases} (\mathbf{L}_\theta^\dagger \mathbf{w}_i^C)_j / (\lambda_i^{C*} - \lambda_j^{H*}); & \text{if } \lambda_i^{C*} \neq \lambda_j^{H*}, \\ 0; & \text{if } \lambda_i^{C*} = \lambda_j^{H*}, \end{cases} \quad (25)$$

$\forall j \in 1, \dots, h$ . Here,  $(\mathbf{L}_\mu^\dagger \mathbf{w}_i^C)_j$  refers to the  $j^{th}$  component of the vector  $(\mathbf{L}_\mu^\dagger \mathbf{w}_i^C)$ , and likewise for  $(\mathbf{L}_\theta^\dagger \mathbf{w}_i^C)_j$ . It may be verified that the definition is consistent with the eigenvalue equation,  $(\lambda_i^{C*} \mathbf{I} - \widehat{\mathbf{L}}^\dagger) \widehat{\mathbf{w}}_i^C = 0$ , for the non-resonant cases, and satisfies generalized eigenvalue equation,  $(\lambda_i^{C*} \mathbf{I} - \widehat{\mathbf{L}}^\dagger)^2 \widehat{\mathbf{w}}_i^C = 0$ , for the resonant cases.

The adjoint parameter modes are given simply by,

$$\widehat{\mathbf{w}}_i^P = [\mathbf{0}; \mathbf{e}_i^P; \mathbf{0}], \quad \forall i \in 1, \dots, l, \quad (26)$$

which satisfies  $(\lambda_i^{P*} \mathbf{I} - \widehat{\mathbf{L}}^\dagger) \widehat{\mathbf{w}}_i^P = 0$ . And the adjoint forcing modes are given by,

$$\widehat{\mathbf{w}}_i^H = [\mathbf{0}; \mathbf{0}; \mathbf{e}_i^H], \quad \forall i \in 1, \dots, h, \quad (27)$$

which satisfies  $(\lambda_i^{H*} \mathbf{I} - \widehat{\mathbf{L}}^\dagger) \widehat{\mathbf{w}}_i^H = 0$ .

The extended eigenspace of  $\widehat{\mathbf{L}}^\dagger$  can be represented with matrices as

$$\widehat{\mathbf{W}}_C = \begin{bmatrix} \mathbf{W} \\ \mathbf{Z}_P \\ \mathbf{Z}_H \end{bmatrix}, \quad \widehat{\mathbf{W}}_P = \begin{bmatrix} \mathbf{0} \\ \mathbf{I} \\ \mathbf{0} \end{bmatrix}, \quad \widehat{\mathbf{W}}_H = \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \\ \mathbf{I} \end{bmatrix}, \quad (28)$$

where,  $\mathbf{W}$  is the matrix of the adjoint critical eigenvectors of  $\mathbf{L}^\dagger$ . The matrix  $\mathbf{Z}_P$  comprises of the  $l$  vectors  $\boldsymbol{\zeta}_i^P$  and,  $\mathbf{Z}_H$  comprises of the  $h$  vectors  $\boldsymbol{\zeta}_i^H$ . The matrix of vectors of the critical adjoint eigenspace is denoted by,  $\widehat{\mathbf{W}} = [\widehat{\mathbf{W}}_C, \widehat{\mathbf{W}}_P, \widehat{\mathbf{W}}_H]$ . It can be verified that,

$$\widehat{\mathbf{W}}^\dagger \widehat{\mathbf{V}} = \mathbf{I}_m, \quad (29)$$

where,  $m = n + l + h$ . This is the analogue of equation (6), and is simply a statement of the biorthogonality relation between the direct and adjoint eigenvectors in the extended space. The projection matrices into the critical and stable subspaces of  $\widehat{\mathbf{L}}$  are given by,

$$\widehat{\mathbf{P}} = \widehat{\mathbf{V}} \widehat{\mathbf{W}}^\dagger \quad (30)$$

$$\Rightarrow \widehat{\mathbf{P}} = \widehat{\mathbf{V}}_C \widehat{\mathbf{W}}_C^\dagger + \widehat{\mathbf{V}}_P \widehat{\mathbf{W}}_P^\dagger + \widehat{\mathbf{V}}_H \widehat{\mathbf{W}}_H^\dagger,$$

$$\widehat{\mathbf{Q}} = \mathbf{I} - \widehat{\mathbf{V}} \widehat{\mathbf{W}}^\dagger, \quad (31)$$

$$\Rightarrow \widehat{\mathbf{Q}} = \mathbf{I} - \widehat{\mathbf{V}}_C \widehat{\mathbf{W}}_C^\dagger - \widehat{\mathbf{V}}_P \widehat{\mathbf{W}}_P^\dagger - \widehat{\mathbf{V}}_H \widehat{\mathbf{W}}_H^\dagger.$$

Finally, a reduced matrix representing the oblique projection of the extended linear operator  $\widehat{\mathbf{L}}$  onto the critical subspace is defined by,

$$\widehat{\mathbf{K}} = \widehat{\mathbf{W}}^\dagger \widehat{\mathbf{L}} \widehat{\mathbf{V}} = \begin{bmatrix} \Lambda_C & \Gamma_P & \Gamma_H \\ \mathbf{0} & \Lambda_P & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \Lambda_H \end{bmatrix}, \quad (32)$$

which is a complex matrix of size  $m \times m$ . The diagonal elements of  $\widehat{\mathbf{K}}$  represent the eigenvalues of the reduced system, that are the critical eigenvalues of the extended system, and may be represented by the vector  $\widehat{\boldsymbol{\lambda}} = \text{diag}(\widehat{\mathbf{K}})$ .

Note that due to the special structure of the adjoint parameter and forcing modes, the projection of the extended nonlinearity  $\widehat{\mathbf{f}}_{NL}$  onto the subspace spanned by the (direct) parameter and forcing modes vanishes identically. That is to say,

$$\widehat{\mathbf{W}}_P^\dagger \widehat{\mathbf{f}}_{NL} = \mathbf{0}, \quad \widehat{\mathbf{W}}_H^\dagger \widehat{\mathbf{f}}_{NL} = \mathbf{0}. \quad (33)$$

This completes the extension of the system and the subspace structure for both the direct and adjoint systems can be readily identified. While the structure of the extended subspaces has been evaluated, reformulation of the problem as an extended system may have detrimental practical implications when considering large systems.

Practical methods for solving such systems often rely on numerical algorithms that exploit inherent structure of the problem (Hermicity for conjugate-gradient solvers for example). In such a scenario, additional degrees of freedom can potential destroy the system symmetries and may require substantial changes in the numerical approach to the system solution. This would be particularly true of asymptotic approaches that require resolvent calculations [1, 2, 3, 4, 5, 6, 7, 8, 9], which itself is a computationally challenging task for large systems. Such a scenario would be highly undesirable. However, as it turns out, at least some of the parameterizations involved in the evaluation of center-manifolds do not in fact require the use of the extended system after the initial evaluation of the extended tangent space. All higher order calculations can be performed using the original linear system  $\mathbf{L}$ . This is clarified in the next section where the parameterization and its asymptotic terms are explicitly built. First however, we prove the following results which will come in handy.

**Lemma 1.** *Given an extended system described by equation (8), with a linearized operator  $\widehat{\mathbf{L}}$ , given by equation (12), if a vector,  $\widehat{\mathbf{u}} = [\mathbf{u}; \boldsymbol{\nu}; \boldsymbol{\theta}]$ , is such that its projection onto the center-subspace of  $\widehat{\mathbf{L}}$  vanishes, then, the extended variables  $\boldsymbol{\nu}$  and  $\boldsymbol{\theta}$  must also vanish identically for  $\widehat{\mathbf{u}}$ .*

*Proof* A vector  $\widehat{\mathbf{u}}$  whose components along the center-subspace vanish implies that  $\widehat{\mathbf{W}}^\dagger \widehat{\mathbf{u}} = \mathbf{0}$ , where  $\widehat{\mathbf{W}} = [\widehat{\mathbf{W}}_C, \widehat{\mathbf{W}}_P, \widehat{\mathbf{W}}_H]$ , are as deduced in sections 1.2 and 1.3.

$$\therefore \widehat{\mathbf{W}}^\dagger \widehat{\mathbf{u}} = \begin{bmatrix} \mathbf{W}^\dagger \mathbf{u} + \mathbf{Z}_P^\dagger \boldsymbol{\nu} + \mathbf{Z}_H^\dagger \boldsymbol{\theta} \\ \mathbf{I}^\dagger \boldsymbol{\nu} \\ \mathbf{I}^\dagger \boldsymbol{\theta} \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \\ \mathbf{0} \end{bmatrix}.$$

For this to hold,

$$\mathbf{I}^\dagger \boldsymbol{\nu} = \mathbf{0}, \implies \boldsymbol{\nu} = \mathbf{0},$$

$$\mathbf{I}^\dagger \boldsymbol{\theta} = \mathbf{0}, \implies \boldsymbol{\theta} = \mathbf{0}.$$

Hence, both the extended variables  $\boldsymbol{\nu}$  and  $\boldsymbol{\theta}$  vanish identically.  $\square$

**Corollary 1.** *Given an extended system described by equation (8), with a linearized operator  $\widehat{\mathbf{L}}$ , given by equation (12), if a vector,  $\widehat{\mathbf{u}} = [\mathbf{u}; \boldsymbol{\nu}; \boldsymbol{\theta}]$ , is such that its projection onto the center-subspace of  $\widehat{\mathbf{L}}$  vanishes, then, the projection of  $\mathbf{u}$  onto the center-subspace of  $\mathbf{L}$  also vanishes.*

*Proof* For a vector  $\widehat{\mathbf{u}}$  that satisfies  $\widehat{\mathbf{W}}^\dagger \widehat{\mathbf{u}} = \mathbf{0}$ , by lemma 1, both  $\boldsymbol{\nu}$  and  $\boldsymbol{\theta}$  vanish identically.

$$\therefore \widehat{\mathbf{W}}^\dagger \widehat{\mathbf{u}} = \begin{bmatrix} \mathbf{W}^\dagger \mathbf{u} \\ \mathbf{0} \\ \mathbf{0} \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \\ \mathbf{0} \end{bmatrix},$$

which implies  $\mathbf{W}^\dagger \mathbf{u} = \mathbf{0}$ . However,  $\mathbf{W}$  spans the center-subspace of  $\mathbf{L}^\dagger$ , and therefore  $\mathbf{W}^\dagger \mathbf{u}$  is the projection of  $\mathbf{u}$  onto the center-subspace of  $\mathbf{L}$ , which vanishes as required.  $\square$

A proof of corollary 1 in the case of infinite dimensional systems with parameter extensions can be found in [4].

## 2 Center Manifold for the extended system

In the context of invariant manifold theory, several previous approaches for the approximation of such manifolds can be effectively viewed as instances of what is referred to as the parametrization method. The framework was introduced in a series of papers by Cabré et al. [10, 11, 2], although, a particular instance of such an approach was presented earlier by Couillet and Spiegel [1] for the evaluation of the normal form of center manifolds. Nevertheless, subsequent works by Carini, Auteri, and Giannetti [5], Jain and Haller [8], Negi [9] can be fruitfully viewed through the lens of the parameterization method. Within this framework the reduced representation of the invariant manifold being approximated can be chosen in an arbitrary manner depending on the particular requirements of the reduction being sought. Amongst the plethora of arbitrary choices, the normal form is an important representation that is often sought since it represents the simplest form of the reduced equations. In this context, it can be shown that for asymptotic normal form evaluation of the extended system the calculations involve only the resolvents of the original linear system and therefore all existing algorithms may be utilized for the calculations.

To simplify notation for multiple indices used hereafter, a multi-dimensional index set  $\mathcal{I}\{(k), [a, b]\}$  is introduced, which is an ordered set



of all  $k$ -indices  $(i_1, i_2, \dots, i_k)$ , defined as

$$\begin{aligned} \mathcal{I}\{(k), [a, b]\} := & \{(i_1, i_2, \dots, i_k) \mid a, b \in \mathbb{N}, \\ & i_1, i_2, \dots, i_k \in \mathbb{N}, \\ & a \leq i_i \leq b, \ i_1 \leq i_2 \leq b, \\ & \dots, \ i_{k-1} \leq i_k \leq b.\}, \end{aligned} \quad (34)$$

where,  $i_1, i_2, \dots, i_k$ , are the index variables and  $[a, b]$  specify the lower and upper bounds of the indices. Note that the higher index always starts from the previous index. Hence the index-set  $\mathcal{I}\{(2), [1, 3]\}$  is the set containing the tuples,  $(i_1, i_2) \in \{(1, 1), (1, 2), (1, 3), (2, 2), (2, 3), (3, 3)\}$ . Similarly, multiple nested summations are denoted as,

$$\sum_{(k)}^{[a, b]} = \sum_{i_1=a}^b \sum_{i_2=i_1}^b \dots \sum_{i_k=i_{k-1}}^b \quad (35)$$

As before, the higher index starts from the previous index. For the asymptotic expansions we will typically have  $a = 1$  and  $b = m$ , where  $m$  is the dimension of the manifold. For such cases the number of  $k$ -tuples in a multi-index tuple set  $\mathcal{I}\{(k), [1, m]\}$  is analytically known to be  $T_m^k = C(m + k - 1, k)$ , where,

$$C(p, q) = \binom{p}{q} = \frac{p!}{q!(p-q)!}, \quad (36)$$

defines the binomial coefficient of  $p$  and  $q$ .

## 2.1 Normal Form

One may find the exposition on the implementation of the parameterization method in previous works of Haro et al. [7] and Jain and Haller [8]. To be clear, the application of the parameterization method to an extended system like equation (8) follows the same steps as outlined in those earlier works. However, in order to show that the resolvent calculations in fact only require the original linear system, and not the extended one, an explanation of the key steps of the method is necessary.

Consider the extended system as defined by equation (8), represented in a compact form by equation (13), which has an  $m = n + l + h$  dimensional critical subspace. Assume  $\mathbf{x} = [x_1; x_2; \dots; x_m]$  is an  $m$ -dimensional vector of coordinates that parameterize the center-manifold of the extended system. The evolution of the

system along the center-manifold in the reduced coordinates is assumed to be given by  $\dot{\mathbf{x}} = \mathcal{G}(\mathbf{x})$ , where,  $\mathcal{G}$  is a vector valued function. The system solution on the center-manifold corresponding to a point  $\mathbf{x}$  on the reduced system is given by  $\mathcal{Y}(\mathbf{x})$ . At this stage neither  $\mathcal{G}(\mathbf{x})$  nor  $\mathcal{Y}(\mathbf{x})$  have been specified in any form, since, the parameterization can be arbitrarily chosen. In order to obtain a normal form of the reduced system one may expand both  $\mathcal{G}(\mathbf{x})$  and  $\mathcal{Y}(\mathbf{x})$  as a formal power series in  $\mathbf{x}$  and constrain the asymptotic form of  $\mathcal{G}$  to be in a normal form for the parameterized coordinates  $\mathbf{x}$ . The asymptotic form of  $\mathcal{Y}$  is obtained by introducing the asymptotic expansions into equation (13) and solving the terms order-by-order.

In what follows, Einstein summation is *not* assumed for repeated indices unless otherwise specified. We assume the power series expansions as,

$$\begin{aligned} \mathcal{G}(\mathbf{x}) = & \sum_{(1)}^{[1, m]} \mathbf{g}_{i_1} x_{i_1} + \sum_{(2)}^{[1, m]} \mathbf{g}_{i_1, i_2} x_{i_1} x_{i_2} + \\ & \sum_{(3)}^{[1, m]} \mathbf{g}_{i_1, i_2, i_3} x_{i_1} x_{i_2} x_{i_3} + \dots, \end{aligned} \quad (37)$$

and,

$$\begin{aligned} \mathcal{Y}(\mathbf{x}) = & \sum_{(1)}^{[1, m]} \hat{\mathbf{y}}_{i_1} x_{i_1} + \sum_{(2)}^{[1, m]} \hat{\mathbf{y}}_{i_1, i_2} x_{i_1} x_{i_2} + \\ & \sum_{(3)}^{[1, m]} \hat{\mathbf{y}}_{i_1, i_2, i_3} x_{i_1} x_{i_2} x_{i_3} + \dots \end{aligned} \quad (38)$$

Here the terms  $\mathbf{g}_{i_1, i_2, i_3, \dots}$  are  $m$ -dimensional vectors while,  $\hat{\mathbf{y}}_{i_1, i_2, i_3, \dots}$  are vectors of size  $(N + l + h)$ , consistent with the total dimension of the extended system. Given the asymptotic form of  $\mathcal{Y}(\mathbf{x})$ , one automatically obtains the nonlinear term  $\hat{\mathbf{f}}_{NL}$  as a power series in  $\mathbf{x}$ . This can be represented as,

$$\begin{aligned} \hat{\mathbf{f}}_{NL} = & \sum_{(2)}^{[1, m]} \hat{\mathbf{f}}_{i_1, i_2} x_{i_1} x_{i_2} + \\ & \sum_{(3)}^{[1, m]} \hat{\mathbf{f}}_{i_1, i_2, i_3} x_{i_1} x_{i_2} x_{i_3} + \dots, \end{aligned} \quad (39)$$

which is obtained via the grouping of terms with similar powers of  $\mathbf{x}$ .

Substitution of the power series expansions into equation (13) results in

$$\frac{\partial}{\partial t} \left( \sum_{(1)}^{[1,m]} \hat{\mathbf{y}}_{i_1} x_{i_1} + \dots \right) = \hat{\mathbf{L}} \left( \sum_{(1)}^{[1,m]} \hat{\mathbf{y}}_{i_1} x_{i_1} + \dots \right) + \left( \sum_{(2)}^{[1,m]} \hat{\mathbf{f}}_{i_1, i_2} x_{i_1} x_{i_2} + \dots \right),$$

resulting in,

$$\sum_{\alpha=1}^m \frac{\partial}{\partial x_{\alpha}} \left( \sum_{(1)}^{[1,m]} \hat{\mathbf{y}}_{i_1} x_{i_1} + \dots \right) \left( \sum_{(1)}^{[1,m]} \mathbf{g}_{i_1}^{\alpha} x_{i_1} + \dots \right) = \hat{\mathbf{L}} \left( \sum_{(1)}^{[1,m]} \hat{\mathbf{y}}_{i_1} x_{i_1} + \dots \right) + \left( \sum_{(2)}^{[1,m]} \hat{\mathbf{f}}_{i_1, i_2} x_{i_1} x_{i_2} + \dots \right). \quad (40)$$

Here  $\mathbf{g}_{i_1}^{\alpha}$  represents the  $\alpha^{th}$  element of the vector  $\mathbf{g}_{i_1}$ .

At first order the non-linear terms do not contribute and one simply obtains,

$$\sum_{\alpha=1}^m \left( \hat{\mathbf{y}}_{\alpha} \sum_{(1)}^{[1,m]} \mathbf{g}_{i_1}^{\alpha} x_{i_1} \right) = \hat{\mathbf{L}} \left( \sum_{(1)}^{[1,m]} \hat{\mathbf{y}}_{i_1} x_{i_1} \right) \Rightarrow \sum_{(1)}^{[1,m]} \left( \sum_{\alpha=1}^m \hat{\mathbf{y}}_{\alpha} \mathbf{g}_{i_1}^{\alpha} - \hat{\mathbf{L}} \hat{\mathbf{y}}_{i_1} \right) = \mathbf{0}$$

## References

- [1] P. H. Coullet and E. A. Spiegel. Amplitude equations for systems with competing instabilities. *SIAM Journal on Applied Mathematics*, 43(4):776–821, 1983. doi: 10.1137/0143052.
- [2] Xavier Cabré, Ernest Fontich, and Rafael De La Llave. The parameterization method for invariant manifolds iii: overview and applications. *Journal of Differential Equations*, 218(2):444–515, 2005.
- [3] D. Sipp and A. Lebedev. Global stability of base and mean flows: a general approach and its applications to cylinder and open cavity flows. *Journal of Fluid*

*Mechanics*, 593:333–358, 2007. doi: 10.1017/S0022112007008907.

- [4] Mariana Haragus and Iooss Gérard. *Local Bifurcations, Center Manifolds, and Normal Forms in Infinite-Dimensional Dynamical Systems*. Springer-Verlag, New York, 2011. doi: 10.1007/978-0-85729-112-7.
- [5] M. Carini, F. Auteri, and F. Giannetti. Centre-manifold reduction of bifurcating flows. *Journal of Fluid Mechanics*, 767: 109–145, 2015. doi: 10.1017/jfm.2015.3.
- [6] Francois Gallaire, Edouard Boujo, Vladislav Mantic-Lugo, Cristobal Arratia, Benjamin Thiria, and Philippe Meliga. Pushing amplitude equations far from threshold: application to the supercritical hopf bifurcation in the cylinder wake. *Fluid Dynamics Research*, 48(6):061401, 2016.
- [7] Alex Haro, Marta Canadell, Jordi-Lluís Figueras, Alejandro Luque, Josep Maria Mondelo, et al. *The parameterization method for invariant manifolds*. Springer, 2016.
- [8] Shobhit Jain and George Haller. How to compute invariant manifolds and their reduced dynamics in high-dimensional finite element models. *Nonlinear dynamics*, 107(2):1417–1450, 2022.
- [9] Prabal S Negi. Asymptotic center-manifold for the navier-stokes. *arXiv preprint arXiv:2411.03727*, 2024.
- [10] Xavier Cabré, Ernest Fontich, and Rafael De la Llave. The parameterization method for invariant manifolds i: manifolds associated to non-resonant subspaces. *Indiana University mathematics journal*, pages 283–328, 2003.
- [11] Xavier Cabré, Ernest Fontich, and Rafael De la Llave. The parameterization method for invariant manifolds ii: regularity with respect to parameters. *Indiana University mathematics journal*, pages 329–360, 2003.
- [12] John Guckenheimer and Philip Holmes. *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*. Springer Science & Business Media, New York, 1983. doi: 10.1007/b97481.
- [13] Stephen Wiggins. *Introduction to Applied Nonlinear Dynamical Systems and Chaos*. Springer-Verlag New York, 2003. doi: 10.1007/b97481.



- [14] Jack Carr. *Applications of centre manifold theory*, volume 35. Springer Science & Business Media, 1982.
- [15] Jan Sijbrand. Properties of center manifolds. *Transactions of the American Mathematical Society*, 289(2):431–469, 1985. ISSN 00029947.
- [16] Alan C. Newell and J. A. Whitehead. Finite bandwidth, finite amplitude convection. *Journal of Fluid Mechanics*, 38(2):279–303, 1969. doi: 10.1017/S0022112069000176.
- [17] M. C. Cross. Derivation of the amplitude equation at the Rayleigh–Bénard instability. *The Physics of Fluids*, 23(9):1727–1731, 09 1980. ISSN 0031-9171. doi: 10.1063/1.863198.
- [18] Michael Cross and Henry Greenside. *Pattern Formation and Dynamics in Nonequilibrium Systems*. Cambridge University Press, 2009.
- [19] J. T. Stuart. On the non-linear mechanics of hydrodynamic stability. *Journal of Fluid Mechanics*, 4(1):1–21, 1958. doi: 10.1017/S0022112058000276.
- [20] J. T. Stuart. On the non-linear mechanics of wave disturbances in stable and unstable parallel flows part 1. the basic behaviour in plane poiseuille flow. *Journal of Fluid Mechanics*, 9(3):353–370, 1960. doi: 10.1017/S002211206000116X.