

Created by Prabal Negi

Navier-Stokes in Cylindrical formulation

Weak form and Nek implementation

Formulation of Navier-Stokes 1

Mapping from Cartesian to Cylindrical 1.1

$$x_c = x \tag{1a}$$

$$y_c = r\cos(\theta) \tag{1b}$$

$$z_c = r\sin(\theta) \tag{1c}$$

$$\begin{cases} dx_c \\ dy_c \\ dz_c \end{cases} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -r\sin(\theta) \\ 0 & \sin(\theta) & r\cos(\theta) \end{pmatrix} \begin{cases} dx \\ dr \\ d\theta \end{cases}$$
 (2)

$$\begin{cases}
 dx_c \\
 dy_c \\
 dz_c
\end{cases} = \begin{pmatrix}
 1 & 0 & 0 \\
 0 & \cos(\theta) & -r\sin(\theta) \\
 0 & \sin(\theta) & r\cos(\theta)
\end{pmatrix} \begin{cases}
 dx \\
 dr \\
 d\theta
\end{cases}$$

$$\Rightarrow \begin{cases}
 dx \\
 dr \\
 d\theta
\end{cases} = \begin{pmatrix}
 1 & 0 & 0 \\
 0 & \cos(\theta) & \sin(\theta) \\
 0 & -\sin(\theta)/r & \cos(\theta)/r
\end{pmatrix} \begin{cases}
 dx_c \\
 dy_c \\
 dz_c
\end{cases}$$
(2)

$$\hat{x} = \hat{x_c} \tag{4a}$$

$$\hat{r} = \cos(\theta)\hat{y_c} + \sin(\theta)\hat{z_c} \tag{4b}$$

$$\hat{\theta} = -\sin(\theta)\hat{y_c} + \cos(\theta)\hat{z_c} \tag{4c}$$

$$\Rightarrow \begin{cases} d\hat{x} \\ d\hat{r} \\ d\hat{\theta} \end{cases} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -\sin(\theta)\hat{y}_c + \cos(\theta)\hat{z}_c \\ 0 & 0 & -\cos(\theta)\hat{y}_c - \sin(\theta)\hat{z}_c \end{pmatrix} \begin{cases} dx \\ dr \\ d\theta \end{cases}$$

$$\Rightarrow \begin{cases} d\hat{x} \\ d\hat{r} \\ d\hat{\theta} \end{cases} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \hat{\theta} \\ 0 & 0 & -\hat{r} \end{pmatrix} \begin{cases} dx \\ dr \\ d\theta \end{cases}$$
(5)

1.2 Gradient/Divergence/Laplace Operators

For scalar fields

$$\nabla(\psi) = \left(\frac{\partial \psi}{\partial x} \frac{\partial x}{\partial x_c} + \frac{\partial \psi}{\partial r} \frac{\partial r}{\partial x_c} + \frac{\partial \psi}{\partial \theta} \frac{\partial \theta}{\partial x_c}\right) \hat{x_c} + \left(\frac{\partial \psi}{\partial x} \frac{\partial x}{\partial y_c} + \frac{\partial \psi}{\partial r} \frac{\partial r}{\partial y_c} + \frac{\partial \psi}{\partial \theta} \frac{\partial \theta}{\partial y_c}\right) \hat{y_c} + \left(\frac{\partial \psi}{\partial x} \frac{\partial x}{\partial z_c} + \frac{\partial \psi}{\partial r} \frac{\partial r}{\partial z_c} + \frac{\partial \psi}{\partial \theta} \frac{\partial \theta}{\partial z_c}\right) \hat{z_c}$$

$$\therefore \nabla(\psi) = \frac{\partial \psi}{\partial x}\hat{x} + \frac{\partial \psi}{\partial r}\hat{r} + \frac{1}{r}\frac{\partial \psi}{\partial \theta}\hat{\theta}$$
 (6)

Laplacian of a scalar field

$$\nabla \cdot \nabla(\psi) = \left[\frac{\partial}{\partial x} \hat{x} + \frac{\partial}{\partial r} \hat{r} + \frac{1}{r} \frac{\partial}{\partial \theta} \hat{\theta} \right] \cdot \left[\hat{x} \frac{\partial \psi}{\partial x} + \hat{r} \frac{\partial \psi}{\partial r} + \hat{\theta} \frac{1}{r} \frac{\partial \psi}{\partial \theta} \right]$$

$$= \begin{cases} \left[\hat{x} \cdot \hat{x} \frac{\partial}{\partial x} \frac{\partial \psi}{\partial x} + \frac{\partial \psi}{\partial x} \frac{\partial \hat{x}}{\partial x} \cdot \hat{x} \right] + \left[\hat{x} \cdot \hat{r} \frac{\partial}{\partial x} \frac{\partial \psi}{\partial r} + \frac{\partial \psi}{\partial r} \frac{\partial \hat{r}}{\partial x} \cdot \hat{x} \right] + \left[\hat{x} \cdot \hat{\theta} \frac{\partial}{\partial x} \left(\frac{1}{r} \frac{\partial \psi}{\partial \theta} \right) + \frac{1}{r} \frac{\partial \psi}{\partial \theta} \frac{\partial \hat{\theta}}{\partial x} \cdot \hat{x} \right] \right] \\ + \left[\hat{r} \cdot \hat{x} \frac{\partial}{\partial r} \frac{\partial \psi}{\partial x} + \frac{\partial \psi}{\partial x} \frac{\partial \hat{x}}{\partial r} \cdot \hat{r} \right] + \left[\hat{r} \cdot \hat{r} \frac{\partial}{\partial r} \frac{\partial \psi}{\partial r} + \frac{\partial \psi}{\partial r} \frac{\partial \hat{r}}{\partial r} \cdot \hat{r} \right] + \left[\hat{r} \cdot \hat{\theta} \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial \psi}{\partial \theta} \right) + \frac{1}{r} \frac{\partial \psi}{\partial \theta} \frac{\partial \hat{\theta}}{\partial r} \cdot \hat{r} \right] \\ + \left[\hat{\theta} \cdot \hat{x} \frac{1}{r} \frac{\partial}{\partial \theta} \frac{\partial \psi}{\partial x} + \frac{1}{r} \frac{\partial \psi}{\partial x} \frac{\partial \hat{x}}{\partial \theta} \cdot \hat{\theta} \right] + \left[\hat{\theta} \cdot \hat{r} \frac{1}{r} \frac{\partial}{\partial \theta} \frac{\partial \psi}{\partial r} + \frac{1}{r} \frac{\partial \psi}{\partial r} \frac{\partial \hat{\theta}}{\partial \theta} \cdot \hat{\theta} \right] \\ + \left[\hat{\theta} \cdot \hat{\theta} \frac{1}{r} \frac{\partial}{\partial \theta} \left(\frac{1}{r} \frac{\partial \psi}{\partial \theta} \right) + \frac{1}{r^2} \frac{\partial \psi}{\partial \theta} \frac{\partial \hat{\theta}}{\partial \theta} \cdot \hat{\theta} \right]$$

$$\therefore \nabla^2(\psi) = \frac{\partial^2 \psi}{\partial x^2} + \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial \psi}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 \psi}{\partial \theta}$$
 (7)

Vector gradient. Highlighted in color are the only terms which are non-zero differentials of unit vectors.

$$\nabla(\boldsymbol{u}) = \left[\hat{x} \frac{\partial}{\partial x} + \hat{r} \frac{\partial}{\partial r} + \hat{\theta} \frac{1}{r} \frac{\partial}{\partial \theta} \right] \left[\hat{x} u_x + \hat{r} u_r + \hat{\theta} u_\theta \right]$$

$$= \left\{ \begin{bmatrix} \hat{x} \frac{\partial u_x}{\partial x} \hat{x} + \hat{x} u_x \frac{\partial \hat{x}}{\partial x} \end{bmatrix} + \left[\hat{x} \frac{\partial u_r}{\partial x} \hat{r} + \hat{x} u_r \frac{\partial \hat{r}}{\partial x} \right] + \left[\hat{x} \frac{\partial u_\theta}{\partial x} \hat{\theta} + \hat{x} u_\theta \frac{\partial \hat{\theta}}{\partial x} \right]$$

$$+ \left[\hat{r} \frac{\partial u_x}{\partial r} \hat{x} + \hat{r} u_x \frac{\partial \hat{x}}{\partial r} \right] + \left[\hat{r} \frac{\partial u_r}{\partial r} \hat{r} + \hat{r} u_r \frac{\partial \hat{r}}{\partial r} \right] + \left[\hat{r} \frac{\partial u_\theta}{\partial r} \hat{\theta} + \hat{r} u_\theta \frac{\partial \hat{\theta}}{\partial r} \right]$$

$$+ \left[\hat{\theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta} \hat{x} + \hat{\theta} \frac{u_x}{r} \frac{\partial \hat{x}}{\partial \theta} \right] + \left[\hat{\theta} \frac{1}{r} \frac{\partial u_r}{\partial \theta} \hat{r} + \hat{\theta} \frac{u_r}{r} \frac{\partial \hat{r}}{\partial \theta} \right] + \left[\hat{\theta} \frac{1}{r} \frac{\partial u_\theta}{\partial \theta} \hat{\theta} + \hat{\theta} \frac{u_\theta}{r} \frac{\partial \hat{\theta}}{\partial \theta} \right]$$

$$(8)$$

$$\therefore \nabla(\boldsymbol{u}) = \begin{bmatrix} \hat{x} \frac{\partial u_x}{\partial x} \hat{x} & \hat{x} \frac{\partial u_r}{\partial x} \hat{r} & \hat{x} \frac{\partial u_\theta}{\partial x} \hat{\theta} \\ \hat{r} \frac{\partial u_x}{\partial r} \hat{x} & \hat{r} \frac{\partial u_r}{\partial r} \hat{r} & \hat{r} \frac{\partial u_\theta}{\partial r} \hat{\theta} \\ \hat{\theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta} \hat{x} & \hat{\theta} \left(-\frac{u_\theta}{r} + \frac{1}{r} \frac{\partial u_r}{\partial \theta} \right) \hat{r} & \hat{\theta} \left(\frac{u_r}{r} + \frac{1}{r} \frac{\partial u_\theta}{\partial \theta} \right) \hat{\theta} \end{bmatrix} \tag{9}$$

Vector Laplacian:

$$\nabla \cdot \nabla (\boldsymbol{u}) = \left(\hat{\boldsymbol{x}} \cdot \frac{\partial}{\partial \boldsymbol{x}} + \hat{\boldsymbol{r}} \cdot \frac{\partial}{\partial r} + \hat{\boldsymbol{\theta}} \cdot \frac{1}{r} \frac{\partial}{\partial \theta} \right) \begin{pmatrix} \hat{\boldsymbol{x}} \frac{\partial u_x}{\partial \boldsymbol{x}} \hat{\boldsymbol{x}} + & \hat{\boldsymbol{x}} \frac{\partial u_r}{\partial \boldsymbol{x}} \hat{\boldsymbol{r}} + & \hat{\boldsymbol{x}} \frac{\partial u_\theta}{\partial \boldsymbol{x}} \hat{\boldsymbol{\theta}} + \\ \hat{\boldsymbol{r}} \frac{\partial u_x}{\partial r} \hat{\boldsymbol{x}} + & \hat{\boldsymbol{r}} \frac{\partial u_r}{\partial r} \hat{\boldsymbol{r}} + & \hat{\boldsymbol{r}} \frac{\partial u_\theta}{\partial r} \hat{\boldsymbol{\theta}} + \\ \hat{\boldsymbol{\theta}} \frac{1}{r} \frac{\partial u_x}{\partial \theta} \hat{\boldsymbol{x}} + & \hat{\boldsymbol{\theta}} \left(-\frac{u_\theta}{r} + \frac{1}{r} \frac{\partial u_r}{\partial \theta} \right) \hat{\boldsymbol{r}} + & \hat{\boldsymbol{\theta}} \left(\frac{u_r}{r} + \frac{1}{r} \frac{\partial u_\theta}{\partial \theta} \right) \hat{\boldsymbol{\theta}} \end{pmatrix}$$

First Column

$$\left(\hat{x} \cdot \frac{\partial}{\partial x} + \hat{r} \cdot \frac{\partial}{\partial r} + \hat{\theta} \cdot \frac{1}{r} \frac{\partial}{\partial \theta}\right) \left(\hat{x} \frac{\partial u_x}{\partial x} + \hat{r} \frac{\partial u_x}{\partial r} + \hat{\theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) \hat{x}$$

$$= \hat{x} \cdot \left(\hat{x} \frac{\partial^2 u_x}{\partial x^2} + \frac{\partial \hat{x}}{\partial x} \frac{\partial u_x}{\partial x}\right) \hat{x} + \hat{x} \cdot \hat{x} \frac{\partial u_x}{\partial x} \frac{\partial \hat{x}}{\partial x}$$

$$+ \hat{x} \cdot \left(\hat{r} \frac{\partial}{\partial x} \frac{\partial u_x}{\partial r} + \frac{\partial \hat{r}}{\partial x} \frac{\partial u_x}{\partial r}\right) \hat{x} + \hat{x} \cdot \hat{r} \frac{\partial u_x}{\partial r} \frac{\partial \hat{x}}{\partial r}$$

$$+ \hat{x} \cdot \left(\hat{\theta} \frac{\partial}{\partial x} \left(\frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) + \frac{\partial \hat{\theta}}{\partial x} \frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) \hat{x} + \hat{x} \cdot \hat{\theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta} \frac{\partial \hat{x}}{\partial x}$$

$$+ \hat{r} \cdot \left(\hat{x} \frac{\partial}{\partial r} \frac{\partial u_x}{\partial x} + \frac{\partial \hat{x}}{\partial r} \frac{\partial u_x}{\partial x}\right) \hat{x} + \hat{r} \cdot \hat{x} \frac{\partial u_x}{\partial x} \frac{\partial \hat{x}}{\partial r}$$

$$+ \hat{r} \cdot \left(\hat{r} \frac{\partial^2 u_x}{\partial r^2} + \frac{\partial \hat{r}}{\partial r} \frac{\partial u_x}{\partial r}\right) \hat{x} + \hat{r} \cdot \hat{r} \frac{\partial u_x}{\partial r} \frac{\partial \hat{x}}{\partial r}$$

$$+ \hat{r} \cdot \left(\hat{\theta} \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) + \frac{\partial \hat{\theta}}{\partial r} \frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) \hat{x} + \hat{r} \cdot \hat{\theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta} \frac{\partial \hat{x}}{\partial r}$$

$$+ \hat{\theta} \cdot \left(\hat{x} \frac{1}{r} \frac{\partial}{\partial \theta} \frac{\partial u_x}{\partial x} + \frac{\partial \hat{x}}{\partial \theta} \frac{1}{r} \frac{\partial u_x}{\partial x}\right) \hat{x} + \hat{\theta} \cdot \hat{x} \frac{\partial u_x}{\partial x} \frac{1}{r} \frac{\partial \hat{x}}{\partial \theta}$$

$$+ \hat{\theta} \cdot \left(\hat{r} \frac{1}{r} \frac{\partial}{\partial \theta} \frac{\partial u_x}{\partial r} + \frac{\partial \hat{r}}{\partial \theta} \frac{1}{r} \frac{\partial u_x}{\partial r}\right) \hat{x} + \hat{\theta} \cdot \hat{r} \frac{\partial u_x}{\partial r} \frac{1}{r} \frac{\partial \hat{x}}{\partial \theta}$$

$$+ \hat{\theta} \cdot \left(\hat{\theta} \frac{1}{r} \frac{\partial}{\partial \theta} \left(\frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) \hat{x} + \frac{\partial \hat{\theta}}{\partial \theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) \hat{x} + \hat{\theta} \cdot \hat{\theta} \cdot \hat{\theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta}$$

$$+ \hat{\theta} \cdot \left(\hat{\theta} \frac{1}{r} \frac{\partial}{\partial \theta} \left(\frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) \hat{x} + \frac{\partial \hat{\theta}}{\partial \theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta}\right) \hat{x} + \hat{\theta} \cdot \hat{\theta} \cdot \hat{\theta} \cdot \hat{\theta} \frac{1}{r} \frac{\partial u_x}{\partial \theta}$$

$$= \left[\frac{\partial^2 u_x}{\partial x^2} + \frac{\partial^2 u_x}{\partial r^2} + \frac{1}{r} \frac{\partial^2 u_x}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u_x}{\partial \theta^2}\right] \hat{x}$$
(10)

Second column:

Third column:

$$\therefore \nabla \cdot \nabla (\boldsymbol{u}) = \begin{cases}
\left[\frac{\partial^2 u_x}{\partial x^2} + \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial u_x}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u_x}{\partial \theta^2} \right] \hat{x} + \\
\left[\frac{\partial^2 u_r}{\partial x^2} + \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial u_r}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u_r}{\partial \theta^2} - \frac{u_r}{r^2} - \frac{2}{r^2} \frac{\partial u_\theta}{\partial \theta} \right] \hat{r} + \\
\left[\frac{\partial^2 u_\theta}{\partial x^2} + \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial u_\theta}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u_\theta}{\partial \theta^2} + \frac{2}{r^2} \frac{\partial u_r}{\partial \theta} - \frac{u_\theta}{r^2} \right] \hat{\theta}
\end{cases} \tag{13}$$

Which can be written using the Laplacian of individual scalar fields, u_x, u_r, u_θ as,

$$\therefore \nabla \cdot \nabla (\boldsymbol{u}) = \nabla^2(u_x)\hat{x} + \left(\nabla^2(u_r) - \frac{u_r}{r^2} - \frac{2}{r^2}\frac{\partial u_\theta}{\partial \theta}\right)\hat{r} + \left(\nabla^2(u_\theta) + \frac{2}{r^2}\frac{\partial u_r}{\partial \theta} - \frac{u_\theta}{r^2}\right)\hat{\theta}$$
(14)

References