

Tutorial 2

EC3301 - Martinmas Semester

Hedonic regression models

In this section we examine the an example of multivariate regression model: the hedonic regression model. These models are widely used in real estate to estimate the value of various characteristics of a house: its size, number of rooms, school access, neighbourhood quality, etc. Having decomposed the price, one can then predict the value of another house using its characteristics. In this case, we will make use a small dataset made available with the Wooldridge (2025, 131) textbook which includes the price and housing characteristic information for 506 nieghbourhoods (unit of observation) in the Boston area.

1. Consider the model below:

$$price_i = \beta_0 + \beta_1 rooms_i + \beta_2 proptax_i + \beta_3 dist_i + \beta_4 crime_i + \beta_5 statio_i + u_i$$

where,

- *price*: median housing price (USD)
- *rooms*: average number of rooms
- *proptax*: property tax *per \$1000*
- *dist*: weighted distance (in miles) to 5 employment centers
- *crime*: crimes committed in last year (per capita)
- *statio*: average student-teacher ratio of schools in the area

Provide an interpretation for each coefficient and predict its sign. For *proptax*, think about a \$100 increase in tax.

2. How does the interpretation of β_1 , β_2 , and β_3 change in the model below:

$$\ln(price_i) = \beta_0 + \beta_1 rooms_i + \beta_2 \ln(proptax_i) + \beta_3 \ln(dist_i) + \beta_4 crime_i + \beta_5 statio_i + u_i$$

3. How would this interpretation of β_3 from the above equation change if *proptax* was measured in \$ and not \$ per \$1000? And β_4 if *crime* was measured in crimes committed per 100 people. Would these changes affect the other coefficients in the model?

The following questions relate to the regression output below:

```
reg lprice rooms lproptax ldist crime stratio
```

Source	SS	df	MS	Number of obs	=	506
Model	52.9136182	5	10.5827236	F(5, 500)	=	167.09
Residual	31.6686068	500	.063337214	Prob > F	=	0.0000
				R-squared	=	0.6256
				Adj R-squared	=	0.6218
Total	84.582225	505	.167489554	Root MSE	=	.25167

lprice	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
rooms	.2633338	.0174317	15.11	0.000	.2290854	.2975821
lproptax	-.2005414	.0403609	-4.97	0.000	-.2798393	-.1212435
ldist	.0043865	.0266419	0.16	0.869	-.0479574	.0567304
crime	-.0127542	.001601	-7.97	0.000	-.0158996	-.0096087
stratio	-.0334352	.0059361	-5.63	0.000	-.0450979	-.0217725
_cons	10.13379	.2869904	35.31	0.000	9.569931	10.69764

4. Provide an accurate interpretation of the **rooms** coefficient (i.e. semi-elasticity).
5. Provide an interpretation of the estimated **ldist** and **crime** coefficients.
6. What is the predicted (median) house price of a 4 bedroom home, subject to a 30,000 property tax, 1 (weighted) mile from 5 economic centers, in a neighbourhood that reports 20 crimes per capita and a student-teacher ratio 16. [You can apply coefficients to the nearest 3 decimal points]

Out of date

The data comes a rather old study by Harrison Jr and Rubinfeld (1978) titled *Hedonic housing prices and the demand for clean air*. The answer you get for this question will not seem realistic given the housing market today. The mean of the **price** variable (which captures median house price in a neighbourhood) is \$22,512. That would not be enough to buy a new car in the US today.

7. Test the null hypothesis $H_0 : \beta_5 \geq -0.025$ at a significance level of 0.05. You will need the following information on $t_{n-k-1}^{-1}(0.05)$

```
dis invt(e(df_r),0.05)
```

-1.6479069

8. Replicate the 95% CI interval for `ldist` shown above. You need the following information on $t_{n-k-1}^{-1}(0.975)$,

```
dis invt(e(df_r),0.975)
```

1.9647198

9. Replicate the reported $\mathbf{R}^2$ using the information reported above under ‘Source’. [Hint: ‘SS’ stands for Sum of Squares.]
10. Replicate the reported F -statistic (‘F(5, 500)’) using the reported $\mathbf{R}^2$. Is the model statistically significant?

Note

Before the next two questions, I will store some relevant Stata output from the first model:

```
scalar r2_ur = e(r2)
scalar ssr_ur = e(rss)
scalar df_ur = e(df_r)
```

This will be wiped from memory after estimating a different model.

11. Test the $H_0 : \beta_{\ln(dist)} = \beta_{\ln(proptax)} = 0$ using the reported information below. [The code the $\mathbf{R}^2$ from a model that excludes `ldist` and `lproptax`.]

```
qui reg lprice rooms crime stratio
dis e(r2)
```

.60187703

12. Test the $H_0 : \beta_{\ln(dist)} = 0$ using the reported information below. [The code the SSR from a model that excludes `ldist`.] Compute the relevant p-value for this test. What is the relationship between the F -statistic computed here and the t -statistic shown in the original output.

```
qui reg lprice rooms lproptax crime stratio
dis e(rss)
```

31.670324

Properties of OLS

Consider the following decomposition of y_i based upon the OLS estimator,

$$y_i = \hat{\beta}_0 + \hat{\beta}_1 x_{1i} + \hat{\beta}_2 x_{2i} + \hat{u}_i$$

corresponding to the multivariate model

$$y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + u_i$$

1. Under MLR. 1-4, you can show that $E[\hat{\beta}_1] = \beta_1$. What is this property called and what does it mean?
2. In [Lecture 3](#), we defined

$$se(\hat{\beta}_1) = \frac{\hat{\sigma}}{\sqrt{n}sd(x_1)\sqrt{1 - R_1^2}}$$

Explain each component of this equation and how you would compute it.

3. What happens to the variance of the estimator if x_1 and x_2 are highly correlated?

! Multicollinearity

In the limit, you can consider what happens as $R_j^2 \rightarrow 1$. This is the problem of multicollinearity discussed on p. 91 of Wooldridge (2025).

4. How does the equation above relate to the material from [Lecture 2](#) on partialling out (i.e. Frisch-Waugh Theorem)?
5. Would the problem of correlated regressors be a problem in the evaluation of a Randomized Control Trial? As we saw in the regression from Karlan, Rigol, and Roth (2026, 24) (in [Lecture 1](#)), authors often estimate the impact of treatment in a randomized control trial by estimating a linear regression model with additional controls.

$$y_i = \alpha + \beta T_i + \mathbf{x}_i \delta + \varepsilon_i$$

6. What would the impact of an increase in sample size be?
7. Randomized control trials are famously low powered. Given the discussion above, what are some of the ways that researchers might increase the power of their test to reject the null hypothesis of no effect ($H_0 : \beta = 0$).
8. If the error term in the underlying model is heteroskedastic, will the estimated variance be smaller or larger?
9. When the error variance is not homoskedastic, why can you no longer use the t or F distribution? [This is an advanced question, not covered by material in the lectures. See what you can find out.]

Harrison Jr, David, and Daniel L Rubinfeld. 1978. “Hedonic Housing Prices and the Demand for Clean Air.” *Journal of Environmental Economics and Management* 5 (1): 81–102.

Karlan, Dean, Natalia Rigol, and Benjamin N. Roth. 2026. “When Microenterprises Grow, Are Consumers Better Off? Evidence from Large Loans to Microenterprises in Chile.” Working Paper 35729. Working Paper Series. National Bureau of Economic Research. <https://doi.org/10.3386/w35729>.

Wooldridge, Jeffrey M. 2025. “Introductory Econometrics: A Modern Approach.”