

Lecture 6

- Coordinates and Basis
- Dimension
- Change of Basis
- Row Space, Column Space, and Null Space



Coordinates and Basis

Definition

A vector space V is said to be *finite-dimensional* if there is a finite set of vectors in V that spans V and is said to be *infinite-dimensional* if no such set exists.

Coordinates and Basis

Definition

A vector space V is said to be *finite-dimensional* if there is a finite set of vectors in V that spans V and is said to be *infinite-dimensional* if no such set exists.

Definition

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a set of vectors in a finite-dimensional vector space V , then S is called a *basis* for V if:

- 1 S spans V ;¹ and
- 2 S is linearly independent.

(The plural of basis is bases.)

¹i.e. if $\text{span } S = V$.

$$\mathbf{i} = (1, 0, 0) \quad \mathbf{j} = (0, 1, 0) \quad \mathbf{k} = (0, 0, 1)$$



Example

Recall from last week that the standard unit vectors

$$\mathbf{e}_1 = (1, 0, 0, \dots, 0), \mathbf{e}_2 = (0, 1, 0, \dots, 0), \dots, \mathbf{e}_n = (0, 0, 0, \dots, 1).$$

span \mathbb{R}^n . These vectors are linearly independent (I only proved it for $n = 3$.)

$$\mathbf{i} = (1, 0, 0) \quad \mathbf{j} = (0, 1, 0) \quad \mathbf{k} = (0, 0, 1)$$



Example

Recall from last week that the standard unit vectors

$$\mathbf{e}_1 = (1, 0, 0, \dots, 0), \mathbf{e}_2 = (0, 1, 0, \dots, 0), \dots, \mathbf{e}_n = (0, 0, 0, \dots, 1).$$

span \mathbb{R}^n . These vectors are linearly independent (I only proved it for $n = 3$.)

Therefore these vectors form a basis for \mathbb{R}^n that we call the *standard basis for \mathbb{R}^n* .

$$\mathbf{i} = (1, 0, 0) \quad \mathbf{j} = (0, 1, 0) \quad \mathbf{k} = (0, 0, 1)$$



Example

Recall from last week that the standard unit vectors

$$\mathbf{e}_1 = (1, 0, 0, \dots, 0), \mathbf{e}_2 = (0, 1, 0, \dots, 0), \dots, \mathbf{e}_n = (0, 0, 0, \dots, 1).$$

span \mathbb{R}^n . These vectors are linearly independent (I only proved it for $n = 3$.)

Therefore these vectors form a basis for \mathbb{R}^n that we call the *standard basis for \mathbb{R}^n* .

Example

$S = \{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$ is called the *standard basis for \mathbb{R}^3* .

Coordinates and Basis



Example (The Standard Basis for \mathbb{P}^n)

Show that $S = \{1, x, x^2, \dots, x^n\}$ is a basis for the vector space \mathbb{P}^n of polynomials of degree n or less.

Coordinates and Basis



Example (The Standard Basis for \mathbb{P}^n)

Show that $S = \{1, x, x^2, \dots, x^n\}$ is a basis for the vector space \mathbb{P}^n of polynomials of degree n or less.

Last week we showed that $\text{span } S = \mathbb{P}^n$ and that S is linearly independent. Therefore S is a basis for \mathbb{P}^n .

Coordinates and Basis

Example (Another Basis for \mathbb{R}^3)

Show that the vectors $\mathbf{v}_1 = (1, 2, 1)$, $\mathbf{v}_2 = (2, 9, 0)$, and $\mathbf{v}_3 = (3, 3, 4)$ form a basis for \mathbb{R}^3 .

We need to prove two things:

- 1 These three vectors are linearly independent; and
- 2 they span \mathbb{R}^3 .

Coordinates and Basis

Example (Another Basis for \mathbb{R}^3)

Show that the vectors $\mathbf{v}_1 = (1, 2, 1)$, $\mathbf{v}_2 = (2, 9, 0)$, and $\mathbf{v}_3 = (3, 3, 4)$ form a basis for \mathbb{R}^3 .

We need to prove two things:

- 1** These three vectors are linearly independent; and
- 2** they span \mathbb{R}^3 .

For **1** we must show that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 = \mathbf{0}$$

has only the trivial solution.

Coordinates and Basis

Example (Another Basis for \mathbb{R}^3)

Show that the vectors $\mathbf{v}_1 = (1, 2, 1)$, $\mathbf{v}_2 = (2, 9, 0)$, and $\mathbf{v}_3 = (3, 3, 4)$ form a basis for \mathbb{R}^3 .

We need to prove two things:

- 1** These three vectors are linearly independent; and
- 2** they span \mathbb{R}^3 .

For **1** we must show that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 = \mathbf{0}$$

has only the trivial solution. For **2**, we must show that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 = \mathbf{b}$$

is consistent for every $\mathbf{b} \in \mathbb{R}^3$.

Coordinates and Basis



So we have two linear systems to consider:

$$\begin{cases} c_1 + 2c_2 + 3c_3 = 0 \\ 2c_1 + 9c_2 + 3c_3 = 0 \\ c_1 + 4c_3 = 0 \end{cases} \quad \text{and} \quad \begin{cases} c_1 + 2c_2 + 3c_3 = b_1 \\ 2c_1 + 9c_2 + 3c_3 = b_2 \\ c_1 + 4c_3 = b_3. \end{cases}$$

Coordinates and Basis



So we have two linear systems to consider:

$$\begin{cases} c_1 + 2c_2 + 3c_3 = 0 \\ 2c_1 + 9c_2 + 3c_3 = 0 \\ c_1 + 4c_3 = 0 \end{cases} \quad \text{and} \quad \begin{cases} c_1 + 2c_2 + 3c_3 = b_1 \\ 2c_1 + 9c_2 + 3c_3 = b_2 \\ c_1 + 4c_3 = b_3. \end{cases}$$

These two linear systems have the same coefficient matrix

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 9 & 3 \\ 1 & 0 & 4 \end{bmatrix}.$$

If we can show that $\det(A) \neq 0$, then we can prove both things at the same time.

Coordinates and Basis



If $\det(A) \neq 0$, then

$$A\mathbf{c} = \mathbf{0} \implies \mathbf{c} = A^{-1}\mathbf{0} = \mathbf{0}$$

and

$$A\mathbf{c} = \mathbf{b} \implies \mathbf{c} = A^{-1}\mathbf{b}.$$

Coordinates and Basis



If $\det(A) \neq 0$, then

$$A\mathbf{c} = \mathbf{0} \implies \mathbf{c} = A^{-1}\mathbf{0} = \mathbf{0}$$

and

$$A\mathbf{c} = \mathbf{b} \implies \mathbf{c} = A^{-1}\mathbf{b}.$$

I leave it for you to prove that $\det(A) = -1$. Hence $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ form a basis for \mathbb{R}^3 .

Coordinates and Basis

Example (The Standard Basis for $\mathbb{R}^{m \times n} = M_{mn}$)

Show that the matrices

$$M_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad M_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad M_4 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

form a basis for the vector space $\mathbb{R}^{2 \times 2}$ of 2×2 matrices.

Coordinates and Basis

Example (The Standard Basis for $\mathbb{R}^{m \times n} = M_{mn}$)

Show that the matrices

$$M_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad M_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad M_4 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

form a basis for the vector space $\mathbb{R}^{2 \times 2}$ of 2×2 matrices.

$$c_1 M_1 + c_2 M_2 + c_3 M_3 + c_4 M_4 = \mathbf{0}$$

$$c_1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + c_4 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Coordinates and Basis

Example (The Standard Basis for $\mathbb{R}^{m \times n} = M_{mn}$)

Show that the matrices

$$M_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad M_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad M_4 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

form a basis for the vector space $\mathbb{R}^{2 \times 2}$ of 2×2 matrices.

$$c_1 M_1 + c_2 M_2 + c_3 M_3 + c_4 M_4 = \mathbf{0}$$

$$\begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix} = c_1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + c_4 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Coordinates and Basis

Example (The Standard Basis for $\mathbb{R}^{m \times n} = M_{mn}$)

Show that the matrices

$$M_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad M_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad M_4 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

form a basis for the vector space $\mathbb{R}^{2 \times 2}$ of 2×2 matrices.

$$c_1 M_1 + c_2 M_2 + c_3 M_3 + c_4 M_4 = \mathbf{0}$$

$$\begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix} = c_1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + c_4 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Hence

$$c_1 = c_2 = c_3 = c_4 = 0$$

which proves that these four matrices are linearly independent.

Coordinates and Basis

Let $B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be any matrix in $\mathbb{R}^{2 \times 2}$.

$$c_1 M_1 + c_2 M_2 + c_3 M_3 + c_4 M_4 = B$$

$$c_1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + c_4 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Coordinates and Basis

Let $B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be any matrix in $\mathbb{R}^{2 \times 2}$.

$$c_1 M_1 + c_2 M_2 + c_3 M_3 + c_4 M_4 = B$$

$$\begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix} = c_1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + c_4 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Coordinates and Basis

Let $B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be any matrix in $\mathbb{R}^{2 \times 2}$.

$$c_1 M_1 + c_2 M_2 + c_3 M_3 + c_4 M_4 = B$$

$$\begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix} = c_1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + c_4 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Hence

$$\begin{cases} c_1 = a \\ c_2 = b \\ c_3 = c \\ c_4 = d. \end{cases}$$

This proves that these four matrices span $\mathbb{R}^{2 \times 2}$.

Therefore M_1, M_2, M_3 and M_4 for a basis for $\mathbb{R}^{2 \times 2}$ called the *standard basis for $\mathbb{R}^{2 \times 2} = M_{22}$* .

Coordinates and Basis

Example

Show that the vector space \mathbb{P} is infinite dimensional².

We need to show that \mathbb{P} does not have a finite spanning set.

² \mathbb{P} = {all polynomials with real coefficients}

Coordinates and Basis

Example

Show that the vector space \mathbb{P} is infinite dimensional².

We need to show that \mathbb{P} does not have a finite spanning set.

If there were a finite spanning set, say $S = \{\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_r\}$, then the degrees of the polynomials in S would have a maximum value, say n ;

² $\mathbb{P} = \{\text{all polynomials with real coefficients}\}$

Coordinates and Basis

Example

Show that the vector space \mathbb{P} is infinite dimensional².

We need to show that \mathbb{P} does not have a finite spanning set.

If there were a finite spanning set, say $S = \{\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_r\}$, then the degrees of the polynomials in S would have a maximum value, say n ; and this in turn would imply that any linear combination of the polynomials in S would have degree at most n .

² $\mathbb{P} = \{\text{all polynomials with real coefficients}\}$

Coordinates and Basis

Example

Show that the vector space \mathbb{P} is infinite dimensional².

We need to show that \mathbb{P} does not have a finite spanning set.

If there were a finite spanning set, say $S = \{\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_r\}$, then the degrees of the polynomials in S would have a maximum value, say n ; and this in turn would imply that any linear combination of the polynomials in S would have degree at most n .

Thus, there would be no way to express the polynomial x^{n+1} as a linear combination of the polynomials in S , contradicting the fact that the vectors in S span \mathbb{P} .

² $\mathbb{P} = \{\text{all polynomials with real coefficients}\}$

Coordinates and Basis

Example

Some other infinite dimensional vector spaces are

$$\mathbb{R}^{\mathbb{N}} = \{\text{all sequences of real numbers}\}$$

$$F(-\infty, \infty) = \{\text{all functions } f : \mathbb{R} \rightarrow \mathbb{R}\}$$

$$C(-\infty, \infty) = \{\text{all continuous functions } f : \mathbb{R} \rightarrow \mathbb{R}\}$$

$$C^k(-\infty, \infty) = \left\{ \begin{array}{l} \text{all functions } f : \mathbb{R} \rightarrow \mathbb{R} \text{ which are} \\ \text{continuously differentiable } k \text{ times} \end{array} \right\}$$

$$C^\infty(-\infty, \infty) = \left\{ \begin{array}{l} \text{all functions } f : \mathbb{R} \rightarrow \mathbb{R} \text{ which} \\ \text{can be differentiated } \infty \text{ times} \end{array} \right\}$$

Coordinates Relative to a Basis

Theorem

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a basis for a vector space V , then every vector \mathbf{v} in V can be expressed in the form

$$\mathbf{v} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_n\mathbf{v}_n$$

in exactly one way.

Now we can start talking about coordinates in a general vector space V .

Coordinates and Basis

Definition

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a basis for a vector space V , and

$$\mathbf{v} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_n\mathbf{v}_n$$

is the expression for a vector \mathbf{v} in terms of the basis S , then the scalars c_1, c_2, \dots, c_n are called the *coordinates* of \mathbf{v} relative to the basis S .

Coordinates and Basis

Definition

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a basis for a vector space V , and

$$\mathbf{v} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_n\mathbf{v}_n$$


is the expression for a vector \mathbf{v} in terms of the basis S , then the scalars c_1, c_2, \dots, c_n are called the *coordinates* of \mathbf{v} relative to the basis S . The vector (c_1, c_2, \dots, c_n) in \mathbb{R}^n constructed from these coordinates is called the *coordinate vector of \mathbf{v} relative to S* ; it is denoted by

$$(\mathbf{v})_S = (c_1, c_2, \dots, c_n).$$

Remark

When we calculate coordinates, the order of the vectors in S is important. Some books use the term *ordered basis* which means a basis in which the order of the vectors is fixed.

Observe that $(\mathbf{v})_S$ is a vector in \mathbb{R}^n , so that once an ordered basis S is given for a vector space V , Theorem 4.4.1 establishes a one-to-one correspondence between vectors in V and vectors in \mathbb{R}^n (Figure 4.4.6).



$$\mathbf{i} = (1, 0, 0) \quad \mathbf{j} = (0, 1, 0) \quad \mathbf{k} = (0, 0, 1)$$



Example

$V = \mathbb{R}^3$. Standard basis $\mathbf{i}, \mathbf{j}, \mathbf{k}$.

If $\mathbf{v} = (a, b, c)$ is any vector in \mathbb{R}^3 , then

$$\mathbf{v} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k} \implies (\mathbf{v})_S = (a, b, c).$$

Coordinates and Basis



Example

Find the coordinate vector for the polynomial

$$\mathbf{p}(x) = c_0 + c_1x + c_2x^2 + \dots + c_nx^n$$

relative to the standard basis for the vector space \mathbb{P}^n .

The standard basis for \mathbb{P}^n is $S = \{1, x, x^2, \dots, x^n\}$. So clearly

$$(\mathbf{p})_S = (c_0, c_1, c_2, \dots, c_n).$$

Coordinates and Basis

Example

Find the coordinate vector of

$$B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

relative to the standard basis for $\mathbb{R}^{2 \times 2}$.

Since

$$B = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = a \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + d \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

we have

$$(B)_S = (a, b, c, d).$$

Coordinates and Basis

Example (Coordinates in \mathbb{R}^3)

We showed in a previous example that the vectors

$$\mathbf{v}_1 = (1, 2, 1), \quad \mathbf{v}_2 = (2, 9, 0), \quad \mathbf{v}_3 = (3, 3, 4)$$

form a basis for \mathbb{R}^3 .

- 1 Find the coordinate vector of $\mathbf{v} = (5, -1, 9)$ relative to the basis $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.
- 2 Find the vector \mathbf{u} in \mathbb{R}^3 whose coordinate vector relative to S is $(\mathbf{u})_S = (-1, 3, 2)$.

Coordinate

$$\mathbf{v}_1 = (1, 2, 1), \quad \mathbf{v}_2 = (2, 9, 0), \quad \mathbf{v}_3 = (3, 3, 4)$$

- 1 Find the coordinate vector of $\mathbf{v} = (5, -1, 9)$ relative to the basis $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

We need to find scalars c_1, c_2, c_3 such that

$$\mathbf{v} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3.$$

Coordinate basis $\mathbf{v}_1 = (1, 2, 1), \mathbf{v}_2 = (2, 9, 0), \mathbf{v}_3 = (3, 3, 4)$

- 1 Find the coordinate vector of $\mathbf{v} = (5, -1, 9)$ relative to the basis $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

We need to find scalars c_1, c_2, c_3 such that

$$\mathbf{v} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3.$$

I.e. such that

$$(5, -1, 9) = c_1(1, 2, 1) + c_2(2, 9, 0) + c_3(3, 3, 4).$$

Coordinate basis $\mathbf{v}_1 = (1, 2, 1), \mathbf{v}_2 = (2, 9, 0), \mathbf{v}_3 = (3, 3, 4)$

- 1 Find the coordinate vector of $\mathbf{v} = (5, -1, 9)$ relative to the basis $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

We need to find scalars c_1, c_2, c_3 such that

$$\mathbf{v} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3.$$

I.e. such that

$$(5, -1, 9) = c_1(1, 2, 1) + c_2(2, 9, 0) + c_3(3, 3, 4).$$

So we need to solve the linear system

$$\begin{cases} c_1 + 2c_2 + 3c_3 = 5 \\ 2c_1 + 9c_2 + 3c_3 = -1 \\ c_1 + 4c_3 = 9. \end{cases}$$

Coordinate basis $\mathbf{v}_1 = (1, 2, 1), \mathbf{v}_2 = (2, 9, 0), \mathbf{v}_3 = (3, 3, 4)$

- 1 Find the coordinate vector of $\mathbf{v} = (5, -1, 9)$ relative to the basis $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

We need to find scalars c_1, c_2, c_3 such that

$$\mathbf{v} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3.$$

I.e. such that

$$(5, -1, 9) = c_1(1, 2, 1) + c_2(2, 9, 0) + c_3(3, 3, 4).$$

So we need to solve the linear system

$$\begin{cases} c_1 + 2c_2 + 3c_3 = 5 \\ 2c_1 + 9c_2 + 3c_3 = -1 \\ c_1 + 4c_3 = 9. \end{cases}$$

I leave it to you to check that the solution is $c_1 = 1, c_2 = -1$ and $c_3 = 2$. Therefore

$$(\mathbf{v})_S = (1, -1, 2).$$

Coordinate basis

$$\mathbf{v}_1 = (1, 2, 1), \quad \mathbf{v}_2 = (2, 9, 0), \quad \mathbf{v}_3 = (3, 3, 4)$$

- 2 Find the vector \mathbf{u} in \mathbb{R}^3 whose coordinate vector relative to S is $(\mathbf{u})_S = (-1, 3, 2)$.

This one is easier.

$$\mathbf{u} = (-1)\mathbf{v}_1 + 3\mathbf{v}_2 + 2\mathbf{v}_3$$

=

=

Coordinate basis

$$\mathbf{v}_1 = (1, 2, 1), \quad \mathbf{v}_2 = (2, 9, 0), \quad \mathbf{v}_3 = (3, 3, 4)$$

- 2 Find the vector \mathbf{u} in \mathbb{R}^3 whose coordinate vector relative to S is $(\mathbf{u})_S = (-1, 3, 2)$.

This one is easier.

$$\begin{aligned}\mathbf{u} &= (-1)\mathbf{v}_1 + 3\mathbf{v}_2 + 2\mathbf{v}_3 \\ &= (-1)(1, 2, 1) + 3(2, 9, 0) + 2(3, 3, 4) \\ &= (11, 31, 7).\end{aligned}$$



Dimension

We have seen two bases for \mathbb{R}^3 .

$$\mathbf{i} = (1, 0, 0), \quad \mathbf{j} = (0, 1, 0), \quad \mathbf{k} = (0, 0, 1)$$

$$\mathbf{v}_1 = (1, 2, 1), \quad \mathbf{v}_2 = (2, 9, 0), \quad \mathbf{v}_3 = (3, 3, 4)$$

Dimension



We have seen two bases for \mathbb{R}^3 .

$$\mathbf{i} = (1, 0, 0), \quad \mathbf{j} = (0, 1, 0), \quad \mathbf{k} = (0, 0, 1)$$

$$\mathbf{v}_1 = (1, 2, 1), \quad \mathbf{v}_2 = (2, 9, 0), \quad \mathbf{v}_3 = (3, 3, 4)$$

Dimension



We have seen two bases for \mathbb{R}^3 .

$$\mathbf{i} = (1, 0, 0),$$



$$\mathbf{j} = (0, 1, 0),$$



$$\mathbf{k} = (0, 0, 1)$$



$$\mathbf{v}_1 = (1, 2, 1),$$



$$\mathbf{v}_2 = (2, 9, 0),$$



$$\mathbf{v}_3 = (3, 3, 4)$$



Dimension

Theorem

All bases for a finite-dimensional vector space have the same number of vectors.

Definition

The dimension of a finite-dimensional vector space V , denoted by $\dim(V)$, is defined to be the number of vectors in a basis for V .

The dimension of the zero vector space $\{\mathbf{0}\}$ is defined to be 0.

Dimension

Theorem

All bases for a finite-dimensional vector space have the same number of vectors.

Definition

The dimension of a finite-dimensional vector space V , denoted by $\dim(V)$, is defined to be the number of vectors in a basis for V .

The dimension of the zero vector space $\{\mathbf{0}\}$ is defined to be 0.

Engineers often use the term *degrees of freedom* as a synonym for dimension.

Theorem

Let V be an n -dimensional vector space, and let $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be any basis.

- 1 If a set in V has more than n vectors, then it is linearly dependent.
- 2 If a set in V has fewer than n vectors, then it does not span V .

Example

- $\dim(\mathbb{R}^n) = n$ because the standard basis has n vectors in it.

Example

- $\dim(\mathbb{R}^n) = n$ because the standard basis has n vectors in it.
- $\dim(\mathbb{P}^n) = n + 1$. Recall that the standard basis for \mathbb{P}^n is $\{1, x, x^2, \dots, x^n\}$.

Dimension

Example

- $\dim(\mathbb{R}^n) = n$ because the standard basis has n vectors in it.
- $\dim(\mathbb{P}^n) = n + 1$. Recall that the standard basis for \mathbb{P}^n is $\{1, x, x^2, \dots, x^n\}$.
- $\dim(\mathbb{R}^{2 \times 2}) = 4$ because the standard basis contains the following four vectors

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

Dimension

Example

- $\dim(\mathbb{R}^n) = n$ because the standard basis has n vectors in it.
- $\dim(\mathbb{P}^n) = n + 1$. Recall that the standard basis for \mathbb{P}^n is $\{1, x, x^2, \dots, x^n\}$.
- $\dim(\mathbb{R}^{2 \times 2}) = 4$ because the standard basis contains the following four vectors

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

- Similarly $\dim(\mathbb{R}^{m \times n}) = mn$.

► EXAMPLE 2 Dimension of $\text{Span}(S)$

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ then every vector in $\text{span}(S)$ is expressible as a linear combination of the vectors in S . Thus, if the vectors in S are *linearly independent*, they automatically form a basis for $\text{span}(S)$, from which we can conclude that

$$\dim[\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}] = r$$

In words, the dimension of the space spanned by a linearly independent set of vectors is equal to the number of vectors in that set.

Example (Dimension of a Solution Space)

Find a basis for and the dimension of the solution space of the homogeneous linear system

$$\begin{cases} x_1 + 3x_2 - 2x_3 + 2x_5 = 0 \\ 2x_1 + 6x_2 - 5x_3 - 2x_4 + 4x_5 - 3x_6 = 0 \\ 5x_3 + 10x_4 + 15x_6 = 0 \\ 2x_1 + 6x_2 + 8x_4 + 4x_5 + 18x_6 = 0. \end{cases}$$

Dimension

I leave it to you to check that the solution is

$$\begin{cases} x_1 = -3x_2 - 4x_4 - 2x_5 \\ x_2 \text{ is free} \\ x_3 = -2x_4 \\ x_4 \text{ is free} \\ x_5 \text{ is free} \\ x_6 = 0. \end{cases}$$

Dimension

I leave it to you to check that the solution is

$$\begin{cases} x_1 = -3x_2 - 4x_4 - 2x_5 \\ x_2 \text{ is free} \\ x_3 = -2x_4 \\ x_4 \text{ is free} \\ x_5 \text{ is free} \\ x_6 = 0. \end{cases}$$

We can write this as

$$\begin{aligned} (x_1, x_2, x_3, x_4, x_5, x_6) &= (-3x_2 - 4x_4 - 2x_5, x_2, -2x_4, x_4, x_5, 0) \\ &= x_2(-3, 1, 0, 0, 0, 0) + x_4(-4, 0, -2, 1, 0, 0) \\ &\quad + x_5(-2, 0, 0, 0, 1, 0). \end{aligned}$$

Dimension



This shows that the vectors

$$\mathbf{v}_1 = (-3, 1, 0, 0, 0, 0), \mathbf{v}_2 = (-4, 0, -2, 1, 0, 0), \mathbf{v}_3 = (-2, 0, 0, 0, 1, 0)$$

span the solution space.

This shows that the vectors

$$\mathbf{v}_1 = (-3, 1, 0, 0, 0, 0), \mathbf{v}_2 = (-4, 0, -2, 1, 0, 0), \mathbf{v}_3 = (-2, 0, 0, 0, 1, 0)$$

span the solution space. I leave it to you to check that these vectors are linearly independent.

This shows that the vectors

$$\mathbf{v}_1 = (-3, 1, 0, 0, 0, 0), \mathbf{v}_2 = (-4, 0, -2, 1, 0, 0), \mathbf{v}_3 = (-2, 0, 0, 0, 1, 0)$$

span the solution space. I leave it to you to check that these vectors are linearly independent.

Therefore the solution space has dimension 3.

Some More Theorems

Theorem (The Basis Theorem³)

Let V be an n -dimensional vector space, and let S be a set in V with exactly n vectors. Then the following are equivalent:

- 1 S is a basis for V ;
- 2 S spans V ;
- 3 S is linearly independent.

³page 245 in your textbook

► EXAMPLE 5 Bases by Inspection

- (a) Explain why the vectors $\mathbf{v}_1 = (-3, 7)$ and $\mathbf{v}_2 = (5, 5)$ form a basis for R^2 .
- (b) Explain why the vectors $\mathbf{v}_1 = (2, 0, -1)$, $\mathbf{v}_2 = (4, 0, 7)$, and $\mathbf{v}_3 = (-1, 1, 4)$ form a basis for R^3 .

Solution (a) Since neither vector is a scalar multiple of the other, the two vectors form a linearly independent set in the two-dimensional space R^2 , and hence they form a basis by Theorem 4.5.4.

Solution (b) The vectors \mathbf{v}_1 and \mathbf{v}_2 form a linearly independent set in the xz -plane (why?). The vector \mathbf{v}_3 is outside of the xz -plane, so the set $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is also linearly independent. Since R^3 is three-dimensional, Theorem 4.5.4 implies that $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is a basis for the vector space R^3 . ◀

Theorem

Let S be a finite set of vectors in a finite-dimensional vector space V .

- 1 If S spans V but is not a basis for V , then S can be reduced to a basis for V by removing appropriate vectors from S .
- 2 If S is a linearly independent set that is not already a basis for V , then S can be enlarged to a basis for V by inserting appropriate vectors into S .

Theorem

If W is a subspace of a finite-dimensional vector space V , then:

- 1** W is finite-dimensional;
- 2** $\dim(W) \leq \dim(V)$; and
- 3** $W = V$ if and only if $\dim(W) = \dim(V)$.





Change of Basis

Change of Basis

Consider the vector space $V = \mathbb{R}^2$. Suppose that $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$ are two bases for V .



Change of Basis

Consider the vector space $V = \mathbb{R}^2$. Suppose that $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$ are two bases for V .





Change of Basis

Consider the vector space $V = \mathbb{R}^2$. Suppose that $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$ are two bases for V .




Change of Basis

Consider the vector space $V = \mathbb{R}^2$. Suppose that $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$ are two bases for V .




Change of Basis

Consider the vector space $V = \mathbb{R}^2$. Suppose that $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$ are two bases for V .




$$\mathbf{x} = 3\mathbf{b}_1 + \mathbf{b}_2$$

$$(\mathbf{x})_{\mathcal{B}} = (3, 1) \quad [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$




Change of Basis

Consider the vector space $V = \mathbb{R}^2$. Suppose that $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$ are two bases for V .



$$\mathbf{x} = 3\mathbf{b}_1 + \mathbf{b}_2$$

$$(\mathbf{x})_{\mathcal{B}} = (3, 1) \quad [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$



$$\mathbf{x} = 6\mathbf{c}_1 + 4\mathbf{c}_2$$

$$(\mathbf{x})_{\mathcal{C}} = (6, 4) \quad [\mathbf{x}]_{\mathcal{C}} = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$$

Change of Basis



$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \xrightarrow{\text{HOW?}} [\mathbf{x}]_{\mathcal{C}} = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$$

Change of Basis



$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \xrightarrow{\text{HOW?}} [\mathbf{x}]_{\mathcal{C}} = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$$

Example

Suppose that $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$ are two bases for a vector space V . Suppose you know that

$$\mathbf{b}_1 = 4\mathbf{c}_1 + \mathbf{c}_2, \quad \mathbf{b}_2 = -6\mathbf{c}_1 + \mathbf{c}_2$$

and that the vector \mathbf{x} has coordinates $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$. Find $[\mathbf{x}]_{\mathcal{C}}$.

Change of Basis

Note that

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \implies \mathbf{x} = 3\mathbf{b}_1 + \mathbf{b}_2.$$

Hence

$$[\mathbf{x}]_{\mathcal{C}} = [3\mathbf{b}_1 + \mathbf{b}_2]_{\mathcal{C}} = 3 [\mathbf{b}_1]_{\mathcal{C}} + [\mathbf{b}_2]_{\mathcal{C}}$$

=

Change of Basis



Note that

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \implies \mathbf{x} = 3\mathbf{b}_1 + \mathbf{b}_2.$$

Hence

$$\begin{aligned} [\mathbf{x}]_{\mathcal{C}} &= [3\mathbf{b}_1 + \mathbf{b}_2]_{\mathcal{C}} = 3 [\mathbf{b}_1]_{\mathcal{C}} + [\mathbf{b}_2]_{\mathcal{C}} \\ &= \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix}. \end{aligned}$$

Change of Basis

Note that

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \quad \Rightarrow \quad \mathbf{x} = 3\mathbf{b}_1 + \mathbf{b}_2.$$

Hence

$$\begin{aligned} [\mathbf{x}]_{\mathcal{C}} &= [3\mathbf{b}_1 + \mathbf{b}_2]_{\mathcal{C}} = 3 [\mathbf{b}_1]_{\mathcal{C}} + [\mathbf{b}_2]_{\mathcal{C}} \\ &= \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix}. \end{aligned}$$

So what are $[\mathbf{b}_1]_{\mathcal{C}}$ and $[\mathbf{b}_2]_{\mathcal{C}}$?

Change of Basis

The question told us $\mathbf{b}_1 = 4\mathbf{c}_1 + \mathbf{c}_2$ and $\mathbf{b}_2 = -6\mathbf{c}_1 + \mathbf{c}_2$. Hence

$$[\mathbf{b}_1]_{\mathcal{C}} = \begin{bmatrix} 4 \\ 1 \end{bmatrix} \quad \text{and} \quad [\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} -6 \\ 1 \end{bmatrix}.$$

Change of Basis

The question told us $\mathbf{b}_1 = 4\mathbf{c}_1 + \mathbf{c}_2$ and $\mathbf{b}_2 = -6\mathbf{c}_1 + \mathbf{c}_2$. Hence

$$[\mathbf{b}_1]_{\mathcal{C}} = \begin{bmatrix} 4 \\ 1 \end{bmatrix} \quad \text{and} \quad [\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} -6 \\ 1 \end{bmatrix}.$$

Therefore

$$\begin{aligned} [\mathbf{x}]_{\mathcal{C}} &= [3\mathbf{b}_1 + \mathbf{b}_2]_{\mathcal{C}} = 3[\mathbf{b}_1]_{\mathcal{C}} + [\mathbf{b}_2]_{\mathcal{C}} \\ &= \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 & -6 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix}. \end{aligned}$$

Change of Basis

The question told us $\mathbf{b}_1 = 4\mathbf{c}_1 + \mathbf{c}_2$ and $\mathbf{b}_2 = -6\mathbf{c}_1 + \mathbf{c}_2$. Hence

$$[\mathbf{b}_1]_{\mathcal{C}} = \begin{bmatrix} 4 \\ 1 \end{bmatrix} \quad \text{and} \quad [\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} -6 \\ 1 \end{bmatrix}.$$

Therefore

$$\begin{aligned} [\mathbf{x}]_{\mathcal{C}} &= [3\mathbf{b}_1 + \mathbf{b}_2]_{\mathcal{C}} = 3[\mathbf{b}_1]_{\mathcal{C}} + [\mathbf{b}_2]_{\mathcal{C}} \\ &= \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 & -6 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix}. \end{aligned}$$

Remark

We did

$$[\mathbf{x}]_{\mathcal{C}} = \begin{bmatrix} 4 & -6 \\ 1 & 1 \end{bmatrix} [\mathbf{x}]_{\mathcal{B}}.$$

Change of Basis

Theorem

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ and $\mathcal{C} = \{\mathbf{c}_1, \dots, \mathbf{c}_n\}$ be bases of a vector space V . Then there is a unique $n \times n$ matrix $P_{\mathcal{C} \leftarrow \mathcal{B}}$ such that

$$[\mathbf{x}]_{\mathcal{C}} = P_{\mathcal{C} \leftarrow \mathcal{B}} [\mathbf{x}]_{\mathcal{B}}.$$

The columns of $P_{\mathcal{C} \leftarrow \mathcal{B}}$ are the \mathcal{C} -coordinate vectors of the vectors in the basis \mathcal{B} . That is,

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} & \cdots & [\mathbf{b}_n]_{\mathcal{C}} \end{bmatrix}.$$

Change of Basis



Remark

Some books write $P_{\mathcal{C} \leftarrow \mathcal{B}}$, other books write $P_{\mathcal{B} \rightarrow \mathcal{C}}$.

Change of Basis

Remark

Some books write $P_{\mathcal{C} \leftarrow \mathcal{B}}$, other books write $P_{\mathcal{B} \rightarrow \mathcal{C}}$.

Definition

The matrix $P_{\mathcal{C} \leftarrow \mathcal{B}}$ is called the *change-of-coordinates matrix from \mathcal{B} to \mathcal{C}* or the *transition matrix from \mathcal{B} to \mathcal{C}* .

Change of Basis



Remark

Some books write $P_{\mathcal{C} \leftarrow \mathcal{B}}$, other books write $P_{\mathcal{B} \rightarrow \mathcal{C}}$.

Definition

The matrix $P_{\mathcal{C} \leftarrow \mathcal{B}}$ is called the *change-of-coordinates matrix from \mathcal{B} to \mathcal{C}* or the *transition matrix from \mathcal{B} to \mathcal{C}* .

Theorem

The change-of-coordinates matrix $P_{\mathcal{C} \leftarrow \mathcal{B}}$ is invertible and

$$\left(P_{\mathcal{C} \leftarrow \mathcal{B}} \right)^{-1} = P_{\mathcal{B} \leftarrow \mathcal{C}}.$$



Break

We will continue at 3pm



Change of Basis



Example

Consider $V = \mathbb{R}^2$ with the bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$, where

$$\mathbf{b}_1 = \begin{bmatrix} -9 \\ 1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -5 \\ -1 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} 1 \\ -4 \end{bmatrix} \text{ and } \mathbf{c}_2 = \begin{bmatrix} 3 \\ -5 \end{bmatrix}.$$

Find the change-of-coordinates matrix from \mathcal{B} to \mathcal{C} .

Change of Basis

Example

Consider $V = \mathbb{R}^2$ with the bases $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$, where

$$\mathbf{b}_1 = \begin{bmatrix} -9 \\ 1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -5 \\ -1 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} 1 \\ -4 \end{bmatrix} \text{ and } \mathbf{c}_2 = \begin{bmatrix} 3 \\ -5 \end{bmatrix}.$$

Find the change-of-coordinates matrix from \mathcal{B} to \mathcal{C} .

Theorem

$$\begin{bmatrix} \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{b}_1 & \mathbf{b}_2 \end{bmatrix} \sim \begin{bmatrix} I & P_{\mathcal{C} \leftarrow \mathcal{B}} \end{bmatrix}.$$

Change of Basis



$$\begin{bmatrix} \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{b}_1 & \mathbf{b}_2 \end{bmatrix} = \begin{bmatrix} 1 & 3 & -9 & -5 \\ -4 & -5 & 1 & -1 \end{bmatrix}$$

Change of Basis

$$\begin{aligned}
 \begin{bmatrix} \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{b}_1 & \mathbf{b}_2 \end{bmatrix} &= \begin{bmatrix} 1 & 3 & -9 & -5 \\ -4 & -5 & 1 & -1 \end{bmatrix} \\
 &\sim \begin{bmatrix} 1 & 3 & -9 & -5 \\ 0 & 7 & -35 & -21 \end{bmatrix} \sim \begin{bmatrix} 1 & 3 & -9 & -5 \\ 0 & 1 & -5 & -3 \end{bmatrix} \\
 &\sim \begin{bmatrix} 1 & 0 & 6 & 4 \\ 0 & 1 & -5 & -3 \end{bmatrix} = \begin{bmatrix} I & P \\ \mathcal{C} \leftarrow \mathcal{B} \end{bmatrix}.
 \end{aligned}$$

Change of Basis

$$\begin{aligned}
 \begin{bmatrix} \mathbf{c}_1 & \mathbf{c}_2 & \mathbf{b}_1 & \mathbf{b}_2 \end{bmatrix} &= \begin{bmatrix} 1 & 3 & -9 & -5 \\ -4 & -5 & 1 & -1 \end{bmatrix} \\
 &\sim \begin{bmatrix} 1 & 3 & -9 & -5 \\ 0 & 7 & -35 & -21 \end{bmatrix} \sim \begin{bmatrix} 1 & 3 & -9 & -5 \\ 0 & 1 & -5 & -3 \end{bmatrix} \\
 &\sim \begin{bmatrix} 1 & 0 & 6 & 4 \\ 0 & 1 & -5 & -3 \end{bmatrix} = \begin{bmatrix} I & P_{\mathcal{C} \leftarrow \mathcal{B}} \end{bmatrix}.
 \end{aligned}$$

Therefore

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \begin{bmatrix} 6 & 4 \\ -5 & -3 \end{bmatrix}.$$

Change of Basis



Example

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$ and $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2\}$ be two bases for $V = \mathbb{R}^2$, where

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix} \quad \text{and} \quad \mathbf{c}_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}.$$

- 1** Find the change-of-coordinates matrix from \mathcal{C} to \mathcal{B} .
- 2** Find the change-of-coordinates matrix from \mathcal{B} to \mathcal{C} .

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}.$$



- 1 Note that we need to find $P_{\mathcal{B} \leftarrow \mathcal{C}}$ this time.

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}.$$

- 1 Note that we need to find $P_{\mathcal{B} \leftarrow \mathcal{C}}$ this time. Since

$$\begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \mathbf{c}_1 & \mathbf{c}_2 \end{bmatrix} = \begin{bmatrix} 1 & -2 & -7 & -5 \\ 3 & 4 & 9 & 7 \end{bmatrix}$$

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}.$$

- 1 Note that we need to find $P_{\mathcal{B} \leftarrow \mathcal{C}}$ this time. Since

$$\begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \mathbf{c}_1 & \mathbf{c}_2 \end{bmatrix} = \begin{bmatrix} 1 & -2 & -7 & -5 \\ 3 & 4 & 9 & 7 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 5 & 3 \\ 0 & 1 & 6 & 4 \end{bmatrix}$$

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}.$$

- 1 Note that we need to find $P_{\mathcal{B} \leftarrow \mathcal{C}}$ this time. Since

$$\begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \mathbf{c}_1 & \mathbf{c}_2 \end{bmatrix} = \begin{bmatrix} 1 & -2 & -7 & -5 \\ 3 & 4 & 9 & 7 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 5 & 3 \\ 0 & 1 & 6 & 4 \end{bmatrix}$$

we have that

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = \begin{bmatrix} 5 & 3 \\ 6 & 4 \end{bmatrix}.$$

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}.$$

- 1 Note that we need to find $P_{\mathcal{B} \leftarrow \mathcal{C}}$ this time. Since

$$\begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \mathbf{c}_1 & \mathbf{c}_2 \end{bmatrix} = \begin{bmatrix} 1 & -2 & -7 & -5 \\ 3 & 4 & 9 & 7 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 5 & 3 \\ 0 & 1 & 6 & 4 \end{bmatrix}$$

we have that

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = \begin{bmatrix} 5 & 3 \\ 6 & 4 \end{bmatrix}.$$

- 2 We calculate that

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \left(P_{\mathcal{B} \leftarrow \mathcal{C}} \right)^{-1} =$$

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}.$$

- 1 Note that we need to find $P_{\mathcal{B} \leftarrow \mathcal{C}}$ this time. Since

$$\begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \mathbf{c}_1 & \mathbf{c}_2 \end{bmatrix} = \begin{bmatrix} 1 & -2 & -7 & -5 \\ 3 & 4 & 9 & 7 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 5 & 3 \\ 0 & 1 & 6 & 4 \end{bmatrix}$$

we have that

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = \begin{bmatrix} 5 & 3 \\ 6 & 4 \end{bmatrix}.$$

- 2 We calculate that

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \left(P_{\mathcal{B} \leftarrow \mathcal{C}} \right)^{-1} = \begin{bmatrix} 5 & 3 \\ 6 & 4 \end{bmatrix}^{-1} =$$

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \quad \mathbf{c}_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}.$$

- 1 Note that we need to find $P_{\mathcal{B} \leftarrow \mathcal{C}}$ this time. Since

$$\begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \mathbf{c}_1 & \mathbf{c}_2 \end{bmatrix} = \begin{bmatrix} 1 & -2 & -7 & -5 \\ 3 & 4 & 9 & 7 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 5 & 3 \\ 0 & 1 & 6 & 4 \end{bmatrix}$$

we have that

$$P_{\mathcal{B} \leftarrow \mathcal{C}} = \begin{bmatrix} 5 & 3 \\ 6 & 4 \end{bmatrix}.$$

- 2 We calculate that

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \left(P_{\mathcal{B} \leftarrow \mathcal{C}} \right)^{-1} = \begin{bmatrix} 5 & 3 \\ 6 & 4 \end{bmatrix}^{-1} = \frac{1}{2} \begin{bmatrix} 4 & -3 \\ -6 & 5 \end{bmatrix} = \begin{bmatrix} 2 & -\frac{3}{2} \\ -3 & \frac{5}{2} \end{bmatrix}.$$



Cep telefonunuzu gözetmene teslim ediniz. Deposit your cell phones to an invigilator.

30 December 2019 [11:00-12:15]

MATH215 Mathematics III Final Exam

Page 3 of 4

3. [25 points] Let $\mathcal{A} = \{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ and $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be bases for the vector space V , and suppose $\mathbf{a}_1 = 4\mathbf{b}_1 - \mathbf{b}_2$, $\mathbf{a}_2 = -\mathbf{b}_1 + \mathbf{b}_2 + \mathbf{b}_3$, and $\mathbf{a}_3 = \mathbf{b}_2 - 2\mathbf{b}_3$.

- (a) Find the change of coordinates matrix from \mathcal{A} to \mathcal{B} .

Solution:

$$[\mathbf{a}_1]_{\mathcal{B}} = \begin{bmatrix} 4 \\ -1 \\ 0 \end{bmatrix}, \quad [\mathbf{a}_2]_{\mathcal{B}} = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}, \quad [\mathbf{a}_3]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix} \Rightarrow P = \begin{bmatrix} 4 & -1 & 0 \\ -1 & 1 & 1 \\ 0 & 1 & -2 \end{bmatrix}$$

- (b) Find $[\mathbf{x}]_{\mathcal{B}}$ for $\mathbf{x} = 3\mathbf{a}_1 + 4\mathbf{a}_2 + \mathbf{a}_3$.

Solution:

$$[\mathbf{x}]_{\mathcal{A}} = \begin{bmatrix} 3 \\ 4 \\ 1 \end{bmatrix} \Rightarrow [\mathbf{x}]_{\mathcal{B}} = P[\mathbf{x}]_{\mathcal{A}} = \begin{bmatrix} 4 & -1 & 0 \\ -1 & 1 & 1 \\ 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \\ 1 \end{bmatrix} = \begin{bmatrix} 8 \\ 2 \\ 2 \end{bmatrix}$$

Change of Basis



Cep telefonunuzu gözetmene teslim ediniz. Deposit your cell ph

5 December 2018 [16:00-17:10]

MATH215, Second Exam

3. Let $V = \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ -5 \end{bmatrix} \right\}$ and $W = \left\{ \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 6 \end{bmatrix} \right\}$ be two bases for \mathbb{R}^3 .

(a) 10 points Find the coordinates of $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ relative to the basis V .

Change of Basis



Cep telefonunuzu gözetmene teslim ediniz. Deposit your cell ph

5 December 2018 [16:00-17:10]

MATH215, Second Exam

3. Let $V = \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ -5 \end{bmatrix} \right\}$ and $W = \left\{ \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 6 \end{bmatrix} \right\}$ be two bases for \mathbb{R}^3 .

(a) 10 points Find the coordinates of $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ relative to the basis V .

$$\begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} = c_1 \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 2 \\ -3 \\ -3 \end{bmatrix} + c_3 \begin{bmatrix} 1 \\ -3 \\ -5 \end{bmatrix} \Rightarrow \begin{bmatrix} -1 & 2 & 1 \\ 1 & -3 & -3 \\ 0 & -3 & -5 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$$

$$\left[\begin{array}{ccc|c} -1 & 2 & 1 & 2 \\ 1 & -3 & -3 & -1 \\ 0 & -3 & -5 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & -2 & -1 & -2 \\ 0 & 1 & 2 & -1 \\ 0 & -3 & -5 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 0 & 3 & -4 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 1 & -2 \end{array} \right] \sim \left[\begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

$$[\mathbf{v}]_V = \begin{bmatrix} 2 \\ 3 \\ -2 \end{bmatrix}$$

Change of Basis



- (b) 10 points Find the change of coordinates matrix $P_{W \leftarrow V}$ from V to W .

Change of Basis



- (b) 10 points Find the change of coordinates matrix $P_{W \leftarrow V}$ from V to W .

Solution: We can use $P_{W \leftarrow V} = W^{-1}V$, $[W|V] \sim \left[I \mid P_{W \leftarrow V} \right]$ or $P_{W \leftarrow V} = [[\mathbf{v}_1]_W \quad [\mathbf{v}_2]_W \quad [\mathbf{v}_3]_W]$

$$[W|V] \sim \left[I \mid P_{W \leftarrow V} \right] \Rightarrow \left[\begin{array}{ccc|ccc} 1 & 1 & 2 & -1 & 2 & 1 \\ 2 & 2 & 3 & 1 & -3 & -3 \\ 4 & 3 & 6 & 0 & -3 & -5 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 1 & 2 & -1 & 2 & 1 \\ 0 & 0 & -1 & 3 & -7 & -5 \\ 0 & -1 & -2 & 4 & -11 & -9 \end{array} \right]$$
$$\sim \left[\begin{array}{ccc|ccc} 1 & 1 & 2 & -1 & 2 & 1 \\ 0 & 1 & 2 & -4 & 11 & 9 \\ 0 & 0 & 1 & -3 & 7 & 5 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 3 & -9 & -8 \\ 0 & 1 & 2 & -4 & 11 & 9 \\ 0 & 0 & 1 & -3 & 7 & 5 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 3 & -9 & -8 \\ 0 & 1 & 0 & 2 & -3 & -1 \\ 0 & 0 & 1 & -3 & 7 & 5 \end{array} \right]$$
$$P_{W \leftarrow V} = \begin{bmatrix} 3 & -9 & -8 \\ 2 & -3 & -1 \\ -3 & 7 & 5 \end{bmatrix}$$

Change of Basis



- (c) 5 points Find the coordinates of \mathbf{v} relative to W by using $P_{W \leftarrow V}$.

Solution:

$$[\mathbf{v}]_W = P_{W \leftarrow V} [\mathbf{v}]_V = \begin{bmatrix} 3 & -9 & -8 \\ 2 & -3 & -1 \\ -3 & 7 & 5 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \\ -2 \end{bmatrix} = \begin{bmatrix} -5 \\ -3 \\ 5 \end{bmatrix}$$

Change of Basis



Cep telefonunuzu gözetmene teslim ediniz. Deposit your cell phone

5 December 2018 [16:00-17:10]

MATH215, Second Exam

3. Let $V = \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ -3 \\ -3 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ -5 \end{bmatrix} \right\}$ and $W = \left\{ \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 6 \end{bmatrix} \right\}$ be two bases for \mathbb{R}^3 .

(a) 10 points Find the coordinates of $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ relative to the basis V .

Remark

I think that the 2019 question (below) is easier than the 2018 question (above), but I remember that the 2019 marks were lower than the 2018 marks.



30 December 2019 [11:00-12:15]

MATH215 Mathematics III Final Exam

Page 3 of 4

3. 25 points Let $\mathcal{A} = \{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ and $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be bases for the vector space V , and suppose $\mathbf{a}_1 = 4\mathbf{b}_1 - \mathbf{b}_2$, $\mathbf{a}_2 = -\mathbf{b}_1 + \mathbf{b}_2 + \mathbf{b}_3$, and $\mathbf{a}_3 = \mathbf{b}_2 - 2\mathbf{b}_3$.
- (a) Find the change of coordinates matrix from \mathcal{A} to \mathcal{B} .



Row Space, Column Space, and Null Space

Row Space, Column Space, and Null Space



Consider an $m \times n$ matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

Row Space, Column Space, and Null Space



Consider an $m \times n$ matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

Definition

The *row vectors* of A are the vectors

$$\mathbf{r}_1 = [a_{11} \quad a_{12} \quad \cdots \quad a_{1n}]$$

$$\mathbf{r}_2 = [a_{21} \quad a_{22} \quad \cdots \quad a_{2n}]$$

⋮

$$\mathbf{r}_m = [a_{m1} \quad a_{m2} \quad \cdots \quad a_{mn}]$$

Note that $\mathbf{r}_k \in \mathbb{R}^n$ for each $1 \leq k \leq m$.

Row Space, Column Space, and Null Space



Consider an $m \times n$ matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

Definition

The *column vectors of A* are the vectors

$$\mathbf{c}_1 = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix}, \quad \dots, \quad \mathbf{c}_m = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

Note that $\mathbf{c}_k \in \mathbb{R}^m$ for each $1 \leq k \leq n$.

Row Space, Column Space, and Null Space



Example

Find the row and column vectors of

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 3 & -1 & 4 \end{bmatrix}.$$

Row Space, Column Space, and Null Space



Example

Find the row and column vectors of

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 3 & -1 & 4 \end{bmatrix}.$$

The row vectors of A are

$$\mathbf{r}_1 = [2 \ 1 \ 0] \quad \text{and} \quad \mathbf{r}_2 = [3 \ -1 \ 4].$$

The column vectors of A are

$$\mathbf{c}_1 = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad \text{and} \quad \mathbf{c}_3 = \begin{bmatrix} 0 \\ 4 \end{bmatrix}.$$

Row Space, Column Space, and Null Space



Let A be an $m \times n$ matrix.

Definition

The subspace of \mathbb{R}^n spanned by the row vectors of A is called the *row space* of A . This is denoted by $\text{Row } A$.

Definition

Definition

Row Space, Column Space, and Null Space



Let A be an $m \times n$ matrix.

Definition

The subspace of \mathbb{R}^n spanned by the row vectors of A is called the *row space* of A . This is denoted by $\text{Row } A$.

Definition

The subspace of \mathbb{R}^m spanned by the column vectors of A is called the *column space* of A . This is denoted by $\text{Col } A$.

Definition

Row Space, Column Space, and Null Space



Let A be an $m \times n$ matrix.

Definition

The subspace of \mathbb{R}^n spanned by the row vectors of A is called the *row space* of A . This is denoted by $\text{Row } A$.

Definition

The subspace of \mathbb{R}^m spanned by the column vectors of A is called the *column space* of A . This is denoted by $\text{Col } A$.

Definition

The *null space* of A is the subspace

$$\text{Nul } A = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\} \subseteq \mathbb{R}^n.$$

Row Space, Column Space, and Null Space

Consider $A\mathbf{x} = \mathbf{b}$ where

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}.$$

Note that

$$\mathbf{b} = A\mathbf{x} = x_1\mathbf{c}_1 + x_2\mathbf{c}_2 + \dots + x_n\mathbf{c}_n$$

where $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$ are the column vectors of A .

Row Space, Column Space, and Null Space



Consider $A\mathbf{x} = \mathbf{b}$ where

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}.$$

Note that

$$\mathbf{b} = A\mathbf{x} = x_1\mathbf{c}_1 + x_2\mathbf{c}_2 + \dots + x_n\mathbf{c}_n$$

where $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$ are the column vectors of A . So $A\mathbf{x} = \mathbf{b}$ is consistent iff \mathbf{b} is a linear combination of the column vectors of A .

Row Space, Column Space, and Null Space



Consider $A\mathbf{x} = \mathbf{b}$ where

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}.$$

Note that

$$\mathbf{b} = A\mathbf{x} = x_1\mathbf{c}_1 + x_2\mathbf{c}_2 + \dots + x_n\mathbf{c}_n$$

where $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$ are the column vectors of A . So $A\mathbf{x} = \mathbf{b}$ is consistent iff \mathbf{b} is a linear combination of the column vectors of A .

Theorem

$$A\mathbf{x} = \mathbf{b} \text{ is consistent} \iff \mathbf{b} \in \text{Col } A$$

► EXAMPLE 2 A Vector \mathbf{b} in the Column Space of A

Let $A\mathbf{x} = \mathbf{b}$ be the linear system

$$\begin{bmatrix} -1 & 3 & 2 \\ 1 & 2 & -3 \\ 2 & 1 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -9 \\ -3 \end{bmatrix}$$

Show that \mathbf{b} is in the column space of A by expressing it as a linear combination of the column vectors of A .

Solution Solving the system by Gaussian elimination yields (verify)

$$x_1 = 2, \quad x_2 = -1, \quad x_3 = 3$$

It follows from this and Formula (2) that

$$2 \begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix} - \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} + 3 \begin{bmatrix} 2 \\ -3 \\ -2 \end{bmatrix} = \begin{bmatrix} 1 \\ -9 \\ -3 \end{bmatrix} \quad \blacktriangleleft$$

Row Space, Column Space, and Null Space



Example

Let

$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

- 1 Is \mathbf{u} in $\text{Nul } A$?
- 3 Is \mathbf{v} in $\text{Nul } A$?
- 2 Is \mathbf{u} in $\text{Col } A$?
- 4 Is \mathbf{v} in $\text{Col } A$?

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \text{ and } \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

1 Is \mathbf{u} in $\text{Nul } A$?

Recall that

$$\text{Nul } A = \{\mathbf{x} \in \mathbb{R}^4 \mid A\mathbf{x} = \mathbf{0}\} \subseteq \mathbb{R}^4$$

since A is a 3×4 matrix. So $\mathbf{u} \in \text{Nul } A$ if and only if $A\mathbf{u} = \mathbf{0}$.

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \text{ and } \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

1 Is \mathbf{u} in $\text{Nul } A$?

Recall that

$$\text{Nul } A = \{\mathbf{x} \in \mathbb{R}^4 \mid A\mathbf{x} = \mathbf{0}\} \subseteq \mathbb{R}^4$$

since A is a 3×4 matrix. So $\mathbf{u} \in \text{Nul } A$ if and only if $A\mathbf{u} = \mathbf{0}$.

We calculate that

$$A\mathbf{u} = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix} \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

Therefore the answer is YES.

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

2 Is \mathbf{u} in $\text{Col } A$?

Since A is a 3×4 matrix, $\text{Col } A$ is a subspace of \mathbb{R}^3 .

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \text{ and } \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

2 Is \mathbf{u} in $\text{Col } A$?

Since A is a 3×4 matrix, $\text{Col } A$ is a subspace of \mathbb{R}^3 . But $\mathbf{u} \in \mathbb{R}^4$. Therefore the answer is NO.

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

3 Is \mathbf{v} in $\text{Nul } A$?

NO, because $\mathbf{v} \in \mathbb{R}^3$, but $\text{Nul } A \subseteq \mathbb{R}^4$.

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

4 Is \mathbf{v} in $\text{Col } A$?

To answer this, we need to reduce $[A \ \mathbf{v}]$ to REF.

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

4 Is \mathbf{v} in $\text{Col } A$?

To answer this, we need to reduce $[A \ \mathbf{v}]$ to REF.

$$[A \ \mathbf{v}] = \begin{bmatrix} 2 & 4 & -2 & 1 & 3 \\ -2 & -5 & 7 & 3 & -1 \\ 3 & 7 & -8 & 6 & 3 \end{bmatrix}$$

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

4 Is \mathbf{v} in $\text{Col } A$?

To answer this, we need to reduce $[A \ \mathbf{v}]$ to REF.

$$[A \ \mathbf{v}] = \begin{bmatrix} 2 & 4 & -2 & 1 & 3 \\ -2 & -5 & 7 & 3 & -1 \\ 3 & 7 & -8 & 6 & 3 \end{bmatrix} \sim \begin{bmatrix} 2 & 4 & -2 & 1 & 3 \\ 0 & 1 & -5 & -4 & -2 \\ 0 & 0 & 0 & 17 & 1 \end{bmatrix}$$

Row Space, Column Space, and Null Space



$$A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 3 \end{bmatrix}.$$

4 Is \mathbf{v} in $\text{Col } A$?

To answer this, we need to reduce $[A \ \mathbf{v}]$ to REF.

$$[A \ \mathbf{v}] = \begin{bmatrix} 2 & 4 & -2 & 1 & 3 \\ -2 & -5 & 7 & 3 & -1 \\ 3 & 7 & -8 & 6 & 3 \end{bmatrix} \sim \begin{bmatrix} 2 & 4 & -2 & 1 & 3 \\ 0 & 1 & -5 & -4 & -2 \\ 0 & 0 & 0 & 17 & 1 \end{bmatrix}$$

We can now see that the linear system $A\mathbf{x} = \mathbf{v}$ is consistent. Therefore the answer to this question is YES.

Theorem

Suppose that

- $A\mathbf{x} = \mathbf{b}$ is consistent;
- \mathbf{x}_0 is any solution of $A\mathbf{x} = \mathbf{b}$;
- $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ is a basis for $\text{Nul } A$.

Theorem

Suppose that

- $A\mathbf{x} = \mathbf{b}$ is consistent;
- \mathbf{x}_0 is any solution of $A\mathbf{x} = \mathbf{b}$;
- $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ is a basis for $\text{Nul } A$.

Then every solution of $A\mathbf{x} = \mathbf{b}$ can be written in the form

$$\mathbf{x} = \mathbf{x}_0 + c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k.$$

Row Space, Column Space, and Null Space



Theorem

Suppose that

- $A\mathbf{x} = \mathbf{b}$ is consistent;
- \mathbf{x}_0 is any solution of $A\mathbf{x} = \mathbf{b}$;
- $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ is a basis for $\text{Nul } A$.

Then every solution of $A\mathbf{x} = \mathbf{b}$ can be written in the form

$$\mathbf{x} = \mathbf{x}_0 + c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k.$$

Conversely, the vector \mathbf{x} in this formula is a solution of $A\mathbf{x} = \mathbf{b}$ for any choice of scalars $c_j \in \mathbb{R}$.

Row Space, Column Space, and Null Space



$$\mathbf{x} = \mathbf{x}_0 + c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_k \mathbf{v}_k$$

Row Space, Column Space, and Null Space



$$\mathbf{x} = \underbrace{\mathbf{x}_0}_{\text{a particular solution of}} + c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_k \mathbf{v}_k$$

a particular
solution of

$$A\mathbf{x} = \mathbf{b}$$

Row Space, Column Space, and Null Space



$$\mathbf{x} = \underbrace{\mathbf{x}_0}_{\text{a particular solution of } A\mathbf{x} = \mathbf{b}} + \underbrace{c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k}_{\text{the general solution of } A\mathbf{x} = \mathbf{0}}$$

Row Space, Column Space, and Null Space



$$\mathbf{x} = \underbrace{\mathbf{x}_0}_{\text{a particular solution of } A\mathbf{x} = \mathbf{b}} + \underbrace{c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k}_{\text{the general solution of } A\mathbf{x} = 0}$$

the general solution of $A\mathbf{x} = \mathbf{b}$

Bases for Row Spaces, Column Spaces, and Null Spaces

Theorem

Elementary row operations do not change the null space of a matrix.

Theorem

Elementary row operations do not change the row space of a matrix.

Row Space, Column Space, and Null Space



Remark

Elementary row operations can change the column space of a matrix.

Row Space, Column Space, and Null Space



Remark

Elementary row operations can change the column space of a matrix.

For example, consider

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 3 \\ 0 & 0 \end{bmatrix}.$$

Note that A and B are row equivalent,

Row Space, Column Space, and Null Space



Remark

Elementary row operations can change the column space of a matrix.

For example, consider

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 3 \\ 0 & 0 \end{bmatrix}.$$

Note that A and B are row equivalent, but

$$\text{Col } A = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$$

Row Space, Column Space, and Null Space



Remark

Elementary row operations can change the column space of a matrix.

For example, consider

$$A = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 3 \\ 0 & 0 \end{bmatrix}.$$

Note that A and B are row equivalent, but

$$\text{Col } A = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$$

and

$$\text{Col } B = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\} \neq \text{Col } A.$$

Row Space, Column Space, and Null Space



Example

Find a basis the null space of the matrix

$$A = \begin{bmatrix} 1 & 3 & -2 & 0 & 2 & 0 \\ 2 & 6 & -5 & -2 & 4 & -3 \\ 0 & 0 & 5 & 10 & 0 & 15 \\ 2 & 6 & 0 & 8 & 4 & 18 \end{bmatrix}.$$

Row Space, Column Space, and Null Space



Example

Find a basis the null space of the matrix

$$A = \begin{bmatrix} 1 & 3 & -2 & 0 & 2 & 0 \\ 2 & 6 & -5 & -2 & 4 & -3 \\ 0 & 0 & 5 & 10 & 0 & 15 \\ 2 & 6 & 0 & 8 & 4 & 18 \end{bmatrix}.$$

We need to find the solution space of $A\mathbf{x} = \mathbf{0}$. I leave it to you to check that

$$\left[\begin{array}{cccccc|c} 1 & 3 & -2 & 0 & 2 & 0 & 0 \\ 2 & 6 & -5 & -2 & 4 & -3 & 0 \\ 0 & 0 & 5 & 10 & 0 & 15 & 0 \\ 2 & 6 & 0 & 8 & 4 & 18 & 0 \end{array} \right] \sim \left[\begin{array}{cccccc|c} 1 & 3 & 0 & 4 & 2 & 0 & 0 \\ 0 & 0 & 1 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right].$$

Row Space, Column Space, and Null Space



Therefore the solution of $A\mathbf{x} = \mathbf{0}$ is

$$\begin{cases} x_1 = -3x_2 - 4x_4 - 2x_5 \\ x_2 \text{ is free} \\ x_3 = -2x_4 \\ x_4 \text{ is free} \\ x_5 \text{ is free} \\ x_6 = 0 \end{cases}$$

Row Space, Column Space, and Null Space



Therefore the solution of $A\mathbf{x} = \mathbf{0}$ is

$$\begin{cases} x_1 = -3x_2 - 4x_4 - 2x_5 \\ x_2 \text{ is free} \\ x_3 = -2x_4 \\ x_4 \text{ is free} \\ x_5 \text{ is free} \\ x_6 = 0 \end{cases}$$

or in vectors,

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} -3x_2 - 4x_4 - 2x_5 \\ x_2 \\ -2x_4 \\ x_4 \\ x_5 \\ 0 \end{bmatrix} = x_2 \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -4 \\ 0 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -2 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}.$$

Row Space, Column Space, and Null Space



Therefore the solution of $A\mathbf{x} = \mathbf{0}$ is

$$\begin{cases} x_1 = -3x_2 - 4x_4 - 2x_5 \\ x_2 \text{ is free} \\ x_3 = -2x_4 \\ x_4 \text{ is free} \\ x_5 \text{ is free} \\ x_6 = 0 \end{cases}$$

or in vectors,

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} -3x_2 - 4x_4 - 2x_5 \\ x_2 \\ -2x_4 \\ x_4 \\ x_5 \\ 0 \end{bmatrix} = x_2 \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -4 \\ 0 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -2 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}.$$

Row Space, Column Space, and Null Space



Therefore

$$\left\{ \mathbf{v}_1 = \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -4 \\ 0 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -2 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

is a basis for $\text{Nul } A$.

Row Space, Column Space, and Null Space



Remark

In the previous example we found
that that

$$\begin{cases} x_1 = -3x_2 - 4x_4 - 2x_5 \\ x_2 \text{ is free} \\ x_3 = -2x_4 \\ x_4 \text{ is free} \\ x_5 \text{ is free} \\ x_6 = 0 \end{cases}$$

is the general solution of $A\mathbf{x} = \mathbf{0}$.

Row Space, Column Space, and Null Space

Remark

In the previous example we found that that

$$\begin{cases} x_1 = -3x_2 - 4x_4 - 2x_5 \\ x_2 \text{ is free} \\ x_3 = -2x_4 \\ x_4 \text{ is free} \\ x_5 \text{ is free} \\ x_6 = 0 \end{cases}$$

is the general solution of $A\mathbf{x} = \mathbf{0}$.

Note that if we set $x_2 = 1$ and $x_4 = x_5 = 0$ then we get

$$\mathbf{v}_1 = \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

which is the first vector in $\text{Nul } A$.

Row Space, Column Space, and Null Space



Remark

In the previous example we found that that

$$\begin{cases} x_1 = -3x_2 - 4x_4 - 2x_5 \\ x_2 \text{ is free} \\ x_3 = -2x_4 \\ x_4 \text{ is free} \\ x_5 \text{ is free} \\ x_6 = 0 \end{cases}$$

is the general solution of $A\mathbf{x} = \mathbf{0}$.

Note that if we set $x_2 = 1$ and $x_4 = x_5 = 0$ then we get

$$\mathbf{v}_1 = \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

which is the first vector in $\text{Nul } A$.

Setting $x_4 = 1$ and then $x_5 = 1$ (and the other two to 0) gives us the other two vectors in $\text{Nul } A$.

Theorem

Suppose that a matrix R is in row echelon form. Then

- the row vectors with the pivots (i.e. the nonzero row vectors) form a basis for the row space of R ; and
- the column vectors with the pivots (i.e. the pivot columns) form a basis for the column space of R .

Row Space, Column Space, and Null Space



Example

Find bases for the row and column spaces of the matrix

$$R = \begin{bmatrix} 1 & -2 & 5 & 0 & 3 \\ 0 & 1 & 3 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

This matrix is in REF, so we can use the previous theorem.

Row Space, Column Space, and Null Space



Example

Find bases for the row and column spaces of the matrix

$$R = \begin{bmatrix} 1 & -2 & 5 & 0 & 3 \\ 0 & 1 & 3 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

This matrix is in REF, so we can use the previous theorem.

There are three pivots (shown in green).

Row Space, Column Space, and Null Space



Example

Find bases for the row and column spaces of the matrix

$$R = \begin{bmatrix} 1 & -2 & 5 & 0 & 3 \\ 0 & 1 & 3 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

This matrix is in REF, so we can use the previous theorem.

There are three pivots (shown in green).

- $\{\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3\}$ is a basis for Row R since the pivots are in rows 1, 2 and 3.

Row Space, Column Space, and Null Space



Example

Find bases for the row and column spaces of the matrix

$$R = \begin{bmatrix} 1 & -2 & 5 & 0 & 3 \\ 0 & 1 & 3 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

This matrix is in REF, so we can use the previous theorem.

There are three pivots (shown in green).

- $\{\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3\}$ is a basis for Row R since the pivots are in rows 1, 2 and 3.
- $\{\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_4\}$ is a basis for Col R since the pivots are in columns 1, 2 and 4.

Example

Find a basis for the row space of the matrix

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix}.$$

Recall that elementary row operations do not change the row space of a matrix.

Example

Find a basis for the row space of the matrix

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix}.$$

Recall that elementary row operations do not change the row space of a matrix. So our method is:

- 1 Reduce A to REF. Call this new matrix R ;
- 2 Take the row vectors of R which contain a pivot.

Row Space, Column Space, and Null Space



As always, I leave it for you to check that

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix} \sim R = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 0 & 0 & 1 & 3 & -2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Therefore

$$\mathbf{r}_1 = [\ 1 \ -3 \ 4 \ -2 \ 5 \ 4 \]$$

$$\mathbf{r}_2 = [\ 0 \ 0 \ 1 \ 3 \ -2 \ -6 \]$$

$$\mathbf{r}_3 = [\ 0 \ 0 \ 0 \ 0 \ 1 \ 5 \]$$

form a basis for $\text{Row } A = \text{Row } R$.



Finding a Basis for Col A

Elementary row operations can change the column space of a matrix A ,

Finding a Basis for Col A

Elementary row operations can change the column space of a matrix A , but they do not change *which columns* of A we want in our basis.

Finding a Basis for $\text{Col } A$

Elementary row operations can change the column space of a matrix A , but they do not change *which columns* of A we want in our basis.

Theorem

The pivot columns of A form a basis of $\text{Col } A$.

Finding a Basis for $\text{Col } A$

Elementary row operations can change the column space of a matrix A , but they do not change *which columns* of A we want in our basis.

Theorem

The pivot columns of A form a basis of $\text{Col } A$.

So our method is:

- 1 Reduce A to REF. Call this new matrix R ;
- 2 Find the pivot columns of R ;
- 3 Go back to looking at the original matrix A ;
- 4 Take the pivot columns of A .

Row Space, Column Space, and Null Space



Example

Find a basis for the column space of the matrix

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix}.$$

Row Space, Column Space, and Null Space



Recall that

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix} \sim R = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 0 & 0 & 1 & 3 & -2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Look at R . The pivot columns of R are the first, third and fifth columns.

Row Space, Column Space, and Null Space



Recall that

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix} \sim R = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 0 & 0 & 1 & 3 & -2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Look at R . The pivot columns of R are the first, third and fifth columns. Therefore the pivot columns of A are also the **first, third and fifth** columns.

Row Space, Column Space, and Null Space



Recall that

$$A = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 2 & -6 & 9 & -1 & 8 & 2 \\ 2 & -6 & 9 & -1 & 9 & 7 \\ -1 & 3 & -4 & 2 & -5 & -4 \end{bmatrix} \sim R = \begin{bmatrix} 1 & -3 & 4 & -2 & 5 & 4 \\ 0 & 0 & 1 & 3 & -2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Look at R . The pivot columns of R are the first, third and fifth columns. Therefore the pivot columns of A are also the **first, third and fifth** columns. Hence

$$\left\{ \mathbf{c}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \\ -1 \end{bmatrix}, \mathbf{c}_3 = \begin{bmatrix} 4 \\ 9 \\ 9 \\ -4 \end{bmatrix}, \mathbf{c}_5 = \begin{bmatrix} 5 \\ 8 \\ 9 \\ -5 \end{bmatrix} \right\}$$

is a basis for $\text{Col } A$.

Row Space, Column Space, and Null Space



Example

Find a subset of the vectors

$$\mathbf{v}_1 = (1, -2, 0, 3), \quad \mathbf{v}_2 = (2, -5, -3, 6), \quad \mathbf{v}_3 = (0, 1, 3, 0),$$

$$\mathbf{v}_4 = (2, -1, 4, -7), \quad \mathbf{v}_5 = (5, -8, 1, 2)$$

that forms a basis for $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_5\}$.

Example

Find a subset of the vectors

$$\mathbf{v}_1 = (1, -2, 0, 3), \quad \mathbf{v}_2 = (2, -5, -3, 6), \quad \mathbf{v}_3 = (0, 1, 3, 0),$$

$$\mathbf{v}_4 = (2, -1, 4, -7), \quad \mathbf{v}_5 = (5, -8, 1, 2)$$

that forms a basis for $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_5\}$.

If we write these vectors as columns of a matrix

$$A = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \mathbf{v}_3 & \mathbf{v}_4 & \mathbf{v}_5 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 & 2 & 5 \\ -2 & -5 & 1 & -1 & -8 \\ 0 & -3 & 3 & 4 & 1 \\ 3 & 6 & 0 & -7 & 2 \end{bmatrix}$$

then we just need to find a basis for $\text{Col } A$.

Row Space, Column Space, and Null Space



Since

$$A \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & -1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

(please check) we can see that the pivot columns are the first, second and fourth columns.

Row Space, Column Space, and Null Space



Since

$$A \sim \begin{bmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & -1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

(please check) we can see that the pivot columns are the first, second and fourth columns.

Therefore $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_4\}$ is a basis for $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_5\}$.



Next Time

- Rank and Nullity
- The Fundamental Matrix Spaces
- Linear Transformations
- Composition and Inverse Transformations
- Isomorphisms