

# Lecture 5

- Vector Spaces
- Subspaces
- Null Spaces, Column Spaces, and Linear Transformations
- Linearly Independent Sets; Bases



# Vector Spaces

# Vector Spaces



You studied vectors, the dot product, etc. in MATH114. Now it is time to generalise these concepts.

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined.

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1** If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1** If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2**  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1** If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2**  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- 3**  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$   
such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$   
for all  $\mathbf{u} \in V$ .

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$   
such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$   
for all  $\mathbf{u} \in V$ .
- 5 For each  $\mathbf{u}$  in  $V$ , there is an  
object  $-\mathbf{u}$  in  $V$  such that  
 $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$   
such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$   
for all  $\mathbf{u} \in V$ .
- 5 For each  $\mathbf{u}$  in  $V$ , there is an  
object  $-\mathbf{u}$  in  $V$  such that  
 $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
- 6 If  $k$  is any scalar (number)  
and  $\mathbf{u}$  is any object in  $V$ ,  
then  $k\mathbf{u}$  is in  $V$ .

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$   
such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$   
for all  $\mathbf{u} \in V$ .
- 5 For each  $\mathbf{u}$  in  $V$ , there is an  
object  $-\mathbf{u}$  in  $V$  such that  
 $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
- 6 If  $k$  is any scalar (number)  
and  $\mathbf{u}$  is any object in  $V$ ,  
then  $k\mathbf{u}$  is in  $V$ .
- 7  $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$   
such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$   
for all  $\mathbf{u} \in V$ .
- 5 For each  $\mathbf{u}$  in  $V$ , there is an  
object  $-\mathbf{u}$  in  $V$  such that  
 $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
- 6 If  $k$  is any scalar (number)  
and  $\mathbf{u}$  is any object in  $V$ ,  
then  $k\mathbf{u}$  is in  $V$ .
- 7  $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$
- 8  $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$   
such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$   
for all  $\mathbf{u} \in V$ .
- 5 For each  $\mathbf{u}$  in  $V$ , there is an  
object  $-\mathbf{u}$  in  $V$  such that  
 $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
- 6 If  $k$  is any scalar (number)  
and  $\mathbf{u}$  is any object in  $V$ ,  
then  $k\mathbf{u}$  is in  $V$ .
- 7  $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$
- 8  $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$
- 9  $k(m\mathbf{u}) = (km)\mathbf{u}$

# Vector Spaces

## Definition

A *vector space* is a nonempty set  $V$  on which two operations are defined. These operations are called *addition* and *scalar multiplication* and must satisfy all of the following ten conditions:

- 1** If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ ,  
then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .
- 2**  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- 3**  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
- 4** There is a *zero vector*  $\mathbf{0}$  in  $V$   
such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$   
for all  $\mathbf{u} \in V$ .
- 5** For each  $\mathbf{u}$  in  $V$ , there is an  
object  $-\mathbf{u}$  in  $V$  such that  
 $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
- 6** If  $k$  is any scalar (number)  
and  $\mathbf{u}$  is any object in  $V$ ,  
then  $k\mathbf{u}$  is in  $V$ .
- 7**  $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$
- 8**  $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$
- 9**  $k(m\mathbf{u}) = (km)\mathbf{u}$
- 10**  $1\mathbf{u} = \mathbf{u}$ .

# Vector Spaces

## Remark

- If all the scalars  $k$  are real numbers, then  $V$  is called a *real vector space*.
- If we allow complex numbers  $k$ , then  $V$  is called a *complex vector space*.
- The objects  $\mathbf{v}$  in  $V$  are called *vectors*.
- Conditions 1-10 are called *axioms*.

# Vector Spaces

## Remark

- If all the scalars  $k$  are real numbers, then  $V$  is called a *real vector space*.
- If we allow complex numbers  $k$ , then  $V$  is called a *complex vector space*.
- The objects  $\mathbf{v}$  in  $V$  are called *vectors*.
- Conditions 1-10 are called *axioms*.
- The *zero vector*  $\mathbf{0}$  is unique.
- The negative of  $\mathbf{u}$ , denoted  $-\mathbf{u}$ , is unique for each  $\mathbf{u} \in V$ .

# Vector Spaces

## Remark

- If all the scalars  $k$  are real numbers, then  $V$  is called a *real vector space*.
- If we allow complex numbers  $k$ , then  $V$  is called a *complex vector space*.
- The objects  $\mathbf{v}$  in  $V$  are called *vectors*.
- Conditions 1-10 are called *axioms*.
- The *zero vector*  $\mathbf{0}$  is unique.
- The negative of  $\mathbf{u}$ , denoted  $-\mathbf{u}$ , is unique for each  $\mathbf{u} \in V$ .
- $0\mathbf{u} = \mathbf{0}$  for any  $\mathbf{u} \in V$

# Vector Spaces

## Remark

- If all the scalars  $k$  are real numbers, then  $V$  is called a *real vector space*.
- If we allow complex numbers  $k$ , then  $V$  is called a *complex vector space*.
- The objects  $\mathbf{v}$  in  $V$  are called *vectors*.
- Conditions 1-10 are called *axioms*.
- The *zero vector*  $\mathbf{0}$  is unique.
- The negative of  $\mathbf{u}$ , denoted  $-\mathbf{u}$ , is unique for each  $\mathbf{u} \in V$ .
- $0\mathbf{u} = \mathbf{0}$  for any  $\mathbf{u} \in V$
- $k\mathbf{0} = \mathbf{0}$  for any number  $k$
- $-\mathbf{u} = (-1)\mathbf{u}$  for any  $\mathbf{u} \in V$

# Vector Spaces

## Remark

- If all the scalars  $k$  are real numbers, then  $V$  is called a *real vector space*.
- If we allow complex numbers  $k$ , then  $V$  is called a *complex vector space*.
- The objects  $\mathbf{v}$  in  $V$  are called *vectors*.
- Conditions 1-10 are called *axioms*.
- The *zero vector*  $\mathbf{0}$  is unique.
- The negative of  $\mathbf{u}$ , denoted  $-\mathbf{u}$ , is unique for each  $\mathbf{u} \in V$ .
- $0\mathbf{u} = \mathbf{0}$  for any  $\mathbf{u} \in V$
- $k\mathbf{0} = \mathbf{0}$  for any number  $k$
- $-\mathbf{u} = (-1)\mathbf{u}$  for any  $\mathbf{u} \in V$
- $k\mathbf{u} = \mathbf{0} \implies k = 0 \text{ or } \mathbf{u} = \mathbf{0}$ .

## To Show That a Set with Two Operations Is a Vector Space

- Step 1.** Identify the set  $V$  of objects that will become vectors.
- Step 2.** Identify the addition and scalar multiplication operations on  $V$ .
- Step 3.** Verify Axioms 1 and 6; that is, adding two vectors in  $V$  produces a vector in  $V$ , and multiplying a vector in  $V$  by a scalar also produces a vector in  $V$ .  
Axiom 1 is called ***closure under addition***, and Axiom 6 is called ***closure under scalar multiplication***.
- Step 4.** Confirm that Axioms 2, 3, 4, 5, 7, 8, 9, and 10 hold.

# Vector Spaces



## Example (Euclidean Vector Spaces)

The spaces  $\mathbb{R}^n$  for  $n \geq 1$  are vector spaces.

# Vector Spaces

## Example (Euclidean Vector Spaces)

The spaces  $\mathbb{R}^n$  for  $n \geq 1$  are vector spaces.

## Example (The Zero Vector Space)

Let  $V$  be a set containing a single object called  $\mathbf{0}$ , and define addition and scalar multiplication by

$$\mathbf{0} + \mathbf{0} = \mathbf{0} \quad \text{and} \quad k\mathbf{0} = \mathbf{0}$$

for all scalars  $k$ .

# Vector Spaces

## Example (Euclidean Vector Spaces)

The spaces  $\mathbb{R}^n$  for  $n \geq 1$  are vector spaces.

## Example (The Zero Vector Space)

Let  $V$  be a set containing a single object called  $\mathbf{0}$ , and define addition and scalar multiplication by

$$\mathbf{0} + \mathbf{0} = \mathbf{0} \quad \text{and} \quad k\mathbf{0} = \mathbf{0}$$

for all scalars  $k$ .

I leave it to you to check that all 10 axioms are satisfied. We call this the *zero vector space*.

# Vector Spaces

## Example (The Vector Space of Sequences)

Let  $V$  consist of objects of the form

$$\mathbf{u} = (u_1, u_2, \dots, u_n, \dots)$$

in which  $u_1, u_2, \dots, u_n, \dots$  is an infinite sequence of real numbers.

# Vector Spaces

## Example (The Vector Space of Sequences)

Let  $V$  consist of objects of the form

$$\mathbf{u} = (u_1, u_2, \dots, u_n, \dots)$$

in which  $u_1, u_2, \dots, u_n, \dots$  is an infinite sequence of real numbers.

We define two infinite sequences to be equal if their corresponding components are equal, and we define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (u_1, u_2, \dots, u_n, \dots) + (v_1, v_2, \dots, v_n, \dots) \\ &= (u_1 + v_1, u_2 + v_2, \dots, u_n + v_n, \dots) \\ k\mathbf{u} &= (ku_1, ku_2, \dots, ku_n, \dots)\end{aligned}$$

Show that  $V$  is a vector space.

# Vector Spaces

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ , then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .

As we know from MATH114, the sum of two sequences is a sequence. ✓

This follows from the  $a + b = b + a$  rule for real numbers.



# Vector Spaces

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ , then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .

As we know from MATH114, the sum of two sequences is a sequence. ✓

- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

This follows from the  $a + b = b + a$  rule for real numbers.



# Vector Spaces

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ , then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .

As we know from MATH114, the sum of two sequences is a sequence. ✓

- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

This follows from the  $a + b = b + a$  rule for real numbers.



- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$

Similar to 2. ✓

# Vector Spaces

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ , then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .

As we know from MATH114, the sum of two sequences is a sequence. ✓

- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

This follows from the  $a + b = b + a$  rule for real numbers.



- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$

Similar to 2. ✓

- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$  such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$  for all  $\mathbf{u} \in V$ .

Just take  $\mathbf{0} = (0, 0, 0, 0, 0, \dots)$ . ✓

# Vector Spaces

- 1 If  $\mathbf{u}$  and  $\mathbf{v}$  are objects in  $V$ , then  $\mathbf{u} + \mathbf{v}$  is in  $V$ .

As we know from MATH114, the sum of two sequences is a sequence. ✓

- 2  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

This follows from the  $a + b = b + a$  rule for real numbers.



- 3  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$

Similar to 2. ✓

- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$  such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$  for all  $\mathbf{u} \in V$ .

Just take  $\mathbf{0} = (0, 0, 0, 0, 0, \dots)$ . ✓

- 5 For each  $\mathbf{u}$  in  $V$ , there is an object  $-\mathbf{u}$  in  $V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ . ✓

We have  $-\mathbf{u} = (-u_1, -u_2, -u_3, -u_4, \dots)$ . ✓

# Vector Spaces



- 6 If  $k$  is any scalar (number) and  $\mathbf{u}$  is any object in  $V$ , then  $k\mathbf{u}$  is in  $V$ .
- A sequence multiplied by a number is a sequence. ✓

# Vector Spaces

- 6 If  $k$  is any scalar (number) and  $\mathbf{u}$  is any object in  $V$ , then  $k\mathbf{u}$  is in  $V$ .  
A sequence multiplied by a number is a sequence. ✓
- 7  $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$   
Follows from the same property for numbers. ✓

# Vector Spaces

- 6 If  $k$  is any scalar (number) and  $\mathbf{u}$  is any object in  $V$ , then  $k\mathbf{u}$  is in  $V$ .

A sequence multiplied by a number is a sequence. ✓

- 7  $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$

Follows from the same property for numbers. ✓

- 8  $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$

See 7. ✓

# Vector Spaces

- 6 If  $k$  is any scalar (number) and  $\mathbf{u}$  is any object in  $V$ , then  $k\mathbf{u}$  is in  $V$ .

A sequence multiplied by a number is a sequence. ✓

- 7  $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$

Follows from the same property for numbers. ✓

- 8  $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$

See 7. ✓

- 9  $k(m\mathbf{u}) = (km)\mathbf{u}$

See 7 again. ✓

# Vector Spaces

- 6 If  $k$  is any scalar (number) and  $\mathbf{u}$  is any object in  $V$ , then  $k\mathbf{u}$  is in  $V$ .

A sequence multiplied by a number is a sequence. ✓

- 7  $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$

Follows from the same property for numbers. ✓

- 8  $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$

See 7. ✓

- 9  $k(m\mathbf{u}) = (km)\mathbf{u}$

See 7 again. ✓

- 10  $1\mathbf{u} = \mathbf{u}$ .

Clearly  $1\mathbf{u} = (1u_1, 1u_2, 1u_3, \dots) = (u_1, u_2, u_3, \dots) = \mathbf{u}$  ✓

Therefore  $V$  is a vector space.

# Vector Spaces

## Example

Let

$$\begin{aligned}\mathbb{P}^n &= \{\text{all polynomials of degree } \leq n\} \\ &= \{\mathbf{p}(t) = a_0 + a_1t + a_2t^2 + \dots + a_nt^n \mid a_j \in \mathbb{R}\}\end{aligned}$$

for  $n \in \mathbb{N} \cup \{0\}$ . (See page 210, example 4 in your textbook.) Let addition and scalar multiplication be defined in the obvious way.

Show that  $\mathbb{P}^n$  is a vector space.

# Vector Spaces

First let's do axioms 1 and 6. If

$$\mathbf{p}(t) = a_0 + a_1 t + a_2 t^2 + \dots + a_n t^n$$

and

$$\mathbf{q}(t) = b_0 + b_1 t + b_2 t^2 + \dots + b_n t^n,$$

then

$$\begin{aligned}(\mathbf{p} + \mathbf{q})(t) &= \mathbf{p}(t) + \mathbf{q}(t) \\&= (a_0 + b_0) + (a_1 + b_1)t + (a_2 + b_2)t^2 + \dots + (a_n + b_n)t^n\end{aligned}$$

is a polynomial of degree  $\leq n$ , and

$$k\mathbf{p}(t) = k a_0 + k a_1 t + k a_2 t^2 + \dots + k a_n t^n$$

is also a polynomial of degree  $\leq n$ . So 1. and 6. are satisfied.

# Vector Spaces



Axioms 2,3 and 7-10 follows from properties of the real numbers. So we just need to look at 4. and 5.

Axioms 2,3 and 7-10 follows from properties of the real numbers. So we just need to look at 4. and 5.

- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$  such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$  for all  $\mathbf{u} \in V$ .
  
- 5 For each  $\mathbf{u}$  in  $V$ , there is an object  $-\mathbf{u}$  in  $V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .

Axioms 2,3 and 7-10 follows from properties of the real numbers. So we just need to look at 4. and 5.

- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$  such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$  for all  $\mathbf{u} \in V$ .

We can just let  $\mathbf{0}(t) = 0$  be the zero function.

- 5 For each  $\mathbf{u}$  in  $V$ , there is an object  $-\mathbf{u}$  in  $V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .

# Vector Spaces

Axioms 2,3 and 7-10 follows from properties of the real numbers. So we just need to look at 4. and 5.

- 4 There is a *zero vector*  $\mathbf{0}$  in  $V$  such that  $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$  for all  $\mathbf{u} \in V$ .

We can just let  $\mathbf{0}(t) = 0$  be the zero function.

- 5 For each  $\mathbf{u}$  in  $V$ , there is an object  $-\mathbf{u}$  in  $V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .

For  $\mathbf{p}(t)$  as before, just let

$$-\mathbf{p}(t) = -a_0 - a_1 t - a_2 t^2 - \dots - a_n t^n.$$

Therefore  $\mathbb{P}^n$  is a vector space.

# Vector Spaces

## Example

Is

$$V = \{\text{all polynomials of degree 2}\}$$

(with the obvious addition and scalar multiplication) a vector space?

# Vector Spaces

## Example

Is

$$V = \{\text{all polynomials of degree 2}\}$$

(with the obvious addition and scalar multiplication) a vector space?

No.

Consider  $\mathbf{p}(t) = 1 + x + x^2$  and  $\mathbf{q}(t) = 2 - x^2$ . Both  $\mathbf{p}$  and  $\mathbf{q}$  are polynomials of degree 2, so are in  $V$ .

# Vector Spaces

## Example

Is

$$V = \{\text{all polynomials of degree 2}\}$$

(with the obvious addition and scalar multiplication) a vector space?

No.

Consider  $\mathbf{p}(t) = 1 + x + x^2$  and  $\mathbf{q}(t) = 2 - x^2$ . Both  $\mathbf{p}$  and  $\mathbf{q}$  are polynomials of degree 2, so are in  $V$ . However

$$(\mathbf{p} + \mathbf{q})(t) = (1 + x + x^2) + (2 - x^2) = 3 + x$$

is a polynomial of degree 1, so is *not* in  $V$ .

# Vector Spaces

## Example

Is

$$V = \{\text{all polynomials of degree 2}\}$$

(with the obvious addition and scalar multiplication) a vector space?

No.

Consider  $\mathbf{p}(t) = 1 + x + x^2$  and  $\mathbf{q}(t) = 2 - x^2$ . Both  $\mathbf{p}$  and  $\mathbf{q}$  are polynomials of degree 2, so are in  $V$ . However

$$(\mathbf{p} + \mathbf{q})(t) = (1 + x + x^2) + (2 - x^2) = 3 + x$$

is a polynomial of degree 1, so is *not* in  $V$ .

$V$  is not closed under addition, so is not a vector space.

**EXAMPLE 5** Let  $V$  be the set of all real-valued functions defined on a set  $\mathbb{D}$ . (Typically,  $\mathbb{D}$  is the set of real numbers or some interval on the real line.) Functions are added in the usual way:  $\mathbf{f} + \mathbf{g}$  is the function whose value at  $t$  in the domain  $\mathbb{D}$  is  $\mathbf{f}(t) + \mathbf{g}(t)$ . Likewise, for a scalar  $c$  and an  $\mathbf{f}$  in  $V$ , the scalar multiple  $c\mathbf{f}$  is the function whose value at  $t$  is  $c\mathbf{f}(t)$ . For instance, if  $\mathbb{D} = \mathbb{R}$ ,  $\mathbf{f}(t) = 1 + \sin 2t$ , and  $\mathbf{g}(t) = 2 + .5t$ , then

$$(\mathbf{f} + \mathbf{g})(t) = 3 + \sin 2t + .5t \quad \text{and} \quad (2\mathbf{g})(t) = 4 + t$$

Two functions in  $V$  are equal if and only if their values are equal for every  $t$  in  $\mathbb{D}$ . Hence the zero vector in  $V$  is the function that is identically zero,  $\mathbf{f}(t) = 0$  for all  $t$ , and the negative of  $\mathbf{f}$  is  $(-1)\mathbf{f}$ . Axioms 1 and 6 are obviously true, and the other axioms follow from properties of the real numbers, so  $V$  is a vector space. ■

# Vector Spaces

## Example

Let

$$\mathbb{R}^{2 \times 2} = \{\text{all } 2 \times 2 \text{ matrices with real number entries}\}$$

and let addition and scalar multiplication be the usual operations on matrices: i.e.

$$\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} + \begin{bmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{bmatrix} = \begin{bmatrix} u_{11} + v_{11} & u_{12} + v_{12} \\ u_{21} + v_{21} & u_{22} + v_{22} \end{bmatrix}$$

$$k\mathbf{u} = k \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \begin{bmatrix} ku_{11} & ku_{12} \\ ku_{21} & ku_{22} \end{bmatrix}.$$

Show that  $\mathbb{R}^{2 \times 2}$  is a vector space.

(Some books use  $M_{22}$  instead of  $\mathbb{R}^{2 \times 2}$ .)

# Vector Spaces

- 1 matrix + matrix = a matrix. ✓
- 6 number × matrix = a matrix. ✓

# Vector Spaces

- 1 matrix + matrix = a matrix. ✓
- 6 number × matrix = a matrix. ✓

2

$$\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} + \begin{bmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{bmatrix} = \begin{bmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{bmatrix} + \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \mathbf{v} + \mathbf{u}. \checkmark$$

# Vector Spaces

- 1 matrix + matrix = a matrix. ✓
- 6 number × matrix = a matrix. ✓
- 2

$$\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} + \begin{bmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{bmatrix} = \begin{bmatrix} v_{11} & v_{12} \\ v_{21} & v_{22} \end{bmatrix} + \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \mathbf{v} + \mathbf{u}. \checkmark$$

I leave axioms 3,7,8,9 for you to prove.

- 4 Let  $\mathbf{0} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ . Then

$$\mathbf{0} + \mathbf{u} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \mathbf{u}$$

and similarly  $\mathbf{u} + \mathbf{0} = \mathbf{u}$ . ✓

# Vector Spaces

5 Let  $-\mathbf{u} = \begin{bmatrix} -u_{11} & -u_{12} \\ -u_{21} & -u_{22} \end{bmatrix}$ . Then

$$\mathbf{u} + (-\mathbf{u}) = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} + \begin{bmatrix} -u_{11} & -u_{12} \\ -u_{21} & -u_{22} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} . \checkmark$$

# Vector Spaces

5 Let  $-\mathbf{u} = \begin{bmatrix} -u_{11} & -u_{12} \\ -u_{21} & -u_{22} \end{bmatrix}$ . Then

$$\mathbf{u} + (-\mathbf{u}) = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} + \begin{bmatrix} -u_{11} & -u_{12} \\ -u_{21} & -u_{22} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \checkmark$$

10 Finally

$$1\mathbf{u} = 1 \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \mathbf{u}. \checkmark$$

# Vector Spaces

5 Let  $-\mathbf{u} = \begin{bmatrix} -u_{11} & -u_{12} \\ -u_{21} & -u_{22} \end{bmatrix}$ . Then

$$\mathbf{u} + (-\mathbf{u}) = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} + \begin{bmatrix} -u_{11} & -u_{12} \\ -u_{21} & -u_{22} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \checkmark$$

10 Finally

$$1\mathbf{u} = 1 \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix} = \mathbf{u}. \checkmark$$

This proves that  $\mathbb{R}^{2 \times 2} = M_{22}$  is a vector space.

# Vector Spaces



So far I have been using  $\mathbf{u} + \mathbf{v}$  and  $k\mathbf{u}$  for addition and scalar multiplication, but this might be confusing for more complicated operations.

So far I have been using  $\mathbf{u} + \mathbf{v}$  and  $k\mathbf{u}$  for addition and scalar multiplication, but this might be confusing for more complicated operations.

## Notation

Let

- $\mathbf{u} \oplus \mathbf{v}$  denote vector addition; and
- $k \odot \mathbf{u}$  denote scalar multiplication.

# Vector Spaces

## Example

Let  $V = \mathbb{R}^2$  and define  $\oplus$  and  $\odot$  as follows: If  $\mathbf{u} = (u_1, u_2)$  and  $\mathbf{v} = (v_1, v_2)$ , then we define

$$\mathbf{u} \oplus \mathbf{v} = (u_1 + v_1, u_2 + v_2)$$

and

$$k \odot \mathbf{u} = (ku_1, 0).$$

# Vector Spaces

## Example

Let  $V = \mathbb{R}^2$  and define  $\oplus$  and  $\odot$  as follows: If  $\mathbf{u} = (u_1, u_2)$  and  $\mathbf{v} = (v_1, v_2)$ , then we define

$$\mathbf{u} \oplus \mathbf{v} = (u_1 + v_1, u_2 + v_2) \xleftarrow{\text{usual addition}}$$

and

$$k \odot \mathbf{u} = (ku_1, 0). \xleftarrow{\text{weird scalar multiplication}}$$

# Vector Spaces

## Example

Let  $V = \mathbb{R}^2$  and define  $\oplus$  and  $\odot$  as follows: If  $\mathbf{u} = (u_1, u_2)$  and  $\mathbf{v} = (v_1, v_2)$ , then we define

$$\mathbf{u} \oplus \mathbf{v} = (u_1 + v_1, u_2 + v_2)$$

and

$$k \odot \mathbf{u} = (ku_1, 0).$$

For example, if  $\mathbf{u} = (2, 4)$ ,  $\mathbf{v} = (-3, 5)$  and  $k = 7$ , then

$$\mathbf{u} \oplus \mathbf{v} = (2 + (-3), 4 + 5) = (-1, 9)$$

$$k \odot \mathbf{u} = (7 \cdot 2, 0) = (14, 0).$$

Is  $V$  a vector space?

# Vector Spaces



$$k \odot \mathbf{u} = (ku_1, 0)$$

Axioms 1-9 are all true. I leave this for you to check.

# Vector Spaces



$$k \odot \mathbf{u} = (ku_1, 0)$$

Axioms 1-9 are all true. I leave this for you to check.

**10** However axiom 10 fails. For example, if  $\mathbf{u} = (0, 1)$  then

$$1 \odot \mathbf{u} = 1 \odot (0, 1) = (1 \cdot 0, 0) = (0, 0) \neq \mathbf{u}.$$

Therefore  $V$  is not a vector space with these operations.

# Vector Spaces



## Example (An Unusual Vector Space)

Let  $V$  be the set of strictly positive real numbers. Let  $\mathbf{u} = u$  and  $\mathbf{v} = v$  be any vectors in  $V$  (i.e. any real numbers  $> 0$ ) and let  $k$  be a scalar. Define  $\oplus$  and  $\odot$  by

$$\mathbf{u} \oplus \mathbf{v} = uv$$

$$k \odot \mathbf{v} = u^k.$$

# Vector Spaces



## Example (An Unusual Vector Space)

Let  $V$  be the set of strictly positive real numbers. Let  $\mathbf{u} = u$  and  $\mathbf{v} = v$  be any vectors in  $V$  (i.e. any real numbers  $> 0$ ) and let  $k$  be a scalar. Define  $\oplus$  and  $\odot$  by

$$\mathbf{u} \oplus \mathbf{v} = uv \quad \text{← weird addition}$$

$$k \odot \mathbf{v} = u^k. \quad \text{← weird scalar multiplication}$$

# Vector Spaces

## Example (An Unusual Vector Space)

Let  $V$  be the set of strictly positive real numbers. Let  $\mathbf{u} = u$  and  $\mathbf{v} = v$  be any vectors in  $V$  (i.e. any real numbers  $> 0$ ) and let  $k$  be a scalar. Define  $\oplus$  and  $\odot$  by

$$\mathbf{u} \oplus \mathbf{v} = uv$$

$$k \odot \mathbf{v} = u^k.$$

For example  $1 \oplus 2 = 2$  and  $2 \odot 1 = 1^2 = 1$ .

# Vector Spaces



## Example (An Unusual Vector Space)

Let  $V$  be the set of strictly positive real numbers. Let  $\mathbf{u} = u$  and  $\mathbf{v} = v$  be any vectors in  $V$  (i.e. any real numbers  $> 0$ ) and let  $k$  be a scalar. Define  $\oplus$  and  $\odot$  by

$$\mathbf{u} \oplus \mathbf{v} = uv$$

$$k \odot \mathbf{v} = u^k.$$

For example  $1 \oplus 2 = 2$  and  $2 \odot 1 = 1^2 = 1$ .

Show that  $V$  is a vector space with these operations.

# Vector Spaces



I will prove axioms 4, 5 and 7. I leave the others for you to prove.

# Vector Spaces

I will prove axioms 4, 5 and 7. I leave the others for you to prove.

- 4 The zero vector in this space is the number 1 (i.e.  $\mathbf{0} = 1$ ) since

$$\mathbf{u} \oplus \mathbf{0} = u \cdot 1 = u.$$

# Vector Spaces

I will prove axioms 4, 5 and 7. I leave the others for you to prove.

- 4 The zero vector in this space is the number 1 (i.e.  $\mathbf{0} = 1$ ) since

$$\mathbf{u} \oplus \mathbf{0} = u \cdot 1 = u.$$

- 5 The negative of a vector  $\mathbf{u} = u$  is  $-\mathbf{u} = \frac{1}{u}$  since

$$\mathbf{u} \oplus (-\mathbf{u}) = u \cdot \frac{1}{u} = 1 (= \mathbf{0}).$$

# Vector Spaces

I will prove axioms 4, 5 and 7. I leave the others for you to prove.

- 4 The zero vector in this space is the number 1 (i.e.  $\mathbf{0} = 1$ ) since

$$\mathbf{u} \oplus \mathbf{0} = u \cdot 1 = u.$$

- 5 The negative of a vector  $\mathbf{u} = u$  is  $-\mathbf{u} = \frac{1}{u}$  since

$$\mathbf{u} \oplus (-\mathbf{u}) = u \cdot \frac{1}{u} = 1 (= \mathbf{0}).$$

- 7 We have

$$k \odot (\mathbf{u} \oplus \mathbf{v}) = (uv)^k = u^k v^k = (k \odot \mathbf{u}) \oplus (k \odot \mathbf{v}).$$

# Vector Spaces

I will prove axioms 4, 5 and 7. I leave the others for you to prove.

- 4** The zero vector in this space is the number 1 (i.e.  $\mathbf{0} = 1$ ) since

$$\mathbf{u} \oplus \mathbf{0} = u \cdot 1 = u.$$

- 5** The negative of a vector  $\mathbf{u} = u$  is  $-\mathbf{u} = \frac{1}{u}$  since

$$\mathbf{u} \oplus (-\mathbf{u}) = u \cdot \frac{1}{u} = 1 (= \mathbf{0}).$$

- 7** We have

$$k \odot (\mathbf{u} \oplus \mathbf{v}) = (uv)^k = u^k v^k = (k \odot \mathbf{u}) \oplus (k \odot \mathbf{v}).$$

Therefore  $V$  is a vector space with these operations.


# Subspaces



# Null Spaces, Column Spaces, and Linear Trans- formations

# Break

We will continue at 3pm





# Linearly Independent Sets; Bases



# Next Time

