

Lecture 9

- 24. Limits
- 25. Continuity
- 26. Differentiation




Limits

24. Limits

Consider the function $f(x) = \frac{x^2 - 1}{x - 1}$, $f : (-\infty, 1) \cup (1, \infty) \rightarrow \mathbb{R}$.


24. Limits

Consider the function $f(x) = \frac{x^2-1}{x-1}$, $f : (-\infty, 1) \cup (1, \infty) \rightarrow \mathbb{R}$.



24. Limits


Consider the function $f(x) = \frac{x^2-1}{x-1}$, $f : (-\infty, 1) \cup (1, \infty) \rightarrow \mathbb{R}$.



Question: How does f behave when x is close to 1?

24. Limits

Consider the function $f(x) = \frac{x^2-1}{x-1}$, $f : (-\infty, 1) \cup (1, \infty) \rightarrow \mathbb{R}$.




x	$f(x)$
0.9	1.9
1.1	2.1

Question: How does f behave when x is close to 1?

24. Limits

Consider the function $f(x) = \frac{x^2-1}{x-1}$, $f : (-\infty, 1) \cup (1, \infty) \rightarrow \mathbb{R}$.




x	$f(x)$
0.9	1.9
1.1	2.1
0.99	1.99
1.01	2.01

Question: How does f behave when x is close to 1?

24. Limits

Consider the function $f(x) = \frac{x^2-1}{x-1}$, $f : (-\infty, 1) \cup (1, \infty) \rightarrow \mathbb{R}$.




x	$f(x)$
0.9	1.9
1.1	2.1
0.99	1.99
1.01	2.01
0.999	1.999
1.001	2.001

Question: How does f behave when x is close to 1?

24. Limits

Consider the function $f(x) = \frac{x^2-1}{x-1}$, $f : (-\infty, 1) \cup (1, \infty) \rightarrow \mathbb{R}$.



x	$f(x)$
0.9	1.9
1.1	2.1
0.99	1.99
1.01	2.01
0.999	1.999
1.001	2.001

“If x is close to 1, then $f(x)$ is close to 2.”

24. Limits



“If x is close to 1, then $f(x)$ is close to 2.”

Mathematically, we write this as


$$\lim_{x \rightarrow 1} f(x) = 2$$

and read it as “the limit, as x tends to 1, of $f(x)$ is equal to 2”.

24. Limits

Example

$$f(x) = \frac{x^2 - 1}{x - 1}$$




Note that $\lim_{x \rightarrow 1} f(x) = 2$,
but f is not defined at $x = 1$.

24. Limits

Example


$$g(x) = \begin{cases} \frac{x^2-1}{x-1} & x \neq 1 \\ 1 & x = 1 \end{cases}$$




24. Limits

Example

$$h(x) = x + 1$$




24. Limits



Example (The Identity Function)

$$f(x) = x$$

24. Limits




Example (The Identity Function)

$$f(x) = x$$

$$\lim_{x \rightarrow x_0} f(x) = \lim_{x \rightarrow x_0} x = x_0$$


24. Limits



Example (A Constant Function)

$$f(x) = 13$$

24. Limits



Example (A Constant Function)

$$f(x) = 13$$

$$\lim_{x \rightarrow x_0} f(x) = \lim_{x \rightarrow x_0} 13 = 13$$

24. Limits




Example (Sometimes Limits Do Not Exist)

Consider the functions


$$u(x) = \begin{cases} 0 & x < 0 \\ 1 & x \geq 0 \end{cases} \quad \text{and} \quad v(x) = \begin{cases} 0 & x \leq 0 \\ \sin \frac{1}{x} & x > 0. \end{cases}$$

24. Limits



Note that $\lim_{x \rightarrow 0} u(x)$ does not exist.


24. Limits



Note that $\lim_{x \rightarrow 0} u(x)$ does not exist. To understand why, we consider x close to 0:

- If x is close to 0 and $x < 0$, then $u(x) = 0$.
- If x is close to 0 and $x > 0$, then $u(x) = 1$.

24. Limits




Note that $\lim_{x \rightarrow 0} u(x)$ does not exist. To understand why, we consider x close to 0:

- If x is close to 0 and $x < 0$, then $u(x) = 0$.
- If x is close to 0 and $x > 0$, then $u(x) = 1$.

Because 0 is not close to 1, the limit as $x \rightarrow 0$ can not exist.

24. Limits



Moreover $\lim_{x \rightarrow 0} v(x)$ does not exist because $v(x)$ oscillates up and down too quickly if $x > 0$ and $x \rightarrow 0$.

24. Limits



Theorem (The Limit Laws)

Suppose that

- $L, M, c, k \in \mathbb{R}$;
- f and g are functions;
- $\lim_{x \rightarrow c} f(x) = L$; and
- $\lim_{x \rightarrow c} g(x) = M$.

Then

24. Limits



Theorem (The Limit Laws)

Suppose that

- $L, M, c, k \in \mathbb{R}$;
- f and g are functions;
- $\lim_{x \rightarrow c} f(x) = L$; and
- $\lim_{x \rightarrow c} g(x) = M$.

Then

- 1 Sum Rule:

$$\lim_{x \rightarrow c} (f(x) + g(x)) = L + M;$$

24. Limits

Theorem (The Limit Laws)

Suppose that

- $L, M, c, k \in \mathbb{R}$;
- f and g are functions;
- $\lim_{x \rightarrow c} f(x) = L$; and
- $\lim_{x \rightarrow c} g(x) = M$.

Then

2 Difference Rule:

$$\lim_{x \rightarrow c} (f(x) - g(x)) = L - M;$$

24. Limits

Theorem (The Limit Laws)

Suppose that

- $L, M, c, k \in \mathbb{R}$;
- f and g are functions;
- $\lim_{x \rightarrow c} f(x) = L$; and
- $\lim_{x \rightarrow c} g(x) = M$.

Then

3 Constant Multiple Rule:

$$\lim_{x \rightarrow c} (kf(x)) = kL;$$

24. Limits



Theorem (The Limit Laws)

Suppose that

- $L, M, c, k \in \mathbb{R}$;
- f and g are functions;
- $\lim_{x \rightarrow c} f(x) = L$; and
- $\lim_{x \rightarrow c} g(x) = M$.

Then

- 4 Product Rule:

$$\lim_{x \rightarrow c} (f(x)g(x)) = LM;$$

24. Limits

Theorem (The Limit Laws)

Suppose that

- $L, M, c, k \in \mathbb{R}$;
- f and g are functions;
- $\lim_{x \rightarrow c} f(x) = L$; and
- $\lim_{x \rightarrow c} g(x) = M$.

Then

- 5 Quotient Rule: if $M \neq 0$, then

$$\lim_{x \rightarrow c} \left(\frac{f(x)}{g(x)} \right) = \frac{L}{M};$$

24. Limits

Theorem (The Limit Laws)

Suppose that

- $L, M, c, k \in \mathbb{R}$;
- f and g are functions;
- $\lim_{x \rightarrow c} f(x) = L$; and
- $\lim_{x \rightarrow c} g(x) = M$.

Then

- 6 Power Rule: if $n \in \mathbb{N}$, then

$$\lim_{x \rightarrow c} (f(x))^n = L^n;$$

24. Limits

Theorem (The Limit Laws)

Suppose that

- $L, M, c, k \in \mathbb{R}$;
- f and g are functions;
- $\lim_{x \rightarrow c} f(x) = L$; and
- $\lim_{x \rightarrow c} g(x) = M$.

Then

- 7 Root Rule: if $n \in \mathbb{N}$ and $\sqrt[n]{L}$ exists, then

$$\lim_{x \rightarrow c} \sqrt[n]{f(x)} = \sqrt[n]{L} = L^{\frac{1}{n}}.$$

24. Limits



Example

$$\text{Find } \lim_{x \rightarrow 2} (x^3 + 4x^2 - 3).$$

24. Limits

Example

Find $\lim_{x \rightarrow 5} \frac{x^4 + x^2 - 1}{x^2 + 5}$.

solution:

$$\lim_{x \rightarrow 5} \frac{x^4 + x^2 - 1}{x^2 + 5} = \frac{\lim_{x \rightarrow 5}(x^4 + x^2 - 1)}{\lim_{x \rightarrow 5}(x^2 + 5)}$$

(quotient rule)

$$= \frac{\lim_{x \rightarrow 5} x^4 + \lim_{x \rightarrow 5} x^2 - \lim_{x \rightarrow 5} 1}{\lim_{x \rightarrow 5} x^2 + \lim_{x \rightarrow 5} 5}$$

(sum and difference rules)

$$= \frac{5^4 + 5^2 - 1}{5^2 + 5}$$

(power rule)

$$= \frac{649}{30}.$$

24. Limits



Theorem (Limits of Polynomial Functions)

If $P(x) = a_nx^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$ is a polynomial function, then

$$\lim_{x \rightarrow c} P(x) = P(c).$$

24. Limits



Theorem (Limits of Rational Functions)

If $P(x)$ and $Q(x)$ are polynomial functions and if $Q(c) \neq 0$, then

$$\lim_{x \rightarrow c} \frac{P(x)}{Q(x)} = \frac{P(c)}{Q(c)}.$$

24. Limits



Example

$$\lim_{x \rightarrow -1} \frac{x^3 + 4x^2 - 3}{x^2 + 5} = \frac{(-1)^3 + 4(-1)^2 - 3}{(-1)^2 + 5} = \frac{0}{6} = 0.$$



Eliminating Zero Denominators Algebraically

$$\lim_{x \rightarrow c} \frac{P(x)}{Q(x)}$$

What can we do if $Q(c) = 0$?

24. Limits



Example

$$\lim_{x \rightarrow 1} \frac{x^2 + x - 2}{x^2 - x}$$

If we just put in $x = 1$, we would get “ $\frac{0}{0}$ ” and we never never never want “ $\frac{0}{0}$ ”.


Instead, we try to factor $x^2 + x - 2$ and $x^2 - x$. If $x \neq 1$, we have that

$$\frac{x^2 + x - 2}{x^2 - x} = \frac{(x - 1)(x + 2)}{x(x - 1)} = \frac{x + 2}{x}.$$

So

$$\lim_{x \rightarrow 1} \frac{x^2 + x - 2}{x^2 - x} = \lim_{x \rightarrow 1} \frac{x + 2}{x} = \frac{1 + 2}{1} = 3.$$

The Sandwich Theorem



24. Limits



Theorem (The Sandwich Theorem)

Suppose that

- $f(x) \leq g(x) \leq h(x)$ for all x “close” to c ($x \neq c$); and
- $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x) = L$.

Then

$$\lim_{x \rightarrow c} g(x) = L$$

also.

24. Limits

Example

The inequality

$$1 - \frac{x^2}{6} < \frac{x \sin x}{2 - 2 \cos x} < 1$$

holds for all x close to 0 ($x \neq 0$). Calculate $\lim_{x \rightarrow 0} \frac{x \sin x}{2 - 2 \cos x}$.

solution: Since $\lim_{x \rightarrow 0} 1 - \frac{x^2}{6} = 1 - \frac{0}{6} = 1$ and $\lim_{x \rightarrow 0} 1 = 1$, it follows by the Sandwich Theorem that $\lim_{x \rightarrow 0} \frac{x \sin x}{2 - 2 \cos x} = 1$.

24. Limits



Theorem

If

- $f(x) \leq g(x)$ for all x close to c ($x \neq c$);
- $\lim_{x \rightarrow c} f(x)$ exists; and
- $\lim_{x \rightarrow c} g(x)$ exists,


then

$$\lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} g(x).$$




Continuity

25. Continuity



25. Continuity



25. Continuity



Definition


The function $f : D \rightarrow \mathbb{R}$ is *continuous at $c \in D$* if

- $f(c)$ exists;
- $\lim_{x \rightarrow c} f(x)$ exists; and
- $\lim_{x \rightarrow c} f(x) = f(c)$.

Definition

If f is not continuous at c , we say that f is *discontinuous at c* – we say that c is a *point of discontinuity* of f .


25. Continuity



Example

Consider the function $f : [0, 4] \rightarrow \mathbb{R}$ above. Where is f continuous? Where is f discontinuous?


25. Continuity



solution:

c	Is f continuous at c ?	Why?
0	Yes	because $\lim_{x \rightarrow 0} f(x) = 1 = f(0)$
$(0, 1)$	Yes	because $\lim_{x \rightarrow c} f(x) = f(c)$
1	No	because $\lim_{x \rightarrow 1} f(x)$ does not exist

25. Continuity




solution:

c	Is f continuous at c ?	Why?
(1, 2)	Yes	because $\lim_{x \rightarrow c} f(x) = f(c)$
2	No	because $\lim_{x \rightarrow 2} f(x) = 1 \neq 2 = f(2)$
(2, 4)	Yes	because $\lim_{x \rightarrow c} f(x) = f(c)$
4	No	because $\lim_{x \rightarrow 4} f(x) = 1 \neq \frac{1}{2} = f(4)$

25. Continuity

Example

$$f : [-2, 2] \rightarrow \mathbb{R}, f(x) = \sqrt{4 - x^2}$$




f is continuous at every $c \in [-2, 2]$.

25. Continuity

Example

$$g : \mathbb{R} \rightarrow \mathbb{R}, g(x) = \begin{cases} 0 & x < 0 \\ 1 & x \geq 0 \end{cases}$$




g has a point of discontinuity at $c = 0$. g is continuous at every point $c \neq 0$.

25. Continuity

Example

$$h : \mathbb{R} \rightarrow \mathbb{R}, h(x) = \begin{cases} \frac{1}{(x-1)^2(x-2)^2} & x \neq 1 \text{ or } 2 \\ 10 & x = 1 \text{ or } 2 \end{cases}$$



h is continuous on $(-\infty, 1)$, $(1, 2)$ and $(2, \infty)$. h has a points of discontinuity at $c = 1$ and $c = 2$.

Continuous Functions

Definition

$f : D \rightarrow \mathbb{R}$ is a *continuous function* if it is continuous at every $c \in D$.

25. Continuity



Theorem

If f and g are continuous at c , then $f + g$, $f - g$, kf ($k \in \mathbb{R}$), fg , $\frac{f}{g}$ (if $g(c) \neq 0$) and f^n ($n \in \mathbb{N}$) are all continuous at c . If $\sqrt[n]{f}$ is defined on $(c - \delta, c + \delta)$, then $\sqrt[n]{f}$ is also continuous at c ($n \in \mathbb{N}$).

25. Continuity



Example

Every polynomial

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

is continuous.

25. Continuity



Example

If

- P and Q are polynomials; and
- $Q(c) \neq 0$,

then $\frac{P(x)}{Q(x)}$ is continuous at c .

25. Continuity



Example


$\sin x$ and $\cos x$ are continuous.

Composites

$$g \circ f(x)$$

$g \circ f(x)$ means $g(f(x))$.

25. Continuity



Theorem

If

- f is continuous at c ; and
- g is continuous at $f(c)$,

then $g \circ f$ is continuous at c .

25. Continuity

Example

Show that $h(x) = \sqrt{x^2 - 2x - 5}$ is continuous on its domain.

solution: The function $g(t) = \sqrt{t}$ is continuous by Theorem 24. The function $f(x) = x^2 - 2x - 5$ is continuous because all polynomials are continuous. Therefore $h(x) = g \circ f(x)$ is continuous.

25. Continuity



Example

Show that $\frac{x^{\frac{2}{3}}}{1+x^4}$ is continuous.

solution: $x^{\frac{2}{3}}$ and $1 + x^4$ are continuous. Because $1 + x^4 \neq 0$ for all x , we have that $\frac{x^{\frac{2}{3}}}{1+x^4}$ is continuous.

25. Continuity

Theorem

If

- $g(x)$ is continuous at $x = b$; and
- $\lim_{x \rightarrow c} f(x) = b$,

then

$$\lim_{x \rightarrow c} g(f(x)) = g\left(\lim_{x \rightarrow c} f(x)\right).$$

25. Continuity

Example

$$\begin{aligned}& \lim_{x \rightarrow \frac{\pi}{2}} \cos \left[2x + \sin \left(\frac{3\pi}{2} + x \right) \right] \\&= \cos \left[\lim_{x \rightarrow \frac{\pi}{2}} \left(2x + \sin \left(\frac{3\pi}{2} + x \right) \right) \right] \\&= \cos \left[\lim_{x \rightarrow \frac{\pi}{2}} (2x) + \lim_{x \rightarrow \frac{\pi}{2}} \left(\sin \left(\frac{3\pi}{2} + x \right) \right) \right] \\&= \cos \left[\pi + \sin \left(\lim_{x \rightarrow \frac{\pi}{2}} \left(\frac{3\pi}{2} + x \right) \right) \right] \\&= \cos [\pi + \sin 2\pi] = \cos [\pi + 0] = -1.\end{aligned}$$




Differentiation


26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation



26. Differentiation




26. Differentiation



26. Differentiation

Example


$$y = x^2$$




26. Differentiation

Example


$$y = x^2$$



26. Differentiation



26. Differentiation




26. Differentiation




How can we calculate the slope of the tangent line?


26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation




26. Differentiation



26. Differentiation



26. Differentiation



26. Differentiation



If h is very very small, then

$$\left(\begin{array}{l} \text{slope of the} \\ \text{tangent line} \end{array} \right) \approx \left(\begin{array}{l} \text{slope of the} \\ \text{orange line} \end{array} \right) = \frac{f(x_0 + h) - f(x_0)}{h}$$

The Derivative of f

Definition

The *derivative of a function f at a point x_0* is

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

if the limit exists.

(f' is pronounced “ f prime”)

26. Differentiation

Example


Consider the function $g(x) = \frac{1}{x}$, $x \neq 0$.

If $x_0 \neq 0$, then

$$\begin{aligned} g'(x_0) &= \lim_{h \rightarrow 0} \frac{g(x_0 + h) - g(x_0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{1}{x_0+h} - \frac{1}{x_0}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\left(\frac{x_0}{x_0(x_0+h)} - \frac{x_0+h}{x_0(x_0+h)} \right)}{h} \\ &= \lim_{h \rightarrow 0} \frac{x_0 - x_0 - h}{hx_0(x_0 + h)} \\ &= \lim_{h \rightarrow 0} \frac{-1}{x_0(x_0 + h)} = -\frac{1}{x_0^2}. \end{aligned}$$

26. Differentiation

$$g'(x_0) = -\frac{1}{x_0^2}$$



26. Differentiation



Definition

If $f'(x_0)$ exists, we say that f is differentiable at x_0 .

26. Differentiation



Definition

Let $f : D \rightarrow \mathbb{R}$ be a function. If f is differentiable at every $x_0 \in D$, we say that f is *differentiable*.

26. Differentiation




If $f : D \rightarrow \mathbb{R}$ is differentiable, then we have a new function
 $f' : D \rightarrow \mathbb{R}$.

Definition

f' is called the *derivative* of f .

26. Differentiation



Example

$$\text{Differentiate } f(x) = \frac{x}{x-1}.$$

26. Differentiation

solution: First note that $f(x + h) = \frac{x+h}{x+h-1}$. Therefore

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{x+h}{x+h-1} - \frac{x}{x-1}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{(x+h)(x-1) - x(x+h-1)}{(x-1)(x+h-1)} \right) \\
 &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{-h}{(x-1)(x+h-1)} \right) \\
 &= \lim_{h \rightarrow 0} \frac{-1}{(x-1)(x+h-1)} \\
 &= \frac{-1}{(x-1)(x+0-1)} \\
 &= \frac{-1}{(x-1)^2}.
 \end{aligned}$$

Notations


There are many ways to write the derivative of $y = f(x)$.

$$f'(x) = y' = \frac{dy}{dx} = \frac{df}{dx} = \frac{d}{dx}f(x) = \dot{y} = \dot{f}(x)$$



“the derivative of y with respect to x ”

26. Differentiation



Sir Isaac Newton
UK, 1642-1726

Gottfried Leibniz
GER, 1646-1716

Joseph-Louis Lagrange
ITA, 1736-1813

Calculus was started by two men who hated each other: Sir Isaac Newton used f' and \dot{y} . Gottfried Leibniz used $\frac{df}{dx}$ and $\frac{dy}{dx}$.

The f' and y' notation came later from Joseph-Louis Lagrange.

26. Differentiation



If we want the derivative of $y = f(x)$ at the point $x = x_0$, we can write

$$f'(x_0) = \frac{dy}{dx} \Big|_{x=x_0} = \frac{df}{dx} \Big|_{x=x_0} = \frac{d}{dx} f(x) \Big|_{x=x_0}$$



“the derivative of y with respect to x at $x = x_0$ ”

26. Differentiation



For example, if $u(x) = \frac{1}{x}$, then

$$u'(4) = \left. \frac{d}{dx} \left(\frac{1}{x} \right) \right|_{x=4} = \left. \frac{-1}{x^2} \right|_{x=4} = \frac{-1}{4^2} = \frac{-1}{16}.$$

26. Differentiation



Example

Show that $f(x) = |x|$ is differentiable on $(-\infty, 0)$ and on $(0, \infty)$, but is not differentiable at $x = 0$.

26. Differentiation



solution: If $x > 0$ then

$$\frac{df}{dx} = \frac{d}{dx}(|x|) = \frac{d}{dx}(x) = \lim_{h \rightarrow 0} \frac{(x+h)-x}{h} = \lim_{h \rightarrow 0} 1 = 1.$$

26. Differentiation




Similarly, if $x < 0$ then

$$\begin{aligned}\frac{df}{dx} &= \frac{d}{dx}(|x|) = \frac{d}{dx}(-x) = \lim_{h \rightarrow 0} \frac{(-x - h) - (-x)}{h} \\ &= \lim_{h \rightarrow 0} -1 = -1.\end{aligned}$$


Therefore f is differentiable on $(-\infty, 0)$ and on $(0, \infty)$.

26. Differentiation


Since $\lim_{h \rightarrow 0} \frac{|0+h| - |0|}{h} = \lim_{h \rightarrow 0} \frac{|h|}{h} = \lim_{h \rightarrow 0} (\pm 1)$ does not exist, f is not differentiable at 0.




When Does a Function Not Have a Derivative at a Point?




When Does a Function Not Have a Derivative at a Point?




When Does a Function Not Have a Derivative at a Point?



When Does a Function Not Have a Derivative at a Point?



When Does a Function Not Have a Derivative at a Point?



26. Differentiation



Theorem

$$\left(\begin{array}{c} f \text{ has a derivative} \\ \text{at } x = x_0 \end{array} \right) \implies \left(\begin{array}{c} f \text{ is continuous} \\ \text{at } x = x_0 \end{array} \right)$$



Next Time

- 27. Differentiation Rules
- 28. Derivatives of Trigonometric Functions
- 29. The Chain Rule