

Lecture 10

- 5.3 Homogeneous Linear Systems with Constant Coefficients
- 5.4 Complex Eigenvalues
- 5.5 Fundamental Matrices



Homogeneous Linear Systems with Constant Coefficients

5.3 Homogeneous Linear Systems with Constant Coefficients



Consider

$$\mathbf{x}' = A\mathbf{x}$$

where $A \in \mathbb{R}^{n \times n}$.

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}' = A\mathbf{x}$$

If $n = 1$, then we just have

$$\frac{dx}{dt} = ax$$

which has general solution $x(t) = ce^{at}$.

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}' = A\mathbf{x}$$

If $n = 1$, then we just have

$$\frac{dx}{dt} = ax$$

which has general solution $x(t) = ce^{at}$.

For $n > 1$, we guess that

$$\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$$

is a solution to $\mathbf{x}' = A\mathbf{x}$, for some number $r \in \mathbb{C}$ and some vector $\boldsymbol{\xi} \in \mathbb{C}^n$.

5.3 Homogeneous Linear Systems with Constant Coefficients



But if $\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$, then

$$\mathbf{x}' = A\mathbf{x}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



But if $\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$, then

$$r\boldsymbol{\xi}e^{rt} = \mathbf{x}' = A\mathbf{x}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



But if $\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$, then

$$r\boldsymbol{\xi}e^{rt} = \mathbf{x}' = A\mathbf{x} = A\boldsymbol{\xi}e^{rt}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



But if $\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$, then

$$r\boldsymbol{\xi}e^{rt} = \mathbf{x}' = A\mathbf{x} = A\boldsymbol{\xi}e^{rt}$$

$$r\boldsymbol{\xi} = A\boldsymbol{\xi}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



But if $\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$, then

$$r\boldsymbol{\xi}e^{rt} = \mathbf{x}' = A\mathbf{x} = A\boldsymbol{\xi}e^{rt}$$

$$r\boldsymbol{\xi} = A\boldsymbol{\xi}$$

$$(A - rI)\boldsymbol{\xi} = \mathbf{0}$$

where I is the identity matrix.

5.3 Homogeneous Linear Systems with Constant Coefficients



But if $\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$, then

$$r\boldsymbol{\xi}e^{rt} = \mathbf{x}' = A\mathbf{x} = A\boldsymbol{\xi}e^{rt}$$

$$r\boldsymbol{\xi} = A\boldsymbol{\xi}$$

$$(A - rI)\boldsymbol{\xi} = \mathbf{0}$$

where I is the identity matrix. Hence r must be an eigenvalue of A and $\boldsymbol{\xi}$ must be a corresponding eigenvector of A .

5.3 Homogeneous Linear Systems with Constant Coefficients



Remark

So the idea is:

- 1 Find the eigenvalues;
- 2 Find the eigenvectors; then
- 3 Write $\mathbf{x}^{(j)}(t) = \boldsymbol{\xi}^{(j)} e^{r_j t}$.

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

First we find the eigenvalues.

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

First we find the eigenvalues. Since

$$\begin{aligned} 0 &= \det(A - rI) = \begin{vmatrix} 1-r & 1 \\ 4 & 1-r \end{vmatrix} = (1-r)^2 - 4 \\ &= r^2 - 2r - 3 = (r+1)(r-3), \end{aligned}$$

the eigenvalues are $r_1 = 3$ and $r_2 = -1$.

5.3 Homogeneous Linear Systems with Constant Coefficients



Using the first eigenvalue $r_1 = 3$, we calculate that

$$\mathbf{0} = (A - r_1 I)\boldsymbol{\xi} = \begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Using the first eigenvalue $r_1 = 3$, we calculate that

$$\mathbf{0} = (A - r_1 I)\boldsymbol{\xi} = \begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Longrightarrow \quad 0 = -2\xi_1 + \xi_2.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Using the first eigenvalue $r_1 = 3$, we calculate that

$$\mathbf{0} = (A - r_1 I)\boldsymbol{\xi} = \begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Rightarrow \quad 0 = -2\xi_1 + \xi_2.$$

Hence we can choose $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

5.3 Homogeneous Linear Systems with Constant Coefficients



Using the first eigenvalue $r_1 = 3$, we calculate that

$$\mathbf{0} = (A - r_1 I)\boldsymbol{\xi} = \begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Rightarrow \quad 0 = -2\xi_1 + \xi_2.$$

Hence we can choose $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$. Then using the second eigenvalue $r_2 = -1$, we calculate that

$$\mathbf{0} = (A - r_2 I)\boldsymbol{\xi} = \begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Using the first eigenvalue $r_1 = 3$, we calculate that

$$\mathbf{0} = (A - r_1 I)\boldsymbol{\xi} = \begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Longrightarrow \quad 0 = -2\xi_1 + \xi_2.$$

Hence we can choose $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$. Then using the second eigenvalue $r_2 = -1$, we calculate that

$$\mathbf{0} = (A - r_2 I)\boldsymbol{\xi} = \begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Longrightarrow \quad 0 = 2\xi_1 + \xi_2.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Using the first eigenvalue $r_1 = 3$, we calculate that

$$\mathbf{0} = (A - r_1 I)\boldsymbol{\xi} = \begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Longrightarrow \quad 0 = -2\xi_1 + \xi_2.$$

Hence we can choose $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$. Then using the second eigenvalue $r_2 = -1$, we calculate that

$$\mathbf{0} = (A - r_2 I)\boldsymbol{\xi} = \begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Longrightarrow \quad 0 = 2\xi_1 + \xi_2.$$

Hence we can choose $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$.

5.3 Homogeneous Linear Systems with Constant Coefficients



Using the first eigenvalue $r_1 = 3$, we calculate that

$$\mathbf{0} = (A - r_1 I)\boldsymbol{\xi} = \begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Longrightarrow \quad 0 = -2\xi_1 + \xi_2.$$

Hence we can choose $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$. Then using the second eigenvalue $r_2 = -1$, we calculate that

$$\mathbf{0} = (A - r_2 I)\boldsymbol{\xi} = \begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \Longrightarrow \quad 0 = 2\xi_1 + \xi_2.$$

Hence we can choose $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$. This gives us two solutions:

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



But are these two solutions linearly independent?

5.3 Homogeneous Linear Systems with Constant Coefficients



But are these two solutions linearly independent? To find out, we calculate the Wronskian of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$:

5.3 Homogeneous Linear Systems with Constant Coefficients



But are these two solutions linearly independent? To find out, we calculate the Wronskian of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$:

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} = \begin{bmatrix} e^{3t} \\ 2e^{3t} \end{bmatrix} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t} = \begin{bmatrix} e^{-t} \\ -2e^{-t} \end{bmatrix}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



But are these two solutions linearly independent? To find out, we calculate the Wronskian of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$:

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} = \begin{bmatrix} e^{3t} \\ 2e^{3t} \end{bmatrix} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t} = \begin{bmatrix} e^{-t} \\ -2e^{-t} \end{bmatrix}.$$

$$W(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})(t) = \begin{vmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{vmatrix} = -4e^{2t} \neq 0.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



But are these two solutions linearly independent? To find out, we calculate the Wronskian of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$:

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} = \begin{bmatrix} e^{3t} \\ 2e^{3t} \end{bmatrix} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t} = \begin{bmatrix} e^{-t} \\ -2e^{-t} \end{bmatrix}.$$

$$W(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})(t) = \begin{vmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{vmatrix} = -4e^{2t} \neq 0.$$

Since $W \neq 0$, we have that $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ are linearly independent.

5.3 Homogeneous Linear Systems with Constant Coefficients



But are these two solutions linearly independent? To find out, we calculate the Wronskian of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$:

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} = \begin{bmatrix} e^{3t} \\ 2e^{3t} \end{bmatrix} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t} = \begin{bmatrix} e^{-t} \\ -2e^{-t} \end{bmatrix}.$$

$$W(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})(t) = \begin{vmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{vmatrix} = -4e^{2t} \neq 0.$$

Since $W \neq 0$, we have that $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ are linearly independent. So $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ form a fundamental set of solutions.

5.3 Homogeneous Linear Systems with Constant Coefficients



But are these two solutions linearly independent? To find out, we calculate the Wronskian of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$:

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} = \begin{bmatrix} e^{3t} \\ 2e^{3t} \end{bmatrix} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t} = \begin{bmatrix} e^{-t} \\ -2e^{-t} \end{bmatrix}.$$

$$W(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})(t) = \begin{vmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{vmatrix} = -4e^{2t} \neq 0.$$

Since $W \neq 0$, we have that $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ are linearly independent. So $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ form a fundamental set of solutions. Therefore the general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} 8 & -1 \\ 6 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 1 \\ -2 \end{bmatrix}. \end{cases}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} 8 & -1 \\ 6 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 1 \\ -2 \end{bmatrix}. \end{cases}$$

The eigenvalues are $r_1 = 7$ and $r_2 = 2$.

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} 8 & -1 \\ 6 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 1 \\ -2 \end{bmatrix}. \end{cases}$$

The eigenvalues are $r_1 = 7$ and $r_2 = 2$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ 6 \end{bmatrix}$.

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} 8 & -1 \\ 6 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 1 \\ -2 \end{bmatrix}. \end{cases}$$

The eigenvalues are $r_1 = 7$ and $r_2 = 2$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ 6 \end{bmatrix}$. Therefore the general solution to the ODE is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

Setting $t = 0$, we have

$$\begin{bmatrix} 1 \\ -2 \end{bmatrix} = \mathbf{x}(0)$$

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

Setting $t = 0$, we have

$$\begin{bmatrix} 1 \\ -2 \end{bmatrix} = \mathbf{x}(0) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} = \begin{bmatrix} c_1 + c_2 \\ c_1 + 6c_2 \end{bmatrix}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

Setting $t = 0$, we have

$$\begin{bmatrix} 1 \\ -2 \end{bmatrix} = \mathbf{x}(0) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} = \begin{bmatrix} c_1 + c_2 \\ c_1 + 6c_2 \end{bmatrix} \implies \begin{cases} c_1 + c_2 = 1 \\ c_1 + 6c_2 = -2 \end{cases}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

Setting $t = 0$, we have

$$\begin{aligned} \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \mathbf{x}(0) &= c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} = \begin{bmatrix} c_1 + c_2 \\ c_1 + 6c_2 \end{bmatrix} \implies \begin{cases} c_1 + c_2 = 1 \\ c_1 + 6c_2 = -2 \end{cases} \\ &\implies \begin{cases} c_1 = \frac{8}{5} \\ c_2 = -\frac{3}{5}. \end{cases} \end{aligned}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

Setting $t = 0$, we have

$$\begin{aligned} \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \mathbf{x}(0) &= c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} = \begin{bmatrix} c_1 + c_2 \\ c_1 + 6c_2 \end{bmatrix} \implies \begin{cases} c_1 + c_2 = 1 \\ c_1 + 6c_2 = -2 \end{cases} \\ &\implies \begin{cases} c_1 = \frac{8}{5} \\ c_2 = -\frac{3}{5} \end{cases}. \end{aligned}$$

Therefore the solution to the IVP is

$$\mathbf{x}(t) = \frac{8}{5} \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} - \frac{3}{5} \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} \mathbf{x}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} \mathbf{x}.$$

The eigenvalues are $r_1 = -1$ and $r_2 = -4$.

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} \mathbf{x}.$$

The eigenvalues are $r_1 = -1$ and $r_2 = -4$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ \sqrt{2} \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$.

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} \mathbf{x}.$$

The eigenvalues are $r_1 = -1$ and $r_2 = -4$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ \sqrt{2} \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$. Hence the general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ \sqrt{2} \end{bmatrix} e^{-t} + c_2 \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix} e^{-4t}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Remark

$$\det(A - rI) = 0$$

There are three possibilities for the eigenvalues of A .

5.3 Homogeneous Linear Systems with Constant Coefficients



Remark

$$\det(A - rI) = 0$$

There are three possibilities for the eigenvalues of A .

- 1 All the eigenvalues are real and different;
- 2 Some eigenvalues occur in complex conjugate pairs;
- 3 Some eigenvalues are repeated.

5.3 Homogeneous Linear Systems with Constant Coefficients



If all the eigenvalues are real and different, then the eigenvectors are linearly independent:

5.3 Homogeneous Linear Systems with Constant Coefficients



If all the eigenvalues are real and different, then the eigenvectors are linearly independent:

So $W(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)})(t) \neq 0$ and $\mathbf{x}^{(1)}(t), \dots, \mathbf{x}^{(n)}(t)$ form a fundamental set of solutions.

5.3 Homogeneous Linear Systems with Constant Coefficients



If some eigenvalues are repeated, *but there are n linearly independent eigenvectors*, then this is also true: $\mathbf{x}^{(1)}(t), \dots, \mathbf{x}^{(n)}(t)$ form a fundamental set of solutions.

5.3 Homogeneous Linear Systems with Constant Coefficients



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \mathbf{x}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



The eigenvalues and eigenvectors are

$$r_1 = 2$$
$$\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$r_2 = -1$$
$$\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$$r_3 = -1$$
$$\boldsymbol{\xi}^{(3)} = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$$

5.3 Homogeneous Linear Systems with Constant Coefficients



The eigenvalues and eigenvectors are

$$\begin{array}{lll} r_1 = 2 & r_2 = -1 & r_3 = -1 \\ \boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} & \boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} & \boldsymbol{\xi}^{(3)} = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \end{array}$$

which gives us the following three solutions

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} e^{2t} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} e^{-t} \quad \mathbf{x}^{(3)}(t) = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} e^{-t}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} e^{2t} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} e^{-t} \quad \mathbf{x}^{(3)}(t) = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} e^{-t}.$$

You can check that the Wronskian of $\mathbf{x}^{(1)}$, $\mathbf{x}^{(2)}$ and $\mathbf{x}^{(3)}$ is non-zero.

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} e^{2t} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} e^{-t} \quad \mathbf{x}^{(3)}(t) = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} e^{-t}.$$

You can check that the Wronskian of $\mathbf{x}^{(1)}$, $\mathbf{x}^{(2)}$ and $\mathbf{x}^{(3)}$ is non-zero. Therefore $\mathbf{x}^{(1)}$, $\mathbf{x}^{(2)}$ and $\mathbf{x}^{(3)}$ form a fundamental set of solutions.

5.3 Homogeneous Linear Systems with Constant Coefficients



$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} e^{2t} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} e^{-t} \quad \mathbf{x}^{(3)}(t) = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} e^{-t}.$$

You can check that the Wronskian of $\mathbf{x}^{(1)}$, $\mathbf{x}^{(2)}$ and $\mathbf{x}^{(3)}$ is non-zero. Therefore $\mathbf{x}^{(1)}$, $\mathbf{x}^{(2)}$ and $\mathbf{x}^{(3)}$ form a fundamental set of solutions. The general solution to the ODE is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} e^{2t} + c_2 \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} e^{-t} + c_3 \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} e^{-t}.$$

5.3 Homogeneous Linear Systems with Constant Coefficients



Remark

So if we have repeated eigenvalues with n linearly independent eigenvectors then there is no problem.

In section 5.6 we will study what to do if we do not have enough eigenvectors.

5.3 Homogeneous Linear Systems with Constant Coefficients



Remark

So if we have repeated eigenvalues with n linearly independent eigenvectors then there is no problem.

In section 5.6 we will study what to do if we do not have enough eigenvectors.

Remark

Next we will study systems with complex eigenvalues.

Complex Eigenvalues

5.4 Complex Eigenvalues



Consider

$$\mathbf{x}' = A\mathbf{x}$$

where $A \in \mathbb{R}^{n \times n}$.

5.4 Complex Eigenvalues



Consider

$$\mathbf{x}' = A\mathbf{x}$$

where $A \in \mathbb{R}^{n \times n}$.

$$\begin{bmatrix} x'_1 \\ x'_2 \\ \vdots \\ x'_n \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

5.4 Complex Eigenvalues



Consider

$$\mathbf{x}' = A\mathbf{x}$$

where $A \in \mathbb{R}^{n \times n}$.

$$\begin{cases} x'_1 = a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n \\ x'_2 = a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n \\ \vdots \\ x'_n = a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n \end{cases}$$

5.4 Complex Eigenvalues



Any complex eigenvalues of A must occur in complex conjugate pairs: If $r_1 = \lambda + i\mu$ is an eigenvalue of A , then $r_2 = \bar{r}_1 = \lambda - i\mu$ is also an eigenvalue of A .

5.4 Complex Eigenvalues



Moreover, if $\xi^{(1)}$ is an eigenvector of A corresponding to r_1 , then $\xi^{(2)} = \overline{\xi^{(1)}}$ is an eigenvector of A corresponding to $r_2 = \bar{r}_1$.

5.4 Complex Eigenvalues



Two solutions of $\mathbf{x}' = A\mathbf{x}$ are

$$\mathbf{x}^{(1)}(t) = \boldsymbol{\xi}^{(1)} e^{r_1 t} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \overline{\boldsymbol{\xi}^{(1)}} e^{\bar{r}_1 t}.$$

5.4 Complex Eigenvalues



Two solutions of $\mathbf{x}' = A\mathbf{x}$ are

$$\mathbf{x}^{(1)}(t) = \boldsymbol{\xi}^{(1)} e^{r_1 t} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \overline{\boldsymbol{\xi}^{(1)}} e^{\bar{r}_1 t}.$$

But $\mathbf{x}^{(1)}, \mathbf{x}^{(2)} : \mathbb{R} \rightarrow \mathbb{C}^n$ and we want solutions $: \mathbb{R} \rightarrow \mathbb{R}^n$.

5.4 Complex Eigenvalues



If $r_1 = \lambda + i\mu$, and $\boldsymbol{\xi}^{(1)} = \mathbf{a} + i\mathbf{b}$ ($\lambda, \mu \in \mathbb{R}$, $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$), then

$$\mathbf{x}^{(1)}(t) = (\mathbf{a} + i\mathbf{b})e^{(\lambda+i\mu)t}$$

5.4 Complex Eigenvalues



If $r_1 = \lambda + i\mu$, and $\boldsymbol{\xi}^{(1)} = \mathbf{a} + i\mathbf{b}$ ($\lambda, \mu \in \mathbb{R}$, $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$), then

$$\begin{aligned}\mathbf{x}^{(1)}(t) &= (\mathbf{a} + i\mathbf{b})e^{(\lambda+i\mu)t} \\ &= (\mathbf{a} + i\mathbf{b})e^{\lambda t} (\cos \mu t + i \sin \mu t)\end{aligned}$$

5.4 Complex Eigenvalues



If $r_1 = \lambda + i\mu$, and $\boldsymbol{\xi}^{(1)} = \mathbf{a} + i\mathbf{b}$ ($\lambda, \mu \in \mathbb{R}$, $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$), then

$$\begin{aligned}\mathbf{x}^{(1)}(t) &= (\mathbf{a} + i\mathbf{b})e^{(\lambda+i\mu)t} \\ &= (\mathbf{a} + i\mathbf{b})e^{\lambda t} (\cos \mu t + i \sin \mu t) \\ &= e^{\lambda t} (\mathbf{a} \cos \mu t - \mathbf{b} \sin \mu t) + ie^{\lambda t} (\mathbf{a} \sin \mu t + \mathbf{b} \cos \mu t)\end{aligned}$$

5.4 Complex Eigenvalues



If $r_1 = \lambda + i\mu$, and $\boldsymbol{\xi}^{(1)} = \mathbf{a} + i\mathbf{b}$ ($\lambda, \mu \in \mathbb{R}$, $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$), then

$$\begin{aligned}\mathbf{x}^{(1)}(t) &= (\mathbf{a} + i\mathbf{b})e^{(\lambda+i\mu)t} \\ &= (\mathbf{a} + i\mathbf{b})e^{\lambda t} (\cos \mu t + i \sin \mu t) \\ &= e^{\lambda t} (\mathbf{a} \cos \mu t - \mathbf{b} \sin \mu t) + ie^{\lambda t} (\mathbf{a} \sin \mu t + \mathbf{b} \cos \mu t) \\ &= \mathbf{u}(t) + i\mathbf{v}(t).\end{aligned}$$

5.4 Complex Eigenvalues



Remark

- The functions $\mathbf{u}(t)$ and $\mathbf{v}(t)$ solve the ODE.

5.4 Complex Eigenvalues



Remark

- The functions $\mathbf{u}(t)$ and $\mathbf{v}(t)$ solve the ODE.
- The functions $\mathbf{u}(t)$ and $\mathbf{v}(t)$ will be linearly independent.

Remark

- The functions $\mathbf{u}(t)$ and $\mathbf{v}(t)$ solve the ODE.
- The functions $\mathbf{u}(t)$ and $\mathbf{v}(t)$ will be linearly independent.
- $\text{span}\{\mathbf{u}(t), \mathbf{v}(t)\} = \text{span}\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}\}$.

Remark

- The functions $\mathbf{u}(t)$ and $\mathbf{v}(t)$ solve the ODE.
- The functions $\mathbf{u}(t)$ and $\mathbf{v}(t)$ will be linearly independent.
- $\text{span}\{\mathbf{u}(t), \mathbf{v}(t)\} = \text{span}\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}\}$.

So we can include $\mathbf{u}(t)$ and $\mathbf{v}(t)$ in our fundamental set of solutions instead of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$.

5.4 Complex Eigenvalues



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} -\frac{1}{2} & 1 \\ -1 & -\frac{1}{2} \end{bmatrix} \mathbf{x}.$$

5.4 Complex Eigenvalues



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} -\frac{1}{2} & 1 \\ -1 & -\frac{1}{2} \end{bmatrix} \mathbf{x}.$$

We calculate that

$$0 = \det(A - rI) = \begin{vmatrix} -\frac{1}{2} - r & 1 \\ -1 & -\frac{1}{2} - r \end{vmatrix} = r^2 + r + \frac{5}{4}$$

and

$$r_{1,2} = \frac{-1 \pm \sqrt{1-5}}{2} = \frac{-1 \pm 2i}{2} = -\frac{1}{2} \pm i.$$

5.4 Complex Eigenvalues



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} -\frac{1}{2} & 1 \\ -1 & -\frac{1}{2} \end{bmatrix} \mathbf{x}.$$

We calculate that

$$0 = \det(A - rI) = \begin{vmatrix} -\frac{1}{2} - r & 1 \\ -1 & -\frac{1}{2} - r \end{vmatrix} = r^2 + r + \frac{5}{4}$$

and

$$r_{1,2} = \frac{-1 \pm \sqrt{1-5}}{2} = \frac{-1 \pm 2i}{2} = -\frac{1}{2} \pm i.$$

So we have $r_1 = -\frac{1}{2} + i$ and $r_2 = -\frac{1}{2} - i$.

5.4 Complex Eigenvalues



Example

Solve

$$\mathbf{x}' = \begin{bmatrix} -\frac{1}{2} & 1 \\ -1 & -\frac{1}{2} \end{bmatrix} \mathbf{x}.$$

We calculate that

$$0 = \det(A - rI) = \begin{vmatrix} -\frac{1}{2} - r & 1 \\ -1 & -\frac{1}{2} - r \end{vmatrix} = r^2 + r + \frac{5}{4}$$

and

$$r_{1,2} = \frac{-1 \pm \sqrt{1-5}}{2} = \frac{-1 \pm 2i}{2} = -\frac{1}{2} \pm i.$$

So we have $r_1 = -\frac{1}{2} + i$ and $r_2 = -\frac{1}{2} - i$. We will use r_1 . We do not need r_2 .

5.4 Complex Eigenvalues



Since

$$0 = (A - r_1 I)\boldsymbol{\xi}^{(1)} = \begin{bmatrix} -i & 1 \\ -1 & -i \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix}$$

5.4 Complex Eigenvalues



Since

$$0 = (A - r_1 I)\boldsymbol{\xi}^{(1)} = \begin{bmatrix} -i & 1 \\ -1 & -i \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \implies \begin{cases} -i\xi_1 + \xi_2 = 0 \\ -\xi_1 - i\xi_2 = 0 \end{cases}$$

5.4 Complex Eigenvalues



Since

$$0 = (A - r_1 I)\boldsymbol{\xi}^{(1)} = \begin{bmatrix} -i & 1 \\ -1 & -i \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \implies \begin{cases} -i\xi_1 + \xi_2 = 0 \\ -\xi_1 - i\xi_2 = 0 \end{cases}$$

we can choose

$$\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ i \end{bmatrix}.$$

5.4 Complex Eigenvalues



Note that we also have

$$\boldsymbol{\xi}^{(2)} = \overline{\boldsymbol{\xi}^{(1)}} = \overline{\begin{bmatrix} 1 \\ i \end{bmatrix}} = \begin{bmatrix} 1 \\ -i \end{bmatrix},$$

5.4 Complex Eigenvalues



Note that we also have

$$\boldsymbol{\xi}^{(2)} = \overline{\boldsymbol{\xi}^{(1)}} = \overline{\begin{bmatrix} 1 \\ i \end{bmatrix}} = \begin{bmatrix} 1 \\ -i \end{bmatrix},$$

but we don't need $\boldsymbol{\xi}^{(2)}$.

5.4 Complex Eigenvalues



Next we look at $\mathbf{x}^{(1)}(t)$:

5.4 Complex Eigenvalues



Next we look at $\mathbf{x}^{(1)}(t)$:

$$\mathbf{x}^{(1)}(t) = \boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 1 \\ i \end{bmatrix} e^{-\frac{t}{2}} (\cos t + i \sin t)$$

=

=

5.4 Complex Eigenvalues



Next we look at $\mathbf{x}^{(1)}(t)$:

$$\begin{aligned}\mathbf{x}^{(1)}(t) &= \boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 1 \\ i \end{bmatrix} e^{-\frac{t}{2}} (\cos t + i \sin t) \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t + i \sin t \\ i \cos t - \sin t \end{bmatrix} \\ &= \end{aligned}$$

5.4 Complex Eigenvalues



Next we look at $\mathbf{x}^{(1)}(t)$:

$$\begin{aligned}\mathbf{x}^{(1)}(t) &= \boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 1 \\ i \end{bmatrix} e^{-\frac{t}{2}} (\cos t + i \sin t) \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t + i \sin t \\ i \cos t - \sin t \end{bmatrix} \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} + i e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}\end{aligned}$$

5.4 Complex Eigenvalues



Next we look at $\mathbf{x}^{(1)}(t)$:

$$\begin{aligned}\mathbf{x}^{(1)}(t) &= \boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 1 \\ i \end{bmatrix} e^{-\frac{t}{2}} (\cos t + i \sin t) \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t + i \sin t \\ i \cos t - \sin t \end{bmatrix} \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} + i e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix} \\ &= \mathbf{u}(t) + i \mathbf{v}(t).\end{aligned}$$

5.4 Complex Eigenvalues



Next we look at $\mathbf{x}^{(1)}(t)$:

$$\begin{aligned}\mathbf{x}^{(1)}(t) &= \boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 1 \\ i \end{bmatrix} e^{-\frac{t}{2}} (\cos t + i \sin t) \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t + i \sin t \\ i \cos t - \sin t \end{bmatrix} \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} + i e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix} \\ &= \mathbf{u}(t) + i \mathbf{v}(t).\end{aligned}$$

Hence we choose

$$\mathbf{u}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} \quad \text{and} \quad \mathbf{v}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}.$$

5.4 Complex Eigenvalues



$$\mathbf{u}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} \quad \text{and} \quad \mathbf{v}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}.$$

But are $\mathbf{u}(t)$ and $\mathbf{v}(t)$ linearly independent?

5.4 Complex Eigenvalues



$$\mathbf{u}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} \quad \text{and} \quad \mathbf{v}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}.$$

But are $\mathbf{u}(t)$ and $\mathbf{v}(t)$ linearly independent? Since

$$\begin{aligned} W(\mathbf{u}(t), \mathbf{v}(t))(t) &= \begin{vmatrix} u_1 & v_1 \\ u_2 & v_2 \end{vmatrix} = \begin{vmatrix} e^{-\frac{t}{2}} \cos t & e^{-\frac{t}{2}} \sin t \\ -e^{-\frac{t}{2}} \sin t & e^{-\frac{t}{2}} \cos t \end{vmatrix} \\ &= e^{-t} \cos^2 t + e^{-t} \sin^2 t = e^{-t} \\ &\neq 0 \end{aligned}$$

the answer is yes.

5.4 Complex Eigenvalues



$$\mathbf{u}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} \quad \text{and} \quad \mathbf{v}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}.$$

But are $\mathbf{u}(t)$ and $\mathbf{v}(t)$ linearly independent? Since

$$\begin{aligned} W(\mathbf{u}(t), \mathbf{v}(t))(t) &= \begin{vmatrix} u_1 & v_1 \\ u_2 & v_2 \end{vmatrix} = \begin{vmatrix} e^{-\frac{t}{2}} \cos t & e^{-\frac{t}{2}} \sin t \\ -e^{-\frac{t}{2}} \sin t & e^{-\frac{t}{2}} \cos t \end{vmatrix} \\ &= e^{-t} \cos^2 t + e^{-t} \sin^2 t = e^{-t} \\ &\neq 0 \end{aligned}$$

the answer is yes. Therefore $\mathbf{u}(t)$ and $\mathbf{v}(t)$ form a fundamental set of solutions.

5.4 Complex Eigenvalues



$$\mathbf{u}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} \quad \text{and} \quad \mathbf{v}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}.$$

But are $\mathbf{u}(t)$ and $\mathbf{v}(t)$ linearly independent? Since

$$\begin{aligned} W(\mathbf{u}(t), \mathbf{v}(t))(t) &= \begin{vmatrix} u_1 & v_1 \\ u_2 & v_2 \end{vmatrix} = \begin{vmatrix} e^{-\frac{t}{2}} \cos t & e^{-\frac{t}{2}} \sin t \\ -e^{-\frac{t}{2}} \sin t & e^{-\frac{t}{2}} \cos t \end{vmatrix} \\ &= e^{-t} \cos^2 t + e^{-t} \sin^2 t = e^{-t} \\ &\neq 0 \end{aligned}$$

the answer is yes. Therefore $\mathbf{u}(t)$ and $\mathbf{v}(t)$ form a fundamental set of solutions.

Therefore the general solution to $\mathbf{x}' = \begin{bmatrix} -\frac{1}{2} & 1 \\ -1 & -\frac{1}{2} \end{bmatrix} \mathbf{x}$ is

$$\mathbf{x}(t) = c_1 e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} + c_2 e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}.$$

Remark


Our method is

1. Find the eigenvalues;
2. Find the eigenvectors;
3.
 - If r_j is real, just use the solution $\mathbf{x}^{(j)}(t) = \boldsymbol{\xi}^{(j)} e^{r_j t}$;
 - But if r_j is complex, write

$$\mathbf{x}^{(j)}(t) = \boldsymbol{\xi}^{(j)} e^{r_j t} = \begin{pmatrix} \text{real valued} \\ \text{function} \end{pmatrix} + i \begin{pmatrix} \text{real valued} \\ \text{function} \end{pmatrix}$$

and use these two functions.

5.4 Complex Eigenvalues



5.4 Complex Eigenvalues



Example

The electric circuit shown above is described by

$$\begin{cases} I' = -I - V \\ V' = 2I - V \end{cases}$$

where

I = the current through the inductor

V = the voltage drop across the capacitor.

(Ask an Electrical Engineer.)

5.4 Complex Eigenvalues



Example

The electric circuit shown above is described by

$$\begin{cases} I' = -I - V \\ V' = 2I - V \end{cases}$$

where

I = the current through the inductor

V = the voltage drop across the capacitor.

(Ask an Electrical Engineer.)

Suppose that at time $t = 0$ the current is 2 amperes and the voltage drop is 2 volts. Find $I(t)$ and $V(t)$.

5.4 Complex Eigenvalues



$$\begin{cases} I' = -I - V \\ V' = 2I - V \end{cases}$$

Suppose that at time $t = 0$ the current is 2 amperes and the voltage drop is 2 volts. Find $I(t)$ and $V(t)$.

We must solve the IVP

$$\begin{cases} \frac{d}{dt} \begin{bmatrix} I \\ V \end{bmatrix} = \begin{bmatrix} -1 & -1 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} I \\ V \end{bmatrix} \\ \begin{bmatrix} I \\ V \end{bmatrix} (0) = \begin{bmatrix} 2 \\ 2 \end{bmatrix} . \end{cases}$$

5.4 Complex Eigenvalues



The eigenvalues of $\begin{bmatrix} -1 & -1 \\ 2 & -1 \end{bmatrix}$ are $r_1 = -1 + i\sqrt{2}$ and $r_2 = -1 - i\sqrt{2}$ (please check). The corresponding eigenvectors are

$$\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ -i\sqrt{2} \end{bmatrix} \quad \text{and} \quad \boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ i\sqrt{2} \end{bmatrix}.$$

5.4 Complex Eigenvalues



The eigenvalues of $\begin{bmatrix} -1 & -1 \\ 2 & -1 \end{bmatrix}$ are $r_1 = -1 + i\sqrt{2}$ and $r_2 = -1 - i\sqrt{2}$ (please check). The corresponding eigenvectors are

$$\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ -i\sqrt{2} \end{bmatrix} \quad \text{and} \quad \boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ i\sqrt{2} \end{bmatrix}.$$

Then we calculate that

$$\begin{aligned} \mathbf{x}^{(1)}(t) &= \boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 1 \\ -i\sqrt{2} \end{bmatrix} e^{(-1+i\sqrt{2})t} \\ &= \begin{bmatrix} 1 \\ -i\sqrt{2} \end{bmatrix} e^{-t} (\cos \sqrt{2}t + i \sin \sqrt{2}t) \\ &= e^{-t} \begin{bmatrix} \cos \sqrt{2}t + i \sin \sqrt{2}t \\ -i\sqrt{2} \cos \sqrt{2}t + \sqrt{2} \sin \sqrt{2}t \end{bmatrix} \\ &= e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} + i e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}. \end{aligned}$$

5.4 Complex Eigenvalues



Hence the general solution to the ODE is

$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = c_1 e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} + c_2 e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}.$$

5.4 Complex Eigenvalues



Hence the general solution to the ODE is

$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = c_1 e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} + c_2 e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}.$$

Using the initial condition, we calculate that

$$\begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} I(0) \\ V(0) \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ -\sqrt{2} \end{bmatrix} \quad \Rightarrow \quad \begin{cases} c_1 = 2 \\ c_2 = -\sqrt{2}. \end{cases}$$

5.4 Complex Eigenvalues



Hence the general solution to the ODE is

$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = c_1 e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} + c_2 e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}.$$

Using the initial condition, we calculate that

$$\begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} I(0) \\ V(0) \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ -\sqrt{2} \end{bmatrix} \quad \Rightarrow \quad \begin{cases} c_1 = 2 \\ c_2 = -\sqrt{2}. \end{cases}$$

Thus

$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = 2 e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} - \sqrt{2} e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}.$$

5.4 Complex Eigenvalues



$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = 2e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} - \sqrt{2}e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}$$

So the answers to this problem are

$$I(t) =$$

and

$$V(t) =$$

5.4 Complex Eigenvalues



$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = 2e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} - \sqrt{2}e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}$$

So the answers to this problem are

$$I(t) = 2e^{-t} \cos \sqrt{2}t - \sqrt{2}e^{-t} \sin \sqrt{2}t$$

and

$$V(t) =$$

5.4 Complex Eigenvalues



$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = 2e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} - \sqrt{2}e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}$$


So the answers to this problem are

$$I(t) = 2e^{-t} \cos \sqrt{2}t - \sqrt{2}e^{-t} \sin \sqrt{2}t$$

and

$$V(t) = 2\sqrt{2}e^{-t} \sin \sqrt{2}t + 2e^{-t} \cos \sqrt{2}t.$$

5.4 Complex Eigenvalues



See <https://tinyurl.com/wm2ogdh> for an animated figure.

5.4 Complex Eigenvalues




Example

For the dynamical system shown above, find $x_1(t)$ and $x_2(t)$.


5.4 Complex Eigenvalues



As the springs are stretched and compressed, they apply forces on the blocks as shown below (Hooke's Law).



5.4 Complex Eigenvalues




We calculate that

$$\text{mass} \times \text{acceleration} = \text{force}$$

$$\text{mass} \times \text{acceleration} = \text{force}$$

5.4 Complex Eigenvalues




We calculate that

$$2 \frac{d^2 x_1}{dt^2} = \text{mass} \times \text{acceleration} = \text{force} = -x_1 + 3(x_2 - x_1)$$

$$\text{mass} \times \text{acceleration} = \text{force}$$

5.4 Complex Eigenvalues



We calculate that

$$2 \frac{d^2 x_1}{dt^2} = \text{mass} \times \text{acceleration} = \text{force} = -x_1 + 3(x_2 - x_1)$$

$$\frac{9}{4} \frac{d^2 x_2}{dt^2} = \text{mass} \times \text{acceleration} = \text{force} = -3(x_2 - x_1) - \frac{15}{4}x_2.$$

5.4 Complex Eigenvalues



$$\begin{aligned}2 \frac{d^2 x_1}{dt^2} &= -x_1 + 3(x_2 - x_1) \\ \frac{9}{4} \frac{d^2 x_2}{dt^2} &= -3(x_2 - x_1) - \frac{15}{4} x_2.\end{aligned}$$

This is a system of 2 second order ODEs.

5.4 Complex Eigenvalues



$$\begin{aligned}2 \frac{d^2 x_1}{dt^2} &= -x_1 + 3(x_2 - x_1) \\ \frac{9}{4} \frac{d^2 x_2}{dt^2} &= -3(x_2 - x_1) - \frac{15}{4} x_2.\end{aligned}$$

This is a system of 2 second order ODEs. We want a system of first order ODEs.

5.4 Complex Eigenvalues



$$2 \frac{d^2 x_1}{dt^2} = -x_1 + 3(x_2 - x_1)$$
$$\frac{9}{4} \frac{d^2 x_2}{dt^2} = -3(x_2 - x_1) - \frac{15}{4} x_2.$$

Now let $y_1 = x_1$, $y_2 = x_2$, $y_3 = x_1'$ and $y_4 = x_2'$.

5.4 Complex Eigenvalues



$$2 \frac{d^2 x_1}{dt^2} = -x_1 + 3(x_2 - x_1)$$
$$\frac{9}{4} \frac{d^2 x_2}{dt^2} = -3(x_2 - x_1) - \frac{15}{4} x_2.$$

Now let $y_1 = x_1$, $y_2 = x_2$, $y_3 = x_1'$ and $y_4 = x_2'$. Then

$$y_1' = x_1' = y_3$$

$$y_2' =$$

$$y_3' =$$

$$y_4' =$$

5.4 Complex Eigenvalues



$$2 \frac{d^2 x_1}{dt^2} = -x_1 + 3(x_2 - x_1)$$
$$\frac{9}{4} \frac{d^2 x_2}{dt^2} = -3(x_2 - x_1) - \frac{15}{4} x_2.$$

Now let $y_1 = x_1$, $y_2 = x_2$, $y_3 = x_1'$ and $y_4 = x_2'$. Then

$$y_1' = x_1' = y_3$$

$$y_2' = x_2' = y_4$$

$$y_3' =$$

$$y_4' =$$

5.4 Complex Eigenvalues



$$2 \frac{d^2 x_1}{dt^2} = -x_1 + 3(x_2 - x_1)$$
$$\frac{9}{4} \frac{d^2 x_2}{dt^2} = -3(x_2 - x_1) - \frac{15}{4} x_2.$$

Now let $y_1 = x_1$, $y_2 = x_2$, $y_3 = x_1'$ and $y_4 = x_2'$. Then

$$y_1' = x_1' = y_3$$

$$y_2' = x_2' = y_4$$

$$y_3' = x_1'' = \frac{1}{2} \left(-x_1 + 3x_2 - 3x_1 \right) = -2y_1 + \frac{3}{2}y_2$$

$$y_4' =$$

5.4 Complex Eigenvalues



$$\begin{aligned}2 \frac{d^2 x_1}{dt^2} &= -x_1 + 3(x_2 - x_1) \\ \frac{9}{4} \frac{d^2 x_2}{dt^2} &= -3(x_2 - x_1) - \frac{15}{4} x_2.\end{aligned}$$

Now let $y_1 = x_1$, $y_2 = x_2$, $y_3 = x_1'$ and $y_4 = x_2'$. Then

$$y_1' = x_1' = y_3$$

$$y_2' = x_2' = y_4$$

$$y_3' = x_1'' = \frac{1}{2} \left(-x_1 + 3x_2 - 3x_1 \right) = -2y_1 + \frac{3}{2}y_2$$

$$y_4' = x_2'' = \frac{4}{9} \left(-3x_2 + 3x_1 - \frac{15}{4}x_2 \right) = \frac{4}{3}y_1 - 3y_2.$$

5.4 Complex Eigenvalues



So

$$\mathbf{y}' = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -2 & \frac{3}{2} & 0 & 0 \\ \frac{4}{3} & -3 & 0 & 0 \end{bmatrix} \mathbf{y}.$$

5.4 Complex Eigenvalues



The characteristic polynomial of this matrix is

$$0 = r^4 + 5r^2 + 4 = (r^2 + 1)(r^2 + 4).$$

5.4 Complex Eigenvalues



The characteristic polynomial of this matrix is

$$0 = r^4 + 5r^2 + 4 = (r^2 + 1)(r^2 + 4).$$

So $r_1 = i$, $r_2 = -i$, $r_3 = 2i$ and $r_4 = -2i$.

5.4 Complex Eigenvalues



The characteristic polynomial of this matrix is

$$0 = r^4 + 5r^2 + 4 = (r^2 + 1)(r^2 + 4).$$

So $r_1 = i$, $r_2 = -i$, $r_3 = 2i$ and $r_4 = -2i$. We will use r_1 and r_3 (we do not need r_2 and r_4).

5.4 Complex Eigenvalues



The characteristic polynomial of this matrix is

$$0 = r^4 + 5r^2 + 4 = (r^2 + 1)(r^2 + 4).$$

So $r_1 = i$, $r_2 = -i$, $r_3 = 2i$ and $r_4 = -2i$. We will use r_1 and r_3 (we do not need r_2 and r_4).

The corresponding eigenvectors (please check) are

$$\xi^{(1)} = \begin{bmatrix} 3 \\ 2 \\ 3i \\ 2i \end{bmatrix} \quad \text{and} \quad \xi^{(3)} = \begin{bmatrix} 3 \\ -4 \\ 6i \\ -8i \end{bmatrix}.$$

5.4 Complex Eigenvalues



It follows that

$$\begin{aligned}\boldsymbol{\xi}^{(1)} e^{r_1 t} &= \begin{bmatrix} 3 \\ 2 \\ 3i \\ 2i \end{bmatrix} (\cos t + i \sin t) = \begin{bmatrix} 3 \cos t \\ 2 \cos t \\ -3 \sin t \\ -2 \sin t \end{bmatrix} + i \begin{bmatrix} 3 \sin t \\ 2 \sin t \\ 3 \cos t \\ 2 \cos t \end{bmatrix} \\ &= \mathbf{u}(t) + i \mathbf{v}(t)\end{aligned}$$

and

5.4 Complex Eigenvalues



It follows that

$$\begin{aligned}\boldsymbol{\xi}^{(1)} e^{r_1 t} &= \begin{bmatrix} 3 \\ 2 \\ 3i \\ 2i \end{bmatrix} (\cos t + i \sin t) = \begin{bmatrix} 3 \cos t \\ 2 \cos t \\ -3 \sin t \\ -2 \sin t \end{bmatrix} + i \begin{bmatrix} 3 \sin t \\ 2 \sin t \\ 3 \cos t \\ 2 \cos t \end{bmatrix} \\ &= \mathbf{u}(t) + i\mathbf{v}(t)\end{aligned}$$

and

$$\begin{aligned}\boldsymbol{\xi}^{(3)} e^{r_3 t} &= \begin{bmatrix} 3 \\ -4 \\ 6i \\ -8i \end{bmatrix} (\cos 2t + i \sin 2t) = \begin{bmatrix} 3 \cos 2t \\ -4 \cos 2t \\ -6 \sin 2t \\ +8 \sin 2t \end{bmatrix} + i \begin{bmatrix} 3 \sin 2t \\ -4 \sin 2t \\ 6 \cos 2t \\ -8 \cos 2t \end{bmatrix} \\ &= \mathbf{w}(t) + i\mathbf{z}(t)\end{aligned}$$

5.4 Complex Eigenvalues



Therefore the general solution is

$$\mathbf{y}(t) = c_1 \mathbf{u}(t) + c_2 \mathbf{v}(t) + c_3 \mathbf{w}(t) + c_4 \mathbf{z}(t)$$

5.4 Complex Eigenvalues



Therefore the general solution is

$$\begin{aligned}\mathbf{y}(t) &= c_1 \mathbf{u}(t) + c_2 \mathbf{v}(t) + c_3 \mathbf{w}(t) + c_4 \mathbf{z}(t) \\ &= c_1 \begin{bmatrix} 3 \cos t \\ 2 \cos t \\ -3 \sin t \\ -2 \sin t \end{bmatrix} + c_2 \begin{bmatrix} 3 \sin t \\ 2 \sin t \\ 3 \cos t \\ 2 \cos t \end{bmatrix} + c_3 \begin{bmatrix} 3 \cos 2t \\ -4 \cos 2t \\ -6 \sin 2t \\ 8 \sin 2t \end{bmatrix} + c_4 \begin{bmatrix} 3 \sin 2t \\ -4 \sin 2t \\ 6 \cos 2t \\ -8 \cos 2t \end{bmatrix}.\end{aligned}$$

5.4 Complex Eigenvalues



Example

Suppose that the above system has initial condition

$$\mathbf{y}(0) = \begin{bmatrix} -1 \\ 4 \\ 1 \\ 1 \end{bmatrix}.$$

Sketch graphs of $y_1(t)$ and $y_2(t)$.

5.4 Complex Eigenvalues



The initial value problem


$$\mathbf{y}' = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -2 & \frac{3}{2} & 0 & 0 \\ \frac{4}{3} & -3 & 0 & 0 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} -1 \\ 4 \\ 1 \\ 1 \end{bmatrix}$$

has solution


$$\mathbf{y}(t) = \frac{4}{9} \begin{bmatrix} 3 \cos t \\ 2 \cos t \\ -3 \sin t \\ -2 \sin t \end{bmatrix} + \frac{7}{18} \begin{bmatrix} 3 \sin t \\ 2 \sin t \\ 3 \cos t \\ 2 \cos t \end{bmatrix} - \frac{7}{9} \begin{bmatrix} 3 \cos 2t \\ -4 \cos 2t \\ -6 \sin 2t \\ 8 \sin 2t \end{bmatrix} - \frac{1}{36} \begin{bmatrix} 3 \sin 2t \\ -4 \sin 2t \\ 6 \cos 2t \\ -8 \cos 2t \end{bmatrix}.$$

Then we can draw the graphs of y_1 and y_2 :

5.4 Complex Eigenvalues



5.4 Complex Eigenvalues



5.4 Complex Eigenvalues



Please see <https://tinyurl.com/s7uww7m>

Fundamental Matrices

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = P(t)\mathbf{x}$$

where $P(t)$ is an $n \times n$ matrix function.



Now consider

$$\mathbf{x}' = P(t)\mathbf{x}$$

where $P(t)$ is an $n \times n$ matrix function.

Suppose that $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(n)}$ are linearly independent solutions to this ODE. In other words, suppose that $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(n)}$ form a *fundamental set of solutions* to this ODE.

Definition

The matrix

$$\Psi(t) = \begin{bmatrix} \mathbf{x}^{(1)} & \mathbf{x}^{(2)} & \dots & \mathbf{x}^{(n)} \end{bmatrix} = \begin{bmatrix} x_1^{(1)}(t) & x_1^{(2)}(t) & \dots & x_1^{(n)}(t) \\ x_2^{(1)}(t) & x_2^{(2)}(t) & \dots & x_2^{(n)}(t) \\ \vdots & \vdots & & \vdots \\ x_n^{(1)}(t) & x_n^{(2)}(t) & \dots & x_n^{(n)}(t) \end{bmatrix}$$

is called a *fundamental matrix* of $\mathbf{x}' = P(t)\mathbf{x}$.

5.5 Fundamental Matrices



Example

Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Example

Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Recall that

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}$$

form a fundamental set of solutions to this ODE.

5.5 Fundamental Matrices



Example

Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Recall that

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}$$

form a fundamental set of solutions to this ODE. Therefore

$$\Psi(t) = \begin{bmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{bmatrix}$$

is a fundamental matrix of this ODE.

5.5 Fundamental Matrices



Now, the general solution to

$$\mathbf{x}' = A\mathbf{x}$$

is

$$\mathbf{x}(t) = c_1\mathbf{x}^{(1)}(t) + c_2\mathbf{x}^{(2)}(t) + \dots + c_n\mathbf{x}^{(n)}(t)$$

5.5 Fundamental Matrices



Now, the general solution to

$$\mathbf{x}' = A\mathbf{x}$$

is

$$\mathbf{x}(t) = c_1\mathbf{x}^{(1)}(t) + c_2\mathbf{x}^{(2)}(t) + \dots + c_n\mathbf{x}^{(n)}(t) = \Psi(t)\mathbf{c}$$

where

$$\mathbf{c} = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \in \mathbb{R}^n.$$

5.5 Fundamental Matrices



Now, the general solution to

$$\mathbf{x}' = A\mathbf{x}$$

is

$$\mathbf{x}(t) = c_1\mathbf{x}^{(1)}(t) + c_2\mathbf{x}^{(2)}(t) + \dots + c_n\mathbf{x}^{(n)}(t) = \Psi(t)\mathbf{c}$$

where

$$\mathbf{c} = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \in \mathbb{R}^n.$$

If we have an initial condition $\mathbf{x}(t_0) = \mathbf{x}^0$, then

$$\Psi(t_0)\mathbf{c} = \mathbf{x}^0.$$

5.5 Fundamental Matrices



$$\mathbf{x}(t) = \Psi(t)\mathbf{c}$$

$$\Psi(t_0)\mathbf{c} = \mathbf{x}^0$$

But

$$\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)}$$

are linearly
independent



$$\mathbf{x}(t) = \Psi(t)\mathbf{c}$$

$$\Psi(t_0)\mathbf{c} = \mathbf{x}^0$$

But

$$\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)}$$

are linearly
independent

\implies

$\Psi(t)$ is invertible

5.5 Fundamental Matrices



$$\mathbf{x}(t) = \Psi(t)\mathbf{c}$$

$$\Psi(t_0)\mathbf{c} = \mathbf{x}^0$$

But

$$\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)}$$

are linearly
independent

$$\implies \Psi(t) \text{ is invertible}$$

$$\implies \mathbf{c} = \Psi^{-1}(t_0)\mathbf{x}^0.$$

5.5 Fundamental Matrices



$$\mathbf{x}(t) = \Psi(t)\mathbf{c} \qquad \Psi(t_0)\mathbf{c} = \mathbf{x}^0$$

But

$$\begin{aligned} \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)} \\ \text{are linearly} & \implies \Psi(t) \text{ is invertible} \\ \text{independent} & \\ & \implies \mathbf{c} = \Psi^{-1}(t_0)\mathbf{x}^0. \end{aligned}$$

Therefore the solution to the IVP

$$\begin{cases} \mathbf{x}' = A\mathbf{x} \\ \mathbf{x}(t_0) = \mathbf{x}^0 \end{cases}$$

is

$$\boxed{\mathbf{x}(t) = \Psi(t)\Psi^{-1}(t_0)\mathbf{x}^0.}$$



Theorem

Suppose that $\Psi(t)$ is a fundamental matrix for $\mathbf{x}' = P(t)\mathbf{x}$. Then $\Psi(t)$ solves the differential equation $\Psi' = P(t)\Psi$.

(You prove)

Remark

It is possible to find a *special fundamental matrix*, $\Phi(t)$, which satisfies

$$\Phi(t_0) = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} = I.$$

Remark

It is possible to find a *special fundamental matrix*, $\Phi(t)$, which satisfies

$$\Phi(t_0) = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} = I.$$

We will use Φ for this special fundamental matrix, and Ψ for any fundamental matrix.

5.5 Fundamental Matrices



Example

Consider

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Find the special fundamental matrix which satisfies $\Phi(0) = I$.

5.5 Fundamental Matrices



To find the matrix Φ which satisfies

$$\Phi(0) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

we must solve two IVPs:

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \end{cases}$$

and

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \end{cases}$$



The general solution to the ODE is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}.$$

5.5 Fundamental Matrices



We calculate that

$$\begin{aligned} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \mathbf{x}(0) &= c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} \implies \begin{aligned} c_1 &= \frac{1}{2} \\ c_2 &= \frac{1}{2} \end{aligned} \\ &\implies \mathbf{x}(t) = \begin{bmatrix} \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} \\ e^{3t} - e^{-t} \end{bmatrix} \end{aligned}$$

5.5 Fundamental Matrices



We calculate that

$$\begin{aligned}\begin{bmatrix} 1 \\ 0 \end{bmatrix} = \mathbf{x}(0) &= c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} \implies \begin{aligned} c_1 &= \frac{1}{2} \\ c_2 &= \frac{1}{2} \end{aligned} \\ \implies \mathbf{x}(t) &= \begin{bmatrix} \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} \\ e^{3t} - e^{-t} \end{bmatrix}\end{aligned}$$

and

$$\begin{aligned}\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \mathbf{x}(0) &= c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} \implies \begin{aligned} c_1 &= \frac{1}{4} \\ c_2 &= -\frac{1}{4} \end{aligned} \\ \implies \mathbf{x}(t) &= \begin{bmatrix} \frac{1}{4}e^{3t} - \frac{1}{4}e^{-t} \\ \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} \end{bmatrix}.\end{aligned}$$

5.5 Fundamental Matrices



Therefore the special fundamental matrix is

$$\Phi(t) = \begin{bmatrix} \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} & \frac{1}{4}e^{3t} - \frac{1}{4}e^{-t} \\ e^{3t} - e^{-t} & \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} \end{bmatrix}.$$

What is e^{At} ?

Recall that the solution to

$$\begin{cases} x' = ax & (a \in \mathbb{R}) \\ x(0) = x_0 \end{cases}$$

is

$$x(t) = x_0 e^{at} = x_0 \exp(at)$$

and recall that

$$\exp(at) = 1 + \sum_{n=1}^{\infty} \frac{a^n t^n}{n!}.$$

5.5 Fundamental Matrices



Now consider

$$\begin{cases} \mathbf{x}' = A\mathbf{x} \\ \mathbf{x}(0) = \mathbf{x}^0 \end{cases}$$

for $A \in \mathbb{R}^{n \times n}$.

Now consider

$$\begin{cases} \mathbf{x}' = A\mathbf{x} \\ \mathbf{x}(0) = \mathbf{x}^0 \end{cases}$$

for $A \in \mathbb{R}^{n \times n}$.

Definition

$$\exp(At) = I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} = I + At + \frac{A^2 t^2}{2!} + \frac{A^3 t^3}{3!} + \dots$$

5.5 Fundamental Matrices



Note that

$$\begin{aligned}\frac{d}{dt} \exp(At) &= \exp(At) A \\ &= A \exp(At) \\ &= \exp(At) A \\ &= A \exp(At) \end{aligned}$$

5.5 Fundamental Matrices



Note that

$$\frac{d}{dt} \exp(At) = \frac{d}{dt} \left(I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} \right) =$$

$$=$$

$$=$$

$$= \exp(At) A = A \exp(At).$$

5.5 Fundamental Matrices



Note that

$$\begin{aligned}\frac{d}{dt} \exp(At) &= \frac{d}{dt} \left(I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} \right) = \mathbf{0} + \sum_{n=1}^{\infty} \frac{d}{dt} \left(\frac{A^n t^n}{n!} \right) \\ &= \\ &= \\ &= \end{aligned}$$

5.5 Fundamental Matrices



Note that

$$\begin{aligned}\frac{d}{dt} \exp(At) &= \frac{d}{dt} \left(I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} \right) = \mathbf{0} + \sum_{n=1}^{\infty} \frac{d}{dt} \left(\frac{A^n t^n}{n!} \right) \\ &= \sum_{n=1}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} = \\ &= \\ &= \end{aligned}$$

5.5 Fundamental Matrices



Note that

$$\begin{aligned}\frac{d}{dt} \exp(At) &= \frac{d}{dt} \left(I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} \right) = \mathbf{0} + \sum_{n=1}^{\infty} \frac{d}{dt} \left(\frac{A^n t^n}{n!} \right) \\ &= \sum_{n=1}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} = A + \sum_{n=2}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} \\ &= \\ &= \end{aligned}$$

5.5 Fundamental Matrices



Note that

$$\begin{aligned}\frac{d}{dt} \exp(At) &= \frac{d}{dt} \left(I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} \right) = \mathbf{0} + \sum_{n=1}^{\infty} \frac{d}{dt} \left(\frac{A^n t^n}{n!} \right) \\ &= \sum_{n=1}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} = A + \sum_{n=2}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} \\ &= A + \sum_{k=1}^{\infty} \frac{A^{k+1} t^k}{k!} \quad (k = n-1) \\ &= \quad \quad \quad = \quad \quad \quad .\end{aligned}$$

5.5 Fundamental Matrices



Note that

$$\begin{aligned}\frac{d}{dt} \exp(At) &= \frac{d}{dt} \left(I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} \right) = \mathbf{0} + \sum_{n=1}^{\infty} \frac{d}{dt} \left(\frac{A^n t^n}{n!} \right) \\&= \sum_{n=1}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} = A + \sum_{n=2}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} \\&= A + \sum_{k=1}^{\infty} \frac{A^{k+1} t^k}{k!} \quad (k = n - 1) \\&= A \left(I + \sum_{k=1}^{\infty} \frac{A^k t^k}{k!} \right) = \end{aligned}$$

Note that

$$\begin{aligned}\frac{d}{dt} \exp(At) &= \frac{d}{dt} \left(I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} \right) = \mathbf{0} + \sum_{n=1}^{\infty} \frac{d}{dt} \left(\frac{A^n t^n}{n!} \right) \\&= \sum_{n=1}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} = A + \sum_{n=2}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} \\&= A + \sum_{k=1}^{\infty} \frac{A^{k+1} t^k}{k!} \quad (k = n - 1) \\&= A \left(I + \sum_{k=1}^{\infty} \frac{A^k t^k}{k!} \right) = A \exp(At).\end{aligned}$$



This means that $\exp(At)$ solves

$$\begin{cases} (\exp(At))' = A \exp(At) \\ \exp(At)|_{t=0} = I. \end{cases}$$



This means that $\exp(At)$ solves

$$\begin{cases} (\exp(At))' = A \exp(At) \\ \exp(At)|_{t=0} = I. \end{cases}$$

But remember that Φ solves

$$\begin{cases} \Phi' = A\Phi \\ \Phi(0) = I. \end{cases}$$

This means that $\exp(At)$ solves

$$\begin{cases} (\exp(At))' = A \exp(At) \\ \exp(At)|_{t=0} = I. \end{cases}$$

But remember that Φ solves

$$\begin{cases} \Phi' = A\Phi \\ \Phi(0) = I. \end{cases}$$

Therefore

$$\boxed{\Phi(t) = \exp(At).}$$

5.5 Fundamental Matrices



Example

Let $A = \begin{bmatrix} 8 & -1 \\ 6 & 1 \end{bmatrix}$. Find $\exp(At)$.

5.5 Fundamental Matrices



Example

Let $A = \begin{bmatrix} 8 & -1 \\ 6 & 1 \end{bmatrix}$. Find $\exp(At)$.

We have previously found that the general solution to $\mathbf{x}' = A\mathbf{x}$ is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

5.5 Fundamental Matrices



To satisfy $\mathbf{x}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ we require $c_1 = \frac{6}{5}$ and $c_2 = -\frac{1}{5}$. Hence

$$\mathbf{x}(t) = \frac{6}{5} \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} - \frac{1}{5} \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t} = \begin{bmatrix} \frac{6}{5}e^{7t} - \frac{1}{5}e^{2t} \\ \frac{6}{5}e^{7t} - \frac{6}{5}e^{2t} \end{bmatrix}.$$

5.5 Fundamental Matrices



To satisfy $\mathbf{x}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ we require $c_1 = \frac{6}{5}$ and $c_2 = -\frac{1}{5}$. Hence

$$\mathbf{x}(t) = \frac{6}{5} \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} - \frac{1}{5} \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t} = \begin{bmatrix} \frac{6}{5}e^{7t} - \frac{1}{5}e^{2t} \\ \frac{6}{5}e^{7t} - \frac{6}{5}e^{2t} \end{bmatrix}.$$

To satisfy $\mathbf{x}(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ we require $c_1 = -\frac{1}{5}$ and $c_2 = \frac{1}{5}$. Hence

$$\mathbf{x}(t) = -\frac{1}{5} \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + \frac{1}{5} \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t} = \begin{bmatrix} -\frac{1}{5}e^{7t} + \frac{1}{5}e^{2t} \\ -\frac{1}{5}e^{7t} + \frac{6}{5}e^{2t} \end{bmatrix}.$$

5.5 Fundamental Matrices



Therefore the answer is

$$\exp(At) = \Phi(t) = \begin{bmatrix} \frac{6}{5}e^{7t} - \frac{1}{5}e^{2t} & -\frac{1}{5}e^{7t} + \frac{1}{5}e^{2t} \\ \frac{6}{5}e^{7t} - \frac{6}{5}e^{2t} & -\frac{1}{5}e^{7t} + \frac{6}{5}e^{2t} \end{bmatrix}.$$

Diagonalisable Matrices

If

$$D = \begin{bmatrix} r_1 & 0 & 0 & \dots & 0 \\ 0 & r_2 & 0 & \dots & 0 \\ 0 & 0 & r_3 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & r_n \end{bmatrix}$$

is a diagonal matrix, then it is easy to calculate $\exp(Dt)$. We simply have

$$\exp(Dt) = \begin{bmatrix} e^{r_1 t} & 0 & 0 & \dots & 0 \\ 0 & e^{r_2 t} & 0 & \dots & 0 \\ 0 & 0 & e^{r_3 t} & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & e^{r_n t} \end{bmatrix}.$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}$$

for $A \in \mathbb{R}^{n \times n}$.

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}$$

for $A \in \mathbb{R}^{n \times n}$. Recall how we diagonalise a matrix: If $\boldsymbol{\xi}^{(1)}, \boldsymbol{\xi}^{(2)}, \dots, \boldsymbol{\xi}^{(n)}$ are the eigenvectors of A , we let

$$T = \begin{bmatrix} \boldsymbol{\xi}^{(1)} & \boldsymbol{\xi}^{(2)} & \dots & \boldsymbol{\xi}^{(n)} \end{bmatrix}.$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}$$

for $A \in \mathbb{R}^{n \times n}$. Recall how we diagonalise a matrix: If $\boldsymbol{\xi}^{(1)}, \boldsymbol{\xi}^{(2)}, \dots, \boldsymbol{\xi}^{(n)}$ are the eigenvectors of A , we let

$$T = \begin{bmatrix} \boldsymbol{\xi}^{(1)} & \boldsymbol{\xi}^{(2)} & \dots & \boldsymbol{\xi}^{(n)} \end{bmatrix}.$$

Then

$$\det(T) \neq 0 \implies \begin{matrix} T^{-1} \\ \text{exists} \end{matrix}$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}$$

for $A \in \mathbb{R}^{n \times n}$. Recall how we diagonalise a matrix: If $\boldsymbol{\xi}^{(1)}, \boldsymbol{\xi}^{(2)}, \dots, \boldsymbol{\xi}^{(n)}$ are the eigenvectors of A , we let

$$T = \begin{bmatrix} \boldsymbol{\xi}^{(1)} & \boldsymbol{\xi}^{(2)} & \dots & \boldsymbol{\xi}^{(n)} \end{bmatrix}.$$

Then

$$\det(T) \neq 0 \implies \begin{matrix} T^{-1} \\ \text{exists} \end{matrix} \implies \begin{matrix} T^{-1}AT \\ \text{is diagonal} \end{matrix}$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}$$

for $A \in \mathbb{R}^{n \times n}$. Recall how we diagonalise a matrix: If $\boldsymbol{\xi}^{(1)}, \boldsymbol{\xi}^{(2)}, \dots, \boldsymbol{\xi}^{(n)}$ are the eigenvectors of A , we let

$$T = \begin{bmatrix} \boldsymbol{\xi}^{(1)} & \boldsymbol{\xi}^{(2)} & \dots & \boldsymbol{\xi}^{(n)} \end{bmatrix}.$$

Then

$$\det(T) \neq 0 \implies \begin{matrix} T^{-1} \\ \text{exists} \end{matrix} \implies \begin{matrix} T^{-1}AT \\ \text{is diagonal} \end{matrix} \implies A \text{ is diagonalisable.}$$

5.5 Fundamental Matrices



Example

Diagonalise

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}.$$

Example

Diagonalise

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}.$$

The eigenvalues are $r_1 = 3$ and $r_2 = -1$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$.

Example

Diagonalise

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}.$$

The eigenvalues are $r_1 = 3$ and $r_2 = -1$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$. Thus

$$T = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix}$$

Example

Diagonalise

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}.$$

The eigenvalues are $r_1 = 3$ and $r_2 = -1$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$. Thus

$$T = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix} \quad \text{and} \quad T^{-1} = \begin{bmatrix} \frac{1}{2} & \frac{1}{4} \\ \frac{1}{2} & -\frac{1}{4} \end{bmatrix}.$$

Example

Diagonalise

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}.$$

The eigenvalues are $r_1 = 3$ and $r_2 = -1$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$. Thus

$$T = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix} \quad \text{and} \quad T^{-1} = \begin{bmatrix} \frac{1}{2} & \frac{1}{4} \\ \frac{1}{2} & -\frac{1}{4} \end{bmatrix}.$$

It follows that

$$D = T^{-1}AT = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix}.$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}.$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}.$$

Define a new variable \mathbf{y} by

$$\mathbf{x} = T\mathbf{y} \quad \text{or} \quad \mathbf{y} = T^{-1}\mathbf{x}.$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}.$$

Define a new variable \mathbf{y} by

$$\mathbf{x} = T\mathbf{y} \quad \text{or} \quad \mathbf{y} = T^{-1}\mathbf{x}.$$

Then we calculate that

$$\mathbf{x}' = A\mathbf{x}$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}.$$

Define a new variable \mathbf{y} by

$$\mathbf{x} = T\mathbf{y} \quad \text{or} \quad \mathbf{y} = T^{-1}\mathbf{x}.$$

Then we calculate that

$$\begin{aligned}\mathbf{x}' &= A\mathbf{x} \\ T\mathbf{y}' &= AT\mathbf{y}\end{aligned}$$

5.5 Fundamental Matrices



Now consider

$$\mathbf{x}' = A\mathbf{x}.$$

Define a new variable \mathbf{y} by

$$\mathbf{x} = T\mathbf{y} \quad \text{or} \quad \mathbf{y} = T^{-1}\mathbf{x}.$$

Then we calculate that

$$\begin{aligned}\mathbf{x}' &= A\mathbf{x} \\ T\mathbf{y}' &= AT\mathbf{y} \\ \mathbf{y}' &= T^{-1}AT\mathbf{y} = D\mathbf{y}.\end{aligned}$$

5.5 Fundamental Matrices



We know that a fundamental matrix for $\mathbf{y}' = D\mathbf{y}$ is

$$\exp(Dt) = \begin{bmatrix} e^{r_1 t} & 0 & \dots & 0 \\ 0 & e^{r_2 t} & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & e^{r_n t} \end{bmatrix}.$$

5.5 Fundamental Matrices



We know that a fundamental matrix for $\mathbf{y}' = D\mathbf{y}$ is

$$\exp(Dt) = \begin{bmatrix} e^{r_1 t} & 0 & \dots & 0 \\ 0 & e^{r_2 t} & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & e^{r_n t} \end{bmatrix}.$$

Therefore a fundamental matrix for $\mathbf{x}' = A\mathbf{x}$ is

$$\Psi = T \exp(Dt) = \begin{bmatrix} \boldsymbol{\xi}^{(1)} e^{r_1 t} & \boldsymbol{\xi}^{(2)} e^{r_2 t} & \dots & \boldsymbol{\xi}^{(n)} e^{r_n t} \end{bmatrix}.$$



Example

Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Example

Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Recall that $T = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix}$.

Example

Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Recall that $T = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix}$. Letting $\mathbf{y} = T^{-1}\mathbf{x}$, we have

$$\mathbf{y}' = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{y}.$$

5.5 Fundamental Matrices



A fundamental matrix for $\mathbf{y}' = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{y}$ is

$$\exp(Dt) = e^{Dt} = \begin{bmatrix} e^{3t} & 0 \\ 0 & e^{-t} \end{bmatrix}.$$

5.5 Fundamental Matrices



A fundamental matrix for $\mathbf{y}' = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{y}$ is

$$\exp(Dt) = e^{Dt} = \begin{bmatrix} e^{3t} & 0 \\ 0 & e^{-t} \end{bmatrix}.$$

Hence a fundamental matrix for $\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}$ is

$$\Psi(t) = T \exp(Dt)$$

A fundamental matrix for $\mathbf{y}' = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{y}$ is

$$\exp(Dt) = e^{Dt} = \begin{bmatrix} e^{3t} & 0 \\ 0 & e^{-t} \end{bmatrix}.$$

Hence a fundamental matrix for $\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}$ is

$$\Psi(t) = T \exp(Dt) = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} e^{3t} & 0 \\ 0 & e^{-t} \end{bmatrix}$$

5.5 Fundamental Matrices



A fundamental matrix for $\mathbf{y}' = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{y}$ is

$$\exp(Dt) = e^{Dt} = \begin{bmatrix} e^{3t} & 0 \\ 0 & e^{-t} \end{bmatrix}.$$

Hence a fundamental matrix for $\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}$ is

$$\Psi(t) = T \exp(Dt) = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} e^{3t} & 0 \\ 0 & e^{-t} \end{bmatrix} = \begin{bmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{bmatrix}.$$

Next Time

- 5.6 Repeated Eigenvalues
- 5.7 Nonhomogeneous Linear Systems