

Differential Equations

Chapter 6 and Chapter 7

Neil Course

2019-20

Contents

	Page
6 The Laplace Transform	1
6.1 Definition of the Laplace Transform	1
6.2 Solving Initial Value Problems	4
6.3 Step Functions	8
6.4 ODEs with Discontinuous Forcing Functions	13
6.6 The Convolution Integral	17
7 Systems of First Order Linear Equations	19
7.1 Introduction	19
7.4 Basic Theory of Systems of First Order Linear Equations	22
7.5 Homogeneous Linear Systems with Constant Coefficients	24
7.6 Complex Eigenvalues	27
7.7 Fundamental Matrices	34
7.8 Repeated Eigenvalues	39
7.9 Nonhomogeneous Linear Systems	44

6

The Laplace Transform

Recall that $\int_a^\infty f(t) dt$ means $\lim_{R \rightarrow \infty} \int_a^R f(t) dt$.

Example 6.1. Let $c \neq 0$. Then

$$\int_0^\infty e^{ct} dt = \lim_{R \rightarrow \infty} \int_0^R e^{ct} dt = \lim_{R \rightarrow \infty} \left[\frac{1}{c} e^{ct} \right]_0^R = \lim_{R \rightarrow \infty} \frac{1}{c} (e^{cR} - 1) = \begin{cases} \infty & c > 0 \\ -\frac{1}{c} & c < 0. \end{cases}$$

Example 6.2.

$$\int_1^\infty \frac{1}{t} dt = \lim_{R \rightarrow \infty} \int_1^R \frac{1}{t} dt = \lim_{R \rightarrow \infty} [\ln t]_1^R = \lim_{R \rightarrow \infty} (\ln R - 0) = \infty$$

6.1 Definition of the Laplace Transform

$\frac{d}{dt}$ changes a function $f(t)$ into a new function $f'(t)$.
 \mathcal{L} changes a function $f(t)$ into a new function $F(s)$.

Definition. Suppose that

- (i) $A > 0$, $K > 0$, $M > 0$, $a \in \mathbb{R}$;
- (ii) f is piecewise continuous on $[0, A]$; and
- (iii) $|f(t)| \leq Ke^{at}$ for all $t \geq M$.

The **Laplace Transform** of $f : [0, \infty) \rightarrow \mathbb{R}$ is a new function defined by

$$F(s) = \mathcal{L}[f](s) = \int_0^\infty e^{-st} f(t) dt.$$

Example 6.3.

$$\mathcal{L}[1](s) = \int_0^\infty e^{-st} dt = - \lim_{R \rightarrow \infty} \left[\frac{e^{-st}}{s} \right]_0^R = \frac{1}{s} \quad \text{if } s > 0.$$



Pierre-Simon
Laplace
FRA, 1749-1827

Example 6.4.

$$\mathcal{L}[e^{at}](s) = \int_0^\infty e^{-st} e^{at} dt = \int_0^\infty e^{-(s-a)t} dt = \frac{1}{s-a} \quad \text{if } s > a.$$

The Laplace Transform of $e^{at} : [0, \infty) \rightarrow \mathbb{R}$ is $\frac{1}{s-a} : (a, \infty) \rightarrow \mathbb{R}$.

Example 6.5. Let

$$f(t) = \begin{cases} 1 & 0 \leq t < 1 \\ k & t = 1 \\ 0 & t > 1. \end{cases}$$

Then

$$F(s) = \mathcal{L}[f](s) = \int_0^\infty e^{-st} f(t) dt = \int_0^1 e^{-st} dt = \left[-\frac{1}{s} e^{-st} \right]_0^1 = \frac{1 - e^{-s}}{s} \quad \text{if } s > 0.$$

Example 6.6. Now consider $g(t) = \sin at$ ($t \geq 0$). Using integration by parts, we have

$$\begin{aligned} G(s) = \mathcal{L}[g](s) &= \int_0^\infty e^{-st} \sin at dt = \lim_{R \rightarrow \infty} \int_0^R e^{-st} \sin at dt \\ &= \lim_{R \rightarrow \infty} \left(\left[-\frac{1}{a} e^{-st} \cos at \right]_0^R - \frac{s}{a} \int_0^R e^{-st} \cos at dt \right) = \frac{1}{a} - \frac{s}{a} \int_0^\infty e^{-st} \cos at dt. \end{aligned}$$

Using integration by parts a second time, we have

$$G(s) = \frac{1}{a} - \frac{s^2}{a^2} \int_0^\infty e^{-st} \sin at dt = \frac{1}{a} - \frac{s^2}{a^2} G(s).$$

Therefore

$$\mathcal{L}[\sin at](s) = G(s) = \frac{a}{s^2 + a^2} \quad \text{if } s > 0.$$

Theorem 6.1.

$$\mathcal{L}[c_1 f_1 + c_2 f_2] = c_1 \mathcal{L}[f_1] + c_2 \mathcal{L}[f_2].$$

You prove.

Example 6.7. If $h(t) = 5e^{-2t} - 3 \sin 4t$ ($t \geq 0$), then

$$H(s) = \mathcal{L}[h](s) = 5\mathcal{L}[e^{-2t}] - 3\mathcal{L}[\sin 4t] = \frac{5}{s+2} - \frac{12}{s^2 + 16} \quad \text{if } s > 0.$$

Theorem 6.2.

$$\mathcal{L}[t^n f(t)] = (-1)^n \frac{d^n F}{ds^n}$$

You prove this. In Exercise 28(f), you are required to prove this formula with $n = 1$.

Example 6.8.

$$\mathcal{L}[t^2 \cosh 2t] = (-1)^2 \frac{d^2}{ds^2} \mathcal{L}[\cosh 2t] = \frac{d^2}{ds^2} \left(\frac{s}{s^2 - 2^2} \right) = \dots = \frac{2s(s^2 + 12)}{(s^2 - 4)^3}$$

$f(t)$	$F(s) = \mathcal{L}[f](s)$	
1	$\frac{1}{s}$	$s > 0$
e^{at}	$\frac{1}{s-a}$	$s > a$
$t^n \quad (n \in \mathbb{N})$	$\frac{n!}{s^{n+1}}$	$s > 0$
$\sin at$	$\frac{a}{s^2+a^2}$	$s > 0$
$\cos at$	$\frac{s}{s^2+a^2}$	$s > 0$
$\sinh at$	$\frac{a}{s^2-a^2}$	$s > a $
$\cosh at$	$\frac{s}{s^2-a^2}$	$s > a $
$e^{at} \sin bt$	$\frac{b}{(s-a)^2+b^2}$	$s > a$
$e^{at} \cos bt$	$\frac{s-a}{(s-a)^2+b^2}$	$s > a$
$t^n e^{at} \quad (n \in \mathbb{N})$	$\frac{n!}{(s-a)^{n+1}}$	$s > a$
$u_c(t)$	$\frac{e^{-cs}}{s}$	$s > 0$
$u_c(t)f(t-c)$	$e^{-cs}F(s)$	
$e^{ct}f(t)$	$F(s-c)$	
$f(ct) \quad (c > 0)$	$\frac{1}{c}F\left(\frac{s}{c}\right)$	
$\int_0^t f(t-\tau)g(\tau)d\tau$	$F(s)G(s)$	
$t^n f(t)$	$(-1)^n F^{(n)}(s)$	
$\mathcal{L}[f'](s) = s\mathcal{L}[f](s) - f(0)$ $\mathcal{L}[f''](s) = s^2\mathcal{L}[f](s) - sf(0) - f'(0)$ $\mathcal{L}[f^{(n)}](s) = s^n\mathcal{L}[f](s) - s^{n-1}f(0) - \dots - sf^{(n-2)}(0) - f^{(n-1)}(0)$		

Inverse Laplace Transforms

$$\mathcal{L}[f] = F \quad \Longleftrightarrow \quad \mathcal{L}^{-1}[F] = f.$$

Example 6.9. Find the inverse Laplace Transform of $F(s) = \frac{9s^2 - 12s + 216}{s(s^2 + 36)}$.

Using partial fractions we calculate that

$$\begin{aligned} A + B &= 9 \\ C &= -12 \\ 36A &= 216 \end{aligned}$$

$$\begin{aligned} A &= 6 \\ B &= 3 \\ C &= -12 \end{aligned}$$

$$\begin{aligned} F(s) &= \frac{9s^2 - 12s + 216}{s(s^2 + 36)} = \frac{A}{s} + \frac{Bs + C}{s^2 + 36} = \frac{As^2 + 36A + Bs + C}{s(s^2 + 36)} \\ &= 6 \left(\frac{1}{s} \right) + 3 \left(\frac{s}{s^2 + 36} \right) - 12 \left(\frac{1}{s^2 + 36} \right) \\ &= 6 \left(\frac{1}{s} \right) + 3 \left(\frac{s}{s^2 + 36} \right) - \frac{12}{6} \left(\frac{6}{s^2 + 36} \right) \\ &= 6\mathcal{L}[1] + 3\mathcal{L}[\cos 6t] - 2\mathcal{L}[\sin 6t]. \end{aligned}$$

and that

$$f(t) = \mathcal{L}^{-1}[F](t) = 6 + 3 \cos 6t - 2 \sin 6t.$$

6.2 Solving Initial Value Problems

Theorem 6.3.

- (i). $\mathcal{L}[f'](s) = s\mathcal{L}[f](s) - f(0)$.
- (ii). $\mathcal{L}[f''](s) = s^2\mathcal{L}[f](s) - sf(0) - f'(0)$.
- (iii). $\mathcal{L}[f'''](s) = s^3\mathcal{L}[f](s) - s^2f(0) - sf'(0) - f''(0)$.
- (iv). $\mathcal{L}[f^{(n)}](s) = s^n\mathcal{L}[f](s) - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - sf^{(n-2)}(0) - f^{(n-1)}(0)$.

Proof.

(i). Using integration-by-parts ($\int uv' = uv - \int u'v$) we calculate that

$$\begin{aligned} \mathcal{L}[f'](s) &= \int_0^\infty e^{-st} f'(t) dt = [e^{-st} f(t)]_0^\infty - \int_0^\infty \left(\frac{d}{dt} e^{-st} \right) f(t) dt \\ &= 0 - f(0) - \int_0^\infty -se^{-st} f(t) dt = -f(0) + s \int_0^\infty e^{-st} f(t) dt \\ &= -f(0) + sF(s) \end{aligned}$$

as required.

(ii). Using (i), but replacing each f by f' we get

$$\mathcal{L}[f''](s) = s\mathcal{L}[f'](s) - f'(0) = s(s\mathcal{L}[f](s) - f(0)) - f'(0) = s^2\mathcal{L}[f](s) - sf(0) - f'(0).$$

You prove (iii) and (iv)

□

Example 6.10. Solve

$$\begin{cases} y'' - y' - 2y = 0 \\ y(0) = 1 \\ y'(0) = 0. \end{cases}$$


solution 1 (method from Chapter 3): The characteristic equation

$$0 = r^2 - r - 2 = (r - 2)(r + 1)$$

has roots $r_1 = -1$ and $r_2 = 2$. So $y = c_1 e^{-t} + c_2 e^{2t}$. Using the initial conditions we find that $c_1 = \frac{2}{3}$ and $c_2 = \frac{1}{3}$. Therefore

$$y(t) = \frac{2}{3}e^{-t} + \frac{1}{3}e^{2t}.$$

solution 2 (Laplace Transform):



First we take the Laplace Transform of the ODE

$$\begin{aligned} y'' - y' - 2y &= 0 \\ \mathcal{L}[y'' - y' - 2y] &= \mathcal{L}[0] \\ \mathcal{L}[y''] - \mathcal{L}[y'] - 2\mathcal{L}[y] &= 0 \\ (s^2 Y - sy(0) - y'(0)) - (sY - y(0)) - 2Y &= 0 \\ (s^2 Y - s - 0) - (sY - 1) - 2Y &= 0 \\ (s^2 - s - 2)Y + (1 - s) &= 0 \end{aligned}$$

Thus

$$Y(s) = \frac{s - 1}{s^2 - s - 2} = \frac{s - 1}{(s - 2)(s + 1)}.$$

Using partial fractions we obtain

$$\begin{aligned} Y(s) &= \frac{s - 1}{(s - 2)(s + 1)} = \frac{A}{s - 2} + \frac{B}{s + 1} = \frac{As + A + Bs - 2B}{(s - 2)(s + 1)} \\ &= \frac{1}{3} \left(\frac{1}{s - 2} \right) + \frac{2}{3} \left(\frac{1}{s + 1} \right). \end{aligned}$$

$$\begin{aligned} A + B &= 1 \\ A - 2B &= -1 \end{aligned}$$

$$\begin{aligned} A &= \frac{1}{3} \\ B &= \frac{2}{3} \end{aligned}$$

But recall that $\mathcal{L}[e^{2t}] = \frac{1}{s-2}$ and $\mathcal{L}[e^{-t}] = \frac{1}{s+1}$. Therefore

$$y(t) = \mathcal{L}^{-1}[Y] = \frac{1}{3}\mathcal{L}^{-1}\left[\frac{1}{s-2}\right] + \frac{2}{3}\mathcal{L}^{-1}\left[\frac{1}{s+1}\right] = \boxed{\frac{1}{3}e^{2t} + \frac{2}{3}e^{-t}}.$$

Example 6.11. Solve

$$\begin{cases} y'' + y = \sin 2t \\ y(0) = 2 \\ y'(0) = 1. \end{cases}$$

$$\begin{aligned} y'' + y &= \sin 2t \\ \mathcal{L}[y''] + \mathcal{L}[y] &= \mathcal{L}[\sin 2t] \\ (s^2 Y - sy(0) - y'(0)) + Y &= \frac{2}{s^2 + 4} \\ s^2 Y - 2s - 1 + Y &= \frac{2}{s^2 + 4} \\ (s^2 + 1)Y &= 2s + 1 + \frac{2}{s^2 + 4} \end{aligned}$$

$$\begin{aligned} Y &= \frac{2s + 1}{s^2 + 1} + \frac{2}{(s^2 + 1)(s^2 + 4)} = \frac{2s + 1}{s^2 + 1} + \frac{As + B}{s^2 + 1} + \frac{Cs + D}{s^2 + 4} \\ &= \frac{2s}{s^2 + 1} + \frac{\frac{5}{3}}{s^2 + 1} - \frac{\frac{2}{3}}{s^2 + 4} = 2\mathcal{L}[\cos t] + \frac{5}{3}\mathcal{L}[\sin t] - \frac{1}{3}\mathcal{L}[\sin 2t] \end{aligned}$$

Therefore

$$y(t) = 2 \cos t + \frac{5}{3} \sin t - \frac{1}{3} \sin 2t.$$

Example 6.12. Solve

$$\begin{cases} y^{(4)} - y = 0 \\ y(0) = 0 \\ y'(0) = 1 \\ y''(0) = 0 \\ y'''(0) = 0. \end{cases}$$

Using the Laplace Transform we calculate that


$$\begin{aligned} 0 &= \mathcal{L}[y^{(4)}] - \mathcal{L}[y] = (s^4 Y - s^3 y(0) - s^2 y'(0) - s y''(0) - y'''(0)) - Y \\ &= s^4 Y - s^2 - Y = (s^4 - 1)Y - s^2. \end{aligned}$$

Thus

$$Y(s) = \frac{s^2}{s^4 - 1} = \frac{s^2}{(s^2 - 1)(s^2 + 1)} = \frac{\frac{1}{2}}{s^2 - 1} + \frac{\frac{1}{2}}{s^2 + 1}$$

Therefore

$$y = \frac{1}{2}\mathcal{L}^{-1}\left[\frac{1}{s^2 - 1}\right] + \frac{1}{2}\mathcal{L}^{-1}\left[\frac{1}{s^2 + 1}\right] = \boxed{\frac{1}{2}\sinh t + \frac{1}{2}\sin t.}$$




6.3 Step Functions


Definition. The *unit step function* $u_c : [0, \infty) \rightarrow \mathbb{R}$ is defined by

$$u_c(t) = \begin{cases} 0 & t < c \\ 1 & t \geq c \end{cases}$$

for $c \geq 0$.




Example 6.13. Draw the graph of $y = 1 - u_c(t)$.




Example 6.14. Draw the graph of $y = u_1(t) - u_2(t)$.

Clearly $t = 1$ and $t = 2$ are important points. So we consider the function on the intervals $[0, 1)$, $[1, 2)$ and $[2, \infty)$.

$$u_1(t) - u_2(t) = \begin{cases} u_1(t) - u_2(t) & 0 \leq t < 1 \\ u_1(t) - u_2(t) & 1 \leq t < 2 \\ u_1(t) - u_2(t) & 2 \leq t \end{cases}$$




$$\begin{aligned} u_1(t) - u_2(t) &= \begin{cases} u_1(t) - u_2(t) & 0 \leq t < 1 \\ u_1(t) - u_2(t) & 1 \leq t < 2 \\ u_1(t) - u_2(t) & 2 \leq t \end{cases} \\ &= \begin{cases} 0 - 0 & 0 \leq t < 1 \\ 1 - 0 & 1 \leq t < 2 \\ 1 - 1 & 2 \leq t \end{cases} = \begin{cases} 0 & 0 \leq t < 1 \\ 1 & 1 \leq t < 2 \\ 0 & 2 \leq t. \end{cases} \end{aligned}$$



Example 6.15. Write the function

$$f(t) = \begin{cases} 2 & 0 \leq t < 4 \\ 5 & 4 \leq t < 7 \\ -1 & 7 \leq t < 9 \\ 1 & 9 \leq t \end{cases}$$

in terms of the unit step function.




The function starts at $f(0) = 2$. So we will have

$$f(t) = 2 + (\text{something}).$$

At $t = 4$, the function jumps from 2 to 5 (it goes “up 3”). So

$$f(t) = 2 + 3u_4(t) + (\text{something}).$$



Then it goes “down 6” when $t = 7$. So

$$f(t) = 2 + 3u_4(t) - 6u_7(t) + (\text{something}).$$


Finally it goes “up 2” when $t = 9$. Therefore

$$f(t) = 2 + 3u_4(t) - 6u_7(t) + 2u_9(t).$$

Example 6.16. Write the function

$$f(t) = \begin{cases} t & 0 \leq t < 1 \\ t - 1 & 1 \leq t < 2 \\ t - 2 & 2 \leq t < 3 \\ 0 & 3 \leq t \end{cases}$$

in terms of the unit step function.



This function starts with $f(t) = t$, then changes when $t = 1$, $t = 2$ and $t = 3$: So we must have

$$f(t) = t + \left(\begin{smallmatrix} \text{first} \\ \text{jump} \end{smallmatrix} \right) u_1(t) + \left(\begin{smallmatrix} \text{second} \\ \text{jump} \end{smallmatrix} \right) u_2(t) + \left(\begin{smallmatrix} \text{third} \\ \text{jump} \end{smallmatrix} \right) u_3(t).$$

At each “jump” we calculate

$$\text{jump} = \left(\begin{array}{c} \text{function} \\ \text{on right} \end{array} \right) - \left(\begin{array}{c} \text{function} \\ \text{on left} \end{array} \right).$$

So we have

$$\left(\begin{array}{c} \text{first} \\ \text{jump} \end{array} \right) = (t - 1) - t = -1$$

$$\left(\begin{array}{c} \text{second} \\ \text{jump} \end{array} \right) = (t - 2) - (t - 1) = -1$$

$$\left(\begin{array}{c} \text{third} \\ \text{jump} \end{array} \right) = 0 - (t - 2) = 2 - t$$

Hence

$$f(t) = t - u_1(t) - u_2(t) + (2 - t)u_3(t).$$

What is the Laplace Transform of the unit step function?

We calculate that

$$\mathcal{L}[u_c](s) = \int_0^{\infty} e^{-st} u_c(t) dt = \int_c^{\infty} e^{-st} dt = \left[-\frac{1}{s} e^{-st} \right]_c^{\infty} = \frac{e^{-cs}}{s}$$


for $s > 0$.

Theorem 6.4.

$$\mathcal{L}[u_c](s) = \frac{e^{-cs}}{s}$$

Now suppose that we have some function $f : [0, \infty) \rightarrow \mathbb{R}$ and we define a new function $g : [0, \infty) \rightarrow \mathbb{R}$ by

$$g(t) = \begin{cases} 0 & t < c \\ f(t - c) & t \geq c. \end{cases}$$



We can write $g(t) = u_c(t)f(t - c)$. What is the Laplace Transform of g ?

$$\begin{aligned} \mathcal{L}[g] &= \mathcal{L}[u_c(t)f(t - c)] = \int_0^{\infty} e^{-st} u_c(t) f(t - c) dt \\ &= \int_c^{\infty} e^{-st} f(t - c) dt. \end{aligned}$$

Let $u = t - c$. Then $du = dt$ and $t = c \iff u = 0$. Therefore

$$\mathcal{L}[g] = \int_0^{\infty} e^{-s(u+c)} f(u) du = e^{-cs} \int_0^{\infty} e^{-su} f(u) du = e^{-cs} \mathcal{L}[f].$$

Theorem 6.5.

$$\mathcal{L}[u_c(t)f(t - c)](s) = e^{-cs} F(s)$$

Example 6.17. Find the Laplace Transform of

$$f(t) = \begin{cases} t & 0 \leq t < 1 \\ t - 1 & 1 \leq t < 2 \\ t - 2 & 2 \leq t < 3 \\ 0 & 3 \leq t. \end{cases}$$

Since

$$\begin{aligned} f(t) &= t - u_1(t) - u_2(t) + (2 - t)u_3(t) \\ &= t - u_1(t) - u_2(t) - u_3(t) - u_3(t)(t - 3) \end{aligned}$$

we have that

$$\begin{aligned} F(s) &= \mathcal{L}[t] - \mathcal{L}[u_1] - \mathcal{L}[u_2] - \mathcal{L}[u_3] - \mathcal{L}[u_3(t)(t - 3)] \\ &= \frac{1}{s^2} - \frac{e^{-s}}{s} - \frac{e^{-2s}}{s} - \frac{e^{-3s}}{s} - \frac{e^{-3s}}{s^2}. \end{aligned}$$

Example 6.18. Find the Laplace Transform of

$$f(t) = \begin{cases} \sin t & 0 \leq t \leq \frac{\pi}{4} \\ \sin t + \cos(t - \frac{\pi}{4}) & \frac{\pi}{4} \leq t. \end{cases}$$

Note that $f(t) = \sin t + g(t)$ where

$$g(t) = \begin{cases} 0 & 0 \leq t \leq \frac{\pi}{4} \\ \cos(t - \frac{\pi}{4}) & \frac{\pi}{4} \leq t \end{cases} = u_{\frac{\pi}{4}}(t) \cos\left(t - \frac{\pi}{4}\right).$$

So

$$\begin{aligned} F(s) &= \mathcal{L}[f] = \mathcal{L}[\sin t] + \mathcal{L}\left[u_{\frac{\pi}{4}}(t) \cos\left(t - \frac{\pi}{4}\right)\right] \\ &= \mathcal{L}[\sin t] + e^{-\frac{\pi s}{4}} \mathcal{L}[\cos t] = \frac{1}{s^2 + 1} + e^{-\frac{\pi s}{4}} \frac{s}{s^2 + 1} \\ &= \frac{1 + se^{-\frac{\pi s}{4}}}{s^2 + 1}. \end{aligned}$$

Example 6.19. Find the inverse Laplace Transform of $F(s) = \frac{1 - e^{-2s}}{s^2}$.

$$\begin{aligned} f(t) &= \mathcal{L}^{-1}[F] = \mathcal{L}^{-1}\left[\frac{1}{s^2}\right] - \mathcal{L}^{-1}\left[\frac{e^{-2s}}{s^2}\right] = t - u_2(t)(t - 2) \\ &= \begin{cases} t & 0 \leq t < 2 \\ 2 & t \geq 2. \end{cases} \end{aligned}$$

And what is the Laplace Transform of $e^{ct}f(t)$?

$$\mathcal{L}[e^{ct}f(t)] = \int_0^{\infty} e^{-st} e^{ct} f(t) dt = \int_0^{\infty} e^{-(s-c)t} f(t) dt = F(s - c).$$

Theorem 6.6.

$$\mathcal{L}[e^{ct}f(t)] = F(s - c)$$

Example 6.20. Find the inverse Laplace Transform of $G(s) = \frac{1}{s^2 - 4s + 5}$.

Note first that

$$G(s) = \frac{1}{s^2 - 4s + 5} = \frac{1}{(s - 2)^2 + 1}.$$

If $F(s) = \frac{1}{s^2 + 1}$, then we have $G(s) = F(s - 2)$. But $\mathcal{L}^{-1}[F] = \mathcal{L}^{-1}\left[\frac{1}{s^2 + 1}\right] = \sin t$. Therefore

$$g(t) = \mathcal{L}^{-1}[G] = \mathcal{L}^{-1}[F(s - 2)] = e^{2t} \mathcal{L}^{-1}[F] = e^{2t} \sin t.$$

6.4 ODEs with Discontinuous Forcing Functions

Example 6.21. Solve

$$\begin{cases} y'' + 4y = f(t) = \begin{cases} 0 & 0 \leq t < 5 \\ \frac{1}{5}(t - 5) & 5 \leq t < 10 \\ 1 & 10 \leq t \end{cases} \\ y(0) = 0 \\ y'(0) = 0. \end{cases}$$

Note that

$$\begin{aligned} f(t) &= 0 + \left(\frac{1}{5}(t - 5) - 0 \right) u_5(t) + \left(1 - \frac{1}{5}(t - 5) \right) u_{10}(t) \\ &= \frac{1}{5} (u_5(t)(t - 5) - u_{10}(t)(t - 10)). \end{aligned}$$

Taking the Laplace transform of the ODE gives

$$(s^2 + 4)Y = \frac{1}{5} \frac{e^{-5s} - e^{-10s}}{s^2}$$

and

$$Y = \frac{1}{5} \frac{e^{-5s} - e^{-10s}}{s^2(s^2 + 4)}.$$

Let

$$H(s) = \frac{1}{s^2(s^2 + 4)}.$$

Then

$$Y(s) = \frac{1}{5} e^{-5s} H(s) - \frac{1}{5} e^{-10s} H(s).$$

Recall that

$$\mathcal{L}[u_c(t)h(t - c)](s) = e^{-cs}H(s).$$

So

$$u_c(t)h(t - c)(s) = \mathcal{L}^{-1}[e^{-cs}H(s)].$$

If we can find $h(t)$, then we can find $y(t)$.

Using partial fractions, we calculate (please check!) that

$$\begin{aligned} H(s) &= \frac{1}{s^2(s^2 + 4)} = \frac{As + B}{s^2} + \frac{Cs + D}{s^2 + 4} \\ &= \frac{As^3 + Bs^2 + 4As + 4B + Cs^3 + Ds^2}{s^2(s^2 + 4)} \\ &= \frac{0s + \frac{1}{4}}{s^2} + \frac{0s - \frac{1}{4}}{s^2 + 4} = \frac{\frac{1}{4}}{s^2} - \frac{\frac{1}{4}}{s^2 + 4}. \end{aligned}$$

Hence

$$h(t) = \frac{1}{4} \mathcal{L}^{-1}\left[\frac{1}{s^2}\right] - \frac{1}{8} \mathcal{L}^{-1}\left[\frac{2}{s^2 + 4}\right] = \frac{t}{4} - \frac{1}{8} \sin 2t.$$

Therefore

$$\begin{aligned} y(t) &= \mathcal{L}^{-1} \left[\frac{1}{5} e^{-5s} H(s) - \frac{1}{5} e^{-10s} H(s) \right] \\ &= \frac{1}{5} u_5(t) h(t-5) - \frac{1}{5} u_{10}(t) h(t-10) \\ &= u_5(t) \left(\frac{t-5}{20} - \frac{1}{40} \sin(2t-10) \right) \\ &\quad - u_{10}(t) \left(\frac{t-10}{20} - \frac{1}{40} \sin(2t-20) \right). \end{aligned}$$

Example 6.22. Solve

$$\begin{cases} y'' + 3y' + 2y = f(t) = \begin{cases} 1 & 0 \leq t < 10 \\ 0 & 10 \leq t \end{cases} \\ y(0) = 1 \\ y'(0) = 0. \end{cases}$$

Since $f(t) = 1 - u_{10}(t)$, the Laplace Transform of the ODE is

$$(s^2 + 3s + 2)Y - (s + 3) = \frac{1 - e^{-10s}}{s}.$$

Thus

$$\begin{aligned} Y(s) &= \frac{1 - e^{-10s}}{s(s^2 + 3s + 2)} + \frac{s + 3}{s^2 + 3s + 2} \\ &= \frac{(s^2 + 3s + 1) - e^{-10s}}{s(s^2 + 3s + 2)}. \end{aligned}$$

Let

$$G(s) = \frac{s^2 + 3s + 1}{s(s^2 + 3s + 2)} \quad \text{and} \quad H(s) = \frac{1}{s(s^2 + 3s + 2)}.$$

Then $Y = G(s) - e^{-10s}H(s)$. If we can find $g(t)$ and $h(t)$, then we can find $y(t)$.

Using partial fractions we get

$$G(s) = \frac{A}{s} + \frac{B}{s+1} + \frac{C}{s+2} = \frac{\frac{1}{2}}{s} + \frac{1}{s+1} - \frac{\frac{1}{2}}{s+2}$$

and

$$H(s) = \frac{D}{s} + \frac{E}{s+1} + \frac{F}{s+2} = \frac{\frac{1}{2}}{s} - \frac{1}{s+1} + \frac{\frac{1}{2}}{s+2}$$

(please check!). It follows that

$$g(t) = \frac{1}{2} (1 + 2e^{-t} - e^{-2t}) \quad \text{and} \quad h(t) = \frac{1}{2} (1 - 2e^{-t} + e^{-2t}).$$

Therefore

$$\begin{aligned} y(t) &= \mathcal{L}^{-1} [Y] \\ &= \mathcal{L}^{-1} [G(s) - e^{-10s}H(s)] \\ &= g(t) - u_{10}(t)h(t-10) \\ &= \frac{1}{2} (1 + 2e^{-t} - e^{-2t}) - \frac{1}{2} u_{10}(t) (1 - 2e^{-(t-10)} + e^{-2(t-10)}). \end{aligned}$$

Example 6.23. Solve

$$\begin{cases} y'' + 4y = u_\pi(t) - u_{3\pi}(t) \\ y(0) = 0 \\ y'(0) = 0. \end{cases}$$

Taking the Laplace Transform of the ODE gives

$$(s^2 + 4)Y(s) = \frac{e^{-\pi s} - e^{-3\pi s}}{s}.$$

Thus

$$Y(s) = \frac{e^{-\pi s} - e^{-3\pi s}}{s(s^2 + 4)}.$$

Let

$$H(s) = \frac{1}{s(s^2 + 4)}.$$

Using partial fractions, we calculate that

$$\begin{aligned} H(s) &= \frac{1}{s(s^2 + 4)} = \frac{A}{s} + \frac{Bs + C}{s^2 + 4} = \frac{\frac{1}{4}}{s} + \frac{-\frac{1}{4}s + 0}{s^2 + 4} \\ &= \frac{1}{4} \left(\frac{1}{s} \right) - \frac{1}{4} \left(\frac{s}{s^2 + 4} \right) = \frac{1}{4} \mathcal{L} [1] - \frac{1}{4} \mathcal{L} [\cos 2t]. \end{aligned}$$

It follows that

$$h(t) = \frac{1}{4} - \frac{1}{4} \cos 2t$$

and the solution to the IVP is

$$\begin{aligned} y(t) &= \mathcal{L}^{-1} [e^{-\pi s} H(s)] - \mathcal{L}^{-1} [e^{-3\pi s} H(s)] \\ &= u_\pi(t)h(t - \pi) - u_{3\pi}(t)h(t - 3\pi) \\ &= \frac{1}{4}u_\pi(t)(1 - \cos(2t - 2\pi)) - \frac{1}{4}u_{3\pi}(t)(1 - \cos(2t - 6\pi)). \end{aligned}$$

6.6 The Convolution Integral

Let $f : [0, \infty) \rightarrow \mathbb{R}$ and $g : [0, \infty) \rightarrow \mathbb{R}$ be piecewise continuous functions.

Definition. The *convolution* of f and g is

$$(f * g)(t) = \int_0^t f(\tau)g(t - \tau) d\tau.$$

Theorem 6.7 (Properties).

- $f * g = g * f$
- $f * (g * h) = (f * g) * h$
- $f * (g + h) = (f * g) + (f * h)$
- $f * 0 = 0 = 0 * f$

Example 6.24.

$$(\cos * 1)(t) = \int_0^t \cos \tau \cdot 1 d\tau = [\sin \tau]_0^t = \sin t - \sin 0 = \sin t$$

$$(1 * \cos)(t) = \int_0^t 1 \cdot \cos(t - \tau) d\tau = [-\sin(t - \tau)]_0^t = -\sin 0 + \sin t = \sin t$$

Note that $f * 1 \neq f$ in general.

Example 6.25.

$$\begin{aligned} (\sin * \sin)(t) &= \int_0^t \sin \tau \sin(t - \tau) d\tau = \int_0^t \sin \tau (\sin t \cos \tau - \cos t \sin \tau) d\tau \\ &= \sin t \int_0^t \sin \tau \cos \tau d\tau - \cos t \int_0^t \sin^2 \tau d\tau \\ &= \sin t \left[-\frac{1}{2} \cos^2 \tau \right]_0^t - \cos t \left[\frac{1}{2} (\tau - \sin \tau \cos \tau) \right]_0^t \\ &= \frac{1}{2} \sin t (1 - \cos^2 t) - \frac{1}{2} \cos t (t - \sin t \cos t) \\ &= \frac{1}{2} \sin t - \frac{t}{2} \cos t. \end{aligned}$$

Note that $f * f \geq 0$ is not true in general.

Theorem 6.8.

$$\mathcal{L}[f * g](s) = F(s)G(s)$$

This means that $\mathcal{L}^{-1}[FG] = f * g$.

Example 6.26. Find the inverse Laplace Transform of $H(s) = \frac{a}{s^2(s^2 + a^2)}$.

Note that $H(s) = \left(\frac{1}{s^2}\right) \left(\frac{a}{s^2 + a^2}\right)$. We know that $\mathcal{L}[t] = \frac{1}{s^2}$ and $\mathcal{L}[\sin at] = \frac{a}{s^2 + a^2}$. So

$$\begin{aligned} h(t) &= \mathcal{L}^{-1} \left[\left(\frac{1}{s^2}\right) \left(\frac{a}{s^2 + a^2}\right) \right] = \mathcal{L}^{-1} \left[\frac{1}{s^2} \right] * \mathcal{L}^{-1} \left[\frac{a}{s^2 + a^2} \right] \\ &= t * \sin at = \int_0^t \tau \sin a(t - \tau) d\tau \\ &= \frac{at - \sin at}{a^2}. \end{aligned}$$

Example 6.27. Solve

$$\begin{cases} y'' + 4y = g(t) \\ y(0) = 3 \\ y'(0) = -1. \end{cases}$$

Taking the Laplace Transform of the ODE gives

$$(s^2 Y - 3s + 1) + 4Y = G(s)$$

which rearranges to

$$\begin{aligned} Y(s) &= \frac{3s - 1}{s^2 + 4} + \frac{G(s)}{s^2 + 4} \\ &= 3 \left(\frac{s}{s^2 + 4} \right) - \frac{1}{2} \left(\frac{2}{s^2 + 4} \right) + \frac{1}{2} \left(\frac{2}{s^2 + 4} \right) G(s). \end{aligned}$$

Hence the solution to the IVP is

$$\begin{aligned} y(t) &= 3\mathcal{L}^{-1} \left[\frac{s}{s^2 + 4} \right] - \frac{1}{2} \mathcal{L}^{-1} \left[\frac{2}{s^2 + 4} \right] + \frac{1}{2} \mathcal{L}^{-1} \left[\left(\frac{2}{s^2 + 4} \right) G(s) \right] \\ &= 3 \cos 2t - \frac{1}{2} \sin 2t + \frac{1}{2} \sin 2t * g(t) \\ &= 3 \cos 2t - \frac{1}{2} \sin 2t + \frac{1}{2} \int_0^t \sin 2(t - \tau) g(\tau) d\tau. \end{aligned}$$


Example 6.28. Find the inverse Laplace Transform of $\frac{2}{(s - 1)(s^2 + 4)}$.

$$\begin{aligned} \mathcal{L}^{-1} \left[\frac{2}{(s - 1)(s^2 + 4)} \right] &= \mathcal{L}^{-1} \left[\left(\frac{2}{s^2 + 4} \right) \left(\frac{1}{s - 1} \right) \right] = \sin 2t * e^t \\ &= \int_0^t e^{t-\tau} \sin 2\tau d\tau = e^t \int_0^t e^{-\tau} \sin 2\tau d\tau \\ &= e^t \left[\frac{e^{-\tau}}{5} (-\sin 2\tau - 2 \cos 2\tau) \right]_0^t \\ &= \frac{2}{5} e^t - \frac{1}{5} \sin 2t - \frac{2}{5} \cos 2t. \end{aligned}$$

7

Systems of First Order Linear Equations

7.1 Introduction



Consider the dynamical system shown above. There are two blocks and three springs. Forces F_1 and F_2 act on the blocks as shown.

See <https://tinyurl.com/wm2ogdh>

We expect that the acceleration of the blocks will depend on

- the displacements x_1 and x_2 ;
- the forces F_1 and F_2 ; and
- the masses of the blocks.

So we expect that:

$$\begin{cases} \frac{d^2 x_1}{dt^2} = f_1(x_1, x_2, F_1, m_1) \\ \frac{d^2 x_2}{dt^2} = f_2(x_1, x_2, F_2, m_2). \end{cases}$$

This is a system of two ODEs. To find $x_1(t)$ and $x_2(t)$, we would need to solve these equations at the same time.

The most famous system of ODEs is the system of *Predator-Prey* equations:

$$\begin{cases} \frac{dx}{dt} = \alpha x - \beta xy \\ \frac{dy}{dt} = \delta xy - \gamma y \end{cases}$$

where

$x(t)$ = number of prey (e.g. mice)

$y(t)$ = number of predators (e.g. owls),

which originate circa 1925.

It is possible to convert an n th order linear ODE into a system of n first order linear ODEs. Or vice versa.

$$a_n y^{(n)} + a_{n-1} y^{(n-1)} + \dots + a_1 y' + a_0 y = g(t) \quad \longleftrightarrow \quad \begin{cases} x'_1 = b_{11}x_1 + \dots + b_{1n}x_n + h_1(t) \\ x'_2 = b_{21}x_1 + \dots + b_{2n}x_n + h_2(t) \\ \vdots \\ x'_n = b_{n1}x_1 + \dots + b_{nn}x_n + h_n(t) \end{cases}$$

Example 7.1. Write

$$u'' + 0.25u' + u = 0$$

as a system of two first order ODEs.

Let $x_1 = u$ and $x_2 = u'$. Then clearly $x'_1 = u' = x_2$ and

$$x'_2 = u'' = -0.25u' - u = -0.25x_2 - x_1.$$

Therefore

$$\begin{cases} x'_1 = x_2 \\ x'_2 = -x_1 - 0.25x_2. \end{cases}$$

Remark. We will need

- matrices,
- eigenvalues,
- eigenvectors,
- the Wronskian,
- linear independence,
- and more

from MATH215 – please either revise your linear algebra lecture notes or read §7.2-7.3 in the textbook.

7.4 Basic Theory of Systems of First Order Linear Equations

$$\begin{cases} x'_1 = p_{11}(t)x_1 + p_{12}(t)x_2 + \dots + p_{1n}(t)x_n + g_1(t) \\ x'_2 = p_{21}(t)x_1 + p_{22}(t)x_2 + \dots + p_{2n}(t)x_n + g_2(t) \\ \vdots \\ x'_n = p_{n1}(t)x_1 + p_{n2}(t)x_2 + \dots + p_{nn}(t)x_n + g_n(t) \end{cases}$$

is a system of n linear ODEs and n variables: x_1, x_2, \dots, x_n .

If we write

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad \mathbf{x}' = \begin{bmatrix} x'_1 \\ x'_2 \\ \vdots \\ x'_n \end{bmatrix}, \quad P = \begin{bmatrix} p_{11} & p_{12} & \dots & p_{1n} \\ p_{21} & p_{22} & \dots & p_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ p_{n1} & p_{n2} & \dots & p_{nn} \end{bmatrix}, \quad \text{and } \mathbf{g} = \begin{bmatrix} g_1 \\ g_2 \\ \vdots \\ g_n \end{bmatrix}$$

then we can write this system as

$$\mathbf{x}' = P(t)\mathbf{x} + \mathbf{g}(t).$$

First we will consider the homogeneous system

$$\mathbf{x}' = P(t)\mathbf{x}.$$

In Chapters 3 and 4 when we had multiple solutions, we wrote them as $y_1(t), y_2(t), \dots$. But we are already using x_1, x_2, \dots to denote coordinates. So we need a new type of notation.

Notation. We use $\mathbf{x}^{(1)}(t), \mathbf{x}^{(2)}(t), \dots$ to denote different vector solutions.

Recall from Chapter 3 that if $y_1(t)$ and $y_2(t)$ are both solutions to

$$ay'' + by' + cy = 0,$$

then

$$c_1y_1 + c_2y_2$$

is also a solution.

Theorem 7.1. If $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ are solutions to $\mathbf{x}' = P(t)\mathbf{x}$, then

$$c_1\mathbf{x}^{(1)} + c_2\mathbf{x}^{(2)}$$

is also a solution for any $c_1, c_2 \in \mathbb{R}$.

Example 7.2. $\mathbf{x}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t}$ and $\mathbf{x}^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}$ are both solutions to $\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}$ (we will see this later). Therefore

$$\mathbf{x}(t) = c_1\mathbf{x}^{(1)} + c_2\mathbf{x}^{(2)} = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}$$

is also a solution to this system.

(Suppose that $P(t)$ is an $n \times n$ matrix.)

Theorem 7.2. If $\mathbf{x}^{(1)}(t), \mathbf{x}^{(2)}(t), \dots, \mathbf{x}^{(n)}(t)$ are linearly independent solutions to $\mathbf{x}' = P(t)\mathbf{x}$, then every solution to this system can be written as

$$\mathbf{x}(t) = c_1\mathbf{x}^{(1)} + c_2\mathbf{x}^{(2)} + \dots + c_n\mathbf{x}^{(n)}$$

in exactly one way.

Definition. In this case, we say that $\mathbf{x}^{(1)}(t), \mathbf{x}^{(2)}(t), \dots, \mathbf{x}^{(n)}(t)$ form a *fundamental set of solutions* to $\mathbf{x}' = P(t)\mathbf{x}$.

Definition. In this case,

$$\mathbf{x}(t) = c_1\mathbf{x}^{(1)} + c_2\mathbf{x}^{(2)} + \dots + c_n\mathbf{x}^{(n)}$$

is called the *general solution* to $\mathbf{x}' = P(t)\mathbf{x}$.

7.5 Homogeneous Linear Systems with Constant Coefficients

Consider

$$\mathbf{x}' = A\mathbf{x}$$

where $A \in \mathbb{R}^{n \times n}$.

If $n = 1$, then we just have

$$\frac{dx}{dt} = ax$$

which has general solution $x(t) = ce^{at}$.

For $n > 1$, we guess that

$$\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$$

is a solution to $\mathbf{x}' = A\mathbf{x}$, for some number $r \in \mathbb{C}$ and some vector $\boldsymbol{\xi} \in \mathbb{C}^n$.

But if $\mathbf{x}(t) = \boldsymbol{\xi}e^{rt}$, then

$$\begin{aligned} r\boldsymbol{\xi}e^{rt} &= \mathbf{x}' = A\mathbf{x} = A\boldsymbol{\xi}e^{rt} \\ r\boldsymbol{\xi} &= A\boldsymbol{\xi} \\ (A - rI)\boldsymbol{\xi} &= \mathbf{0} \end{aligned}$$

where I is the identity matrix. Hence r must be an eigenvalue of A and $\boldsymbol{\xi}$ must be a corresponding eigenvector of A .

Remark. So the idea is:

- (i). Find the eigenvalues;
- (ii). Find the eigenvectors; then
- (iii). Write $\mathbf{x}^{(j)}(t) = \boldsymbol{\xi}^{(j)}e^{r_j t}$.

Example 7.3. Solve

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

First we find the eigenvalues. Since

$$\begin{aligned} 0 &= \det(A - rI) = \begin{vmatrix} 1-r & 1 \\ 4 & 1-r \end{vmatrix} = (1-r)^2 - 4 \\ &= r^2 - 2r - 3 = (r+1)(r-3), \end{aligned}$$

the eigenvalues are $r_1 = 3$ and $r_2 = -1$.

Using the first eigenvalue $r_1 = 3$, we calculate that

$$\mathbf{0} = (A - r_1 I)\boldsymbol{\xi} = \begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \implies 0 = -2\xi_1 + \xi_2.$$

Hence we can choose $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$. Then using the second eigenvalue $r_2 = -1$, we calculate that

$$\mathbf{0} = (A - r_2 I)\boldsymbol{\xi} = \begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \implies 0 = 2\xi_1 + \xi_2.$$

Hence we can choose $\xi^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$. This gives us two solutions:

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}.$$

But are these two solutions linearly independent? To find out, we calculate the Wronskian of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$:

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} = \begin{bmatrix} e^{3t} \\ 2e^{3t} \end{bmatrix} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t} = \begin{bmatrix} e^{-t} \\ -2e^{-t} \end{bmatrix}.$$

$$W(\mathbf{x}^{(1)}, \mathbf{x}^{(2)})(t) = \begin{vmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{vmatrix} = -4e^{2t} \neq 0.$$

Since $W \neq 0$, we have that $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ are linearly independent. So $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ form a fundamental set of solutions. Therefore the general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}.$$

Example 7.4. Solve

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} 8 & -1 \\ 6 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 1 \\ -2 \end{bmatrix}. \end{cases}$$

The eigenvalues are $r_1 = 7$ and $r_2 = 2$. The corresponding eigenvectors are $\xi^{(1)} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\xi^{(2)} = \begin{bmatrix} 1 \\ 6 \end{bmatrix}$. Therefore the general solution to the ODE is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

Setting $t = 0$, we have

$$\begin{bmatrix} 1 \\ -2 \end{bmatrix} = \mathbf{x}(0) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 6 \end{bmatrix} = \begin{bmatrix} c_1 + c_2 \\ c_1 + 6c_2 \end{bmatrix} \implies \begin{cases} c_1 + c_2 = 1 \\ c_1 + 6c_2 = -2 \end{cases} \implies \begin{cases} c_1 = \frac{8}{5} \\ c_2 = -\frac{3}{5} \end{cases}.$$

Therefore the solution to the IVP is

$$\mathbf{x}(t) = \frac{8}{5} \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{7t} - \frac{3}{5} \begin{bmatrix} 1 \\ 6 \end{bmatrix} e^{2t}.$$

Example 7.5. Solve

$$\mathbf{x}' = \begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} \mathbf{x}.$$

The eigenvalues are $r_1 = -1$ and $r_2 = -4$. The corresponding eigenvectors are $\xi^{(1)} = \begin{bmatrix} 1 \\ \sqrt{2} \end{bmatrix}$ and $\xi^{(2)} = \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$. Hence the general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ \sqrt{2} \end{bmatrix} e^{-t} + c_2 \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix} e^{-4t}.$$

Remark.

$$\det(A - rI) = 0$$

There are three possibilities for the eigenvalues of A .

- (i). All the eigenvalues are real and different;
- (ii). Some eigenvalues occur in complex conjugate pairs;
- (iii). Some eigenvalues are repeated.

If all the eigenvalues are real and different, then the eigenvectors are linearly independent: So $W(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)})(t) \neq 0$ and $\mathbf{x}^{(1)}(t), \dots, \mathbf{x}^{(n)}(t)$ form a fundamental set of solutions.

If some eigenvalues are repeated, *but there are n linearly independent eigenvectors*, then this is also true: $\mathbf{x}^{(1)}(t), \dots, \mathbf{x}^{(n)}(t)$ form a fundamental set of solutions.

Example 7.6. Solve

$$\mathbf{x}' = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \mathbf{x}.$$

The eigenvalues and eigenvectors are

$$\begin{array}{ccc} r_1 = 2 & r_2 = -1 & r_3 = -1 \\ \boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} & \boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} & \boldsymbol{\xi}^{(3)} = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \end{array}$$

which gives us the following three solutions

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} e^{2t} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} e^{-t} \quad \mathbf{x}^{(3)}(t) = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} e^{-t}.$$

You can check that the Wronskian of $\mathbf{x}^{(1)}$, $\mathbf{x}^{(2)}$ and $\mathbf{x}^{(3)}$ is non-zero. Therefore $\mathbf{x}^{(1)}$, $\mathbf{x}^{(2)}$ and $\mathbf{x}^{(3)}$ form a fundamental set of solutions. The general solution to the ODE is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} e^{2t} + c_2 \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} e^{-t} + c_3 \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} e^{-t}.$$

Remark. Next we will study systems with complex eigenvalues.

7.6 Complex Eigenvalues

Consider

$$\mathbf{x}' = A\mathbf{x}$$

where $A \in \mathbb{R}^{n \times n}$.

Any complex eigenvalues of A must occur in complex conjugate pairs: If $r_1 = \lambda + i\mu$ is an eigenvalue of A , then $r_2 = \bar{r}_1 = \lambda - i\mu$ is also an eigenvalue of A .

Moreover, if $\boldsymbol{\xi}^{(1)}$ is an eigenvector of A corresponding to r_1 , then $\boldsymbol{\xi}^{(2)} = \overline{\boldsymbol{\xi}^{(1)}}$ is an eigenvector of A corresponding to $r_2 = \bar{r}_1$.

Two solutions of $\mathbf{x}' = A\mathbf{x}$ are

$$\mathbf{x}^{(1)}(t) = \boldsymbol{\xi}^{(1)} e^{r_1 t} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \overline{\boldsymbol{\xi}^{(1)}} e^{\bar{r}_1 t}.$$

But $\mathbf{x}^{(1)}, \mathbf{x}^{(2)} : \mathbb{R} \rightarrow \mathbb{C}^n$ and we want solutions $\mathbb{R} \rightarrow \mathbb{R}^n$.

If $r_1 = \lambda + i\mu$, and $\boldsymbol{\xi}^{(1)} = \mathbf{a} + i\mathbf{b}$ ($\lambda, \mu \in \mathbb{R}$, $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$), then

$$\begin{aligned} \mathbf{x}^{(1)}(t) &= (\mathbf{a} + i\mathbf{b}) e^{(\lambda + i\mu)t} \\ &= (\mathbf{a} + i\mathbf{b}) e^{\lambda t} (\cos \mu t + i \sin \mu t) \\ &= e^{\lambda t} (\mathbf{a} \cos \mu t - \mathbf{b} \sin \mu t) + i e^{\lambda t} (\mathbf{a} \sin \mu t + \mathbf{b} \cos \mu t) \\ &= \mathbf{u}(t) + i\mathbf{v}(t). \end{aligned}$$

The functions $\mathbf{u}(t)$ and $\mathbf{v}(t)$ will be linearly independent. Furthermore

$$\text{span}\{\mathbf{u}(t), \mathbf{v}(t)\} = \text{span}\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}\}.$$

So we can include $\mathbf{u}(t)$ and $\mathbf{v}(t)$ in our fundamental set of solutions instead of $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$.

Example 7.7. Solve

$$\mathbf{x}' = \begin{bmatrix} -\frac{1}{2} & 1 \\ -1 & -\frac{1}{2} \end{bmatrix} \mathbf{x}.$$

We calculate that

$$0 = \det(A - rI) = \begin{vmatrix} -\frac{1}{2} - r & 1 \\ -1 & -\frac{1}{2} - r \end{vmatrix} = r^2 + r + \frac{5}{4}$$

and

$$r_{1,2} = \frac{-1 \pm \sqrt{1-5}}{2} = \frac{-1 \pm 2i}{2} = -\frac{1}{2} \pm i.$$

So we have $r_1 = -\frac{1}{2} + i$ and $r_2 = -\frac{1}{2} - i$. We will use r_1 . We do not need r_2 .

Since

$$0 = (A - r_1 I) \boldsymbol{\xi}^{(1)} = \begin{bmatrix} -i & 1 \\ -1 & -i \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \implies \begin{cases} -i\xi_1 + \xi_2 = 0 \\ -\xi_1 - i\xi_2 = 0 \end{cases}$$

we can choose

$$\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ i \end{bmatrix}.$$

Note that we also have

$$\boldsymbol{\xi}^{(2)} = \overline{\boldsymbol{\xi}^{(1)}} = \overline{\begin{bmatrix} 1 \\ i \end{bmatrix}} = \begin{bmatrix} 1 \\ -i \end{bmatrix},$$

but we don't need $\boldsymbol{\xi}^{(2)}$.

Next we look at $\mathbf{x}^{(1)}(t)$:

$$\begin{aligned}\mathbf{x}^{(1)}(t) &= \boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 1 \\ i \end{bmatrix} e^{-\frac{t}{2}} (\cos t + i \sin t) \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t + i \sin t \\ i \cos t - \sin t \end{bmatrix} \\ &= e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} + i e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix} \\ &= \mathbf{u}(t) + i \mathbf{v}(t).\end{aligned}$$

Hence we choose

$$\mathbf{u}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} \quad \text{and} \quad \mathbf{v}(t) = e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}.$$

But are $\mathbf{u}(t)$ and $\mathbf{v}(t)$ linearly independent? Since

$$\begin{aligned}W(\mathbf{u}(t), \mathbf{v}(t))(t) &= \begin{vmatrix} u_1 & v_1 \\ u_2 & v_2 \end{vmatrix} = \begin{vmatrix} e^{-\frac{t}{2}} \cos t & e^{-\frac{t}{2}} \sin t \\ -e^{-\frac{t}{2}} \sin t & e^{-\frac{t}{2}} \cos t \end{vmatrix} \\ &= e^{-t} \cos^2 t + e^{-t} \sin^2 t = e^{-t} \\ &\neq 0\end{aligned}$$

the answer is yes. Therefore $\mathbf{u}(t)$ and $\mathbf{v}(t)$ form a fundamental set of solutions.

Therefore the general solution to $\mathbf{x}' = \begin{bmatrix} -\frac{1}{2} & 1 \\ -1 & -\frac{1}{2} \end{bmatrix} \mathbf{x}$ is


$$\mathbf{x}(t) = c_1 e^{-\frac{t}{2}} \begin{bmatrix} \cos t \\ -\sin t \end{bmatrix} + c_2 e^{-\frac{t}{2}} \begin{bmatrix} \sin t \\ \cos t \end{bmatrix}.$$

Remark. Our method is

1. Find the eigenvalues;
2. Find the eigenvectors;
3.
 - If r_j is real, just use the solution $\mathbf{x}^{(j)}(t) = \boldsymbol{\xi}^{(j)} e^{r_j t}$;
 - But if r_j is complex, write

$$\mathbf{x}^{(j)}(t) = \boldsymbol{\xi}^{(j)} e^{r_j t} = \begin{pmatrix} \text{real valued} \\ \text{function} \end{pmatrix} + i \begin{pmatrix} \text{real valued} \\ \text{function} \end{pmatrix}$$

and use these two functions.



Example 7.8. The electric circuit shown above is described by

$$\begin{cases} I' = -I - V \\ V' = 2I - V \end{cases}$$

where

I = the current through the inductor

V = the voltage drop across the capacitor.

(Ask an Electrical Engineer.)

Suppose that at time $t = 0$ the current is 2 amperes and the voltage drop is 2 volts. Find $I(t)$ and $V(t)$.

We must solve the IVP

$$\begin{cases} \frac{d}{dt} \begin{bmatrix} I \\ V \end{bmatrix} = \begin{bmatrix} -1 & -1 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} I \\ V \end{bmatrix} \\ \begin{bmatrix} I \\ V \end{bmatrix} (0) = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \end{cases}.$$

The eigenvalues of $\begin{bmatrix} -1 & -1 \\ 2 & -1 \end{bmatrix}$ are $r_1 = -1 + i\sqrt{2}$ and $r_2 = -1 - i\sqrt{2}$ (please check). The corresponding eigenvectors are

$$\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ -i\sqrt{2} \end{bmatrix} \quad \text{and} \quad \boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ i\sqrt{2} \end{bmatrix}.$$

Then we calculate that

$$\begin{aligned} \mathbf{x}^{(1)}(t) &= \boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 1 \\ -i\sqrt{2} \end{bmatrix} e^{(-1+i\sqrt{2})t} \\ &= \begin{bmatrix} 1 \\ -i\sqrt{2} \end{bmatrix} e^{-t} (\cos \sqrt{2}t + i \sin \sqrt{2}t) \\ &= e^{-t} \begin{bmatrix} \cos \sqrt{2}t + i \sin \sqrt{2}t \\ -i\sqrt{2} \cos \sqrt{2}t + \sqrt{2} \sin \sqrt{2}t \end{bmatrix} \\ &= e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} + i e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}. \end{aligned}$$

Hence the general solution to the ODE is

$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = c_1 e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} + c_2 e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}.$$

Using the initial condition, we calculate that

$$\begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} I(0) \\ V(0) \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ -\sqrt{2} \end{bmatrix} \quad \Rightarrow \quad \begin{cases} c_1 = 2 \\ c_2 = -\sqrt{2}. \end{cases}$$

Thus


$$\begin{bmatrix} I(t) \\ V(t) \end{bmatrix} = 2 e^{-t} \begin{bmatrix} \cos \sqrt{2}t \\ \sqrt{2} \sin \sqrt{2}t \end{bmatrix} - \sqrt{2} e^{-t} \begin{bmatrix} \sin \sqrt{2}t \\ -\sqrt{2} \cos \sqrt{2}t \end{bmatrix}.$$

So the answers to this problem are

$$\boxed{I(t) = 2e^{-t} \cos \sqrt{2}t - \sqrt{2}e^{-t} \sin \sqrt{2}t}$$

and


$$\boxed{V(t) = 2\sqrt{2}e^{-t} \sin \sqrt{2}t + 2e^{-t} \cos \sqrt{2}t.}$$



See <https://tinyurl.com/wm2ogdh> for an animated figure.

Example 7.9. For the dynamical system shown above, find $x_1(t)$ and $x_2(t)$.

As the springs are stretched and compressed, they apply forces on the blocks as shown below (Hooke's Law).



We calculate that

$$2 \frac{d^2 x_1}{dt^2} = \text{mass} \times \text{acceleration} = \text{force} = -x_1 + 3(x_2 - x_1)$$

$$\frac{9}{4} \frac{d^2 x_2}{dt^2} = \text{mass} \times \text{acceleration} = \text{force} = -3(x_2 - x_1) - \frac{15}{4}x_2.$$

This is a system of 2 second order ODEs. We want a system of first order ODEs.

Now let $y_1 = x_1$, $y_2 = x_2$, $y_3 = x_1'$ and $y_4 = x_2'$. Then

$$y_1' = x_1' = y_3$$

$$y_2' = x_2' = y_4$$

$$y_3' = x_1'' = \frac{1}{2} \left(-x_1 + 3x_2 - 3x_1 \right) = -2y_1 + \frac{3}{2}y_2$$

$$y_4' = x_2'' = \frac{4}{9} \left(-3x_2 + 3x_1 - \frac{15}{4}x_2 \right) = \frac{4}{3}y_1 - 3y_2.$$

So

$$\mathbf{y}' = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -2 & \frac{3}{2} & 0 & 0 \\ \frac{4}{3} & -3 & 0 & 0 \end{bmatrix} \mathbf{y}.$$

The characteristic polynomial of this matrix is

$$0 = r^4 + 5r^2 + 4 = (r^2 + 1)(r^2 + 4).$$

So $r_1 = i$, $r_2 = -i$, $r_3 = 2i$ and $r_4 = -2i$. We will use r_1 and r_3 (we do not need r_2 and r_4).

The corresponding eigenvectors (please check) are

$$\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 3 \\ 2 \\ 3i \\ 2i \end{bmatrix} \quad \text{and} \quad \boldsymbol{\xi}^{(3)} = \begin{bmatrix} 3 \\ -4 \\ 6i \\ -8i \end{bmatrix}.$$

It follows that

$$\boldsymbol{\xi}^{(1)} e^{r_1 t} = \begin{bmatrix} 3 \\ 2 \\ 3i \\ 2i \end{bmatrix} (\cos t + i \sin t) = \begin{bmatrix} 3 \cos t \\ 2 \cos t \\ -3 \sin t \\ -2 \sin t \end{bmatrix} + i \begin{bmatrix} 3 \sin t \\ 2 \sin t \\ 3 \cos t \\ 2 \cos t \end{bmatrix} = \mathbf{u}(t) + i\mathbf{v}(t)$$

and

$$\boldsymbol{\xi}^{(3)} e^{r_3 t} = \begin{bmatrix} 3 \\ -4 \\ 6i \\ -8i \end{bmatrix} (\cos 2t + i \sin 2t) = \begin{bmatrix} 3 \cos 2t \\ -4 \cos 2t \\ -6 \sin 2t \\ 8 \sin 2t \end{bmatrix} + i \begin{bmatrix} 3 \sin 2t \\ -4 \sin 2t \\ 6 \cos 2t \\ -8 \cos 2t \end{bmatrix} = \mathbf{w}(t) + i\mathbf{z}(t)$$

Therefore the general solution is

$$\begin{aligned} \mathbf{y}(t) &= c_1 \mathbf{u}(t) + c_2 \mathbf{v}(t) + c_3 \mathbf{w}(t) + c_4 \mathbf{z}(t) \\ &= c_1 \begin{bmatrix} 3 \cos t \\ 2 \cos t \\ -3 \sin t \\ -2 \sin t \end{bmatrix} + c_2 \begin{bmatrix} 3 \sin t \\ 2 \sin t \\ 3 \cos t \\ 2 \cos t \end{bmatrix} + c_3 \begin{bmatrix} 3 \cos 2t \\ -4 \cos 2t \\ -6 \sin 2t \\ 8 \sin 2t \end{bmatrix} + c_4 \begin{bmatrix} 3 \sin 2t \\ -4 \sin 2t \\ 6 \cos 2t \\ -8 \cos 2t \end{bmatrix}. \end{aligned}$$

Example 7.10. Suppose that the above system has initial condition

$$\mathbf{y}(0) = \begin{bmatrix} -1 \\ 4 \\ 1 \\ 1 \end{bmatrix}.$$

Sketch graphs of $y_1(t)$ and $y_2(t)$.



The initial value problem

$$\mathbf{y}' = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -2 & \frac{3}{2} & 0 & 0 \\ \frac{4}{3} & -3 & 0 & 0 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} -1 \\ 4 \\ 1 \\ 1 \end{bmatrix}$$

has solution

$$\mathbf{y}(t) = \frac{4}{9} \begin{bmatrix} 3 \cos t \\ 2 \cos t \\ -3 \sin t \\ -2 \sin t \end{bmatrix} + \frac{7}{18} \begin{bmatrix} 3 \sin t \\ 2 \sin t \\ 3 \cos t \\ 2 \cos t \end{bmatrix} - \frac{7}{9} \begin{bmatrix} 3 \cos 2t \\ -4 \cos 2t \\ -6 \sin 2t \\ 8 \sin 2t \end{bmatrix} - \frac{1}{36} \begin{bmatrix} 3 \sin 2t \\ -4 \sin 2t \\ 6 \cos 2t \\ -8 \cos 2t \end{bmatrix}.$$

Then we can draw the graphs of y_1 and y_2 :



Please see <https://tinyurl.com/s7uw7m>

7.7 Fundamental Matrices

Now consider

$$\mathbf{x}' = P(t)\mathbf{x}$$

where P is an $n \times n$ matrix. Suppose that $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(n)}$ are linearly independent solutions to this ODE. In other words, suppose that $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(n)}$ form a *fundamental set of solutions* to this ODE.

Definition. The matrix

$$\Psi(t) = \begin{bmatrix} \mathbf{x}^{(1)} & \mathbf{x}^{(2)} & \dots & \mathbf{x}^{(n)} \end{bmatrix} = \begin{bmatrix} x_1^{(1)}(t) & x_1^{(2)}(t) & \dots & x_1^{(n)}(t) \\ x_2^{(1)}(t) & x_2^{(2)}(t) & \dots & x_2^{(n)}(t) \\ \vdots & \vdots & & \vdots \\ x_n^{(1)}(t) & x_n^{(2)}(t) & \dots & x_n^{(n)}(t) \end{bmatrix}$$

is called a *fundamental matrix* of $\mathbf{x}' = P(t)\mathbf{x}$.

Example 7.11. Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Recall that

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} \quad \text{and} \quad \mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}$$

form a fundamental set of solutions to this ODE. Therefore

$$\Psi(t) = \begin{bmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{bmatrix}$$

is a fundamental matrix of this ODE.

Now, the general solution to

$$\mathbf{x}' = A\mathbf{x}$$

is

$$\mathbf{x}(t) = c_1 \mathbf{x}^{(1)}(t) + c_2 \mathbf{x}^{(2)}(t) + \dots + c_n \mathbf{x}^{(n)}(t) = \Psi(t)\mathbf{c}$$

where

$$\mathbf{c} = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \in \mathbb{R}^n.$$

If we have an initial condition $\mathbf{x}(t_0) = \mathbf{x}^0$, then

$$\Psi(t_0)\mathbf{c} = \mathbf{x}^0.$$

But

$$\begin{array}{l} \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)} \\ \text{are linearly} \\ \text{independent} \end{array} \implies \Psi(t) \text{ is invertible} \implies \mathbf{c} = \Psi^{-1}(t_0)\mathbf{x}^0.$$

Therefore the solution to the IVP

$$\begin{cases} \mathbf{x}' = A\mathbf{x} \\ \mathbf{x}(t_0) = \mathbf{x}^0 \end{cases}$$

is

$$\boxed{\mathbf{x}(t) = \Psi(t)\Psi^{-1}(t_0)\mathbf{x}^0.}$$

Remark. The matrix $\Psi(t)$ solves the differential equation $\Psi' = P(t)\Psi$. (Homework)

Remark. It is possible to find a *special fundamental matrix*, $\Phi(t)$, which satisfies

$$\Phi(t_0) = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} = I.$$

We will use Φ for this special fundamental matrix, and Ψ for any fundamental matrix.

Example 7.12. Consider

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Find the special fundamental matrix which satisfies $\Phi(0) = I$.

To find the matrix Φ which satisfies

$$\Phi(0) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

we must solve two IVPs:

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \end{cases} \quad \text{and} \quad \begin{cases} \mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x} \\ \mathbf{x}(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{cases}.$$

The general solution to the ODE is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-t}.$$

We calculate that

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} = \mathbf{x}(0) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} \implies \begin{matrix} c_1 = \frac{1}{2} \\ c_2 = \frac{1}{2} \end{matrix} \implies \mathbf{x}(t) = \begin{bmatrix} \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} \\ e^{3t} - e^{-t} \end{bmatrix}$$

and

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \mathbf{x}(0) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} \implies \begin{matrix} c_1 = \frac{1}{4} \\ c_2 = -\frac{1}{4} \end{matrix} \implies \mathbf{x}(t) = \begin{bmatrix} \frac{1}{4}e^{3t} - \frac{1}{4}e^{-t} \\ \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} \end{bmatrix}.$$

Therefore the special fundamental matrix is

$$\Phi(t) = \begin{bmatrix} \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} & \frac{1}{4}e^{3t} - \frac{1}{4}e^{-t} \\ e^{3t} - e^{-t} & \frac{1}{2}e^{3t} + \frac{1}{2}e^{-t} \end{bmatrix}.$$

What is e^{At} ?

Recall that the solution to

$$\begin{cases} x' = ax & (a \in \mathbb{R}) \\ x(0) = x_0 \end{cases}$$

is

$$x(t) = x_0 e^{at} = x_0 \exp(at)$$

and recall that

$$\exp(at) = 1 + \sum_{n=1}^{\infty} \frac{a^n t^n}{n!}.$$

Now consider

$$\begin{cases} \mathbf{x}' = A\mathbf{x} \\ \mathbf{x}(0) = \mathbf{x}^0 \end{cases}$$

for $A \in \mathbb{R}^{n \times n}$.

Definition.

$$\exp(At) = I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} = I + At + \frac{A^2 t^2}{2!} + \frac{A^3 t^3}{3!} + \dots$$

Note that

$$\begin{aligned} \frac{d}{dt} \exp(At) &= \frac{d}{dt} \left(I + \sum_{n=1}^{\infty} \frac{A^n t^n}{n!} \right) = 0 + \sum_{n=1}^{\infty} \frac{d}{dt} \left(\frac{A^n t^n}{n!} \right) \\ &= \sum_{n=1}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} = A + \sum_{n=2}^{\infty} \frac{A^n t^{n-1}}{(n-1)!} \\ &= A + \sum_{k=1}^{\infty} \frac{A^{k+1} t^k}{(k)!} \quad (k = n-1) \\ &= A \left(I + \sum_{k=1}^{\infty} \frac{A^k t^k}{(k)!} \right) = A \exp(At). \end{aligned}$$

This means that $\exp(At)$ solves

$$\begin{cases} (\exp(At))' = A \exp(At) \\ \exp(At)|_{t=0} = I. \end{cases}$$

But remember that Φ solves

$$\begin{cases} \Phi' = A\Phi \\ \Phi(0) = I. \end{cases}$$

Therefore

$$\boxed{\Phi(t) = \exp(At).}$$

Diagonalisable Matrices

If

$$D = \begin{bmatrix} r_1 & 0 & 0 & \dots & 0 \\ 0 & r_2 & 0 & \dots & 0 \\ 0 & 0 & r_3 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & r_n \end{bmatrix}$$

is a diagonal matrix, then it is easy to calculate $\exp(Dt)$. We simply have

$$\exp(Dt) = \begin{bmatrix} e^{r_1 t} & 0 & 0 & \dots & 0 \\ 0 & e^{r_2 t} & 0 & \dots & 0 \\ 0 & 0 & e^{r_3 t} & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & e^{r_n t} \end{bmatrix}.$$

Now consider

$$\mathbf{x}' = A\mathbf{x}$$

for $A \in \mathbb{R}^{n \times n}$. Recall how we diagonalise a matrix: If $\boldsymbol{\xi}^{(1)}, \boldsymbol{\xi}^{(2)}, \dots, \boldsymbol{\xi}^{(n)}$ are the eigenvectors of A , we let

$$T = \begin{bmatrix} \boldsymbol{\xi}^{(1)} & \boldsymbol{\xi}^{(2)} & \dots & \boldsymbol{\xi}^{(n)} \end{bmatrix}.$$

Then

$$\det(T) \neq 0 \implies T^{-1} \text{ exists} \implies \begin{matrix} T^{-1}AT \\ \text{is diagonal} \end{matrix} \implies \begin{matrix} A \text{ is} \\ \text{diagonalisable.} \end{matrix}$$

Example 7.13. Diagonalise

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}.$$

The eigenvalues are $r_1 = 3$ and $r_2 = -1$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$. Thus

$$T = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix} \quad \text{and} \quad T^{-1} = \begin{bmatrix} \frac{1}{2} & \frac{1}{4} \\ \frac{1}{2} & -\frac{1}{4} \end{bmatrix}.$$

It follows that

$$D = T^{-1}AT = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix}.$$

Now consider

$$\mathbf{x}' = A\mathbf{x}.$$

Define a new variable \mathbf{y} by

$$\mathbf{x} = T\mathbf{y} \quad \text{or} \quad \mathbf{y} = T^{-1}\mathbf{x}.$$

Then we calculate that

$$\begin{aligned} \mathbf{x}' &= A\mathbf{x} \\ T\mathbf{y}' &= AT\mathbf{y} \\ \mathbf{y}' &= T^{-1}AT\mathbf{y} = D\mathbf{y}. \end{aligned}$$

We know that a fundamental matrix for $\mathbf{y}' = D\mathbf{y}$ is

$$\exp(Dt) = \begin{bmatrix} e^{r_1 t} & 0 & \dots & 0 \\ 0 & e^{r_2 t} & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & e^{r_n t} \end{bmatrix}.$$

Therefore a fundamental matrix for $\mathbf{x}' = A\mathbf{x}$ is

$$\Psi = T \exp(Dt) = \begin{bmatrix} \boldsymbol{\xi}^{(1)} e^{r_1 t} & \boldsymbol{\xi}^{(2)} e^{r_2 t} & \dots & \boldsymbol{\xi}^{(n)} e^{r_n t} \end{bmatrix}.$$

Example 7.14. Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \mathbf{x}.$$

Recall that $T = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix}$. Letting $\mathbf{y} = T^{-1}\mathbf{x}$, we have

$$\mathbf{y}' = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{y}.$$

A fundamental matrix of $\mathbf{y}' = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{y}$ is

$$\exp(Dt) = e^{Dt} = \begin{bmatrix} e^{3t} & 0 \\ 0 & e^{-t} \end{bmatrix}.$$

Hence

$$\Psi(t) = T \exp(Dt) = \begin{bmatrix} 1 & 1 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} e^{3t} & 0 \\ 0 & e^{-t} \end{bmatrix} = \begin{bmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{bmatrix}.$$

7.8 Repeated Eigenvalues

Example 7.15. Find the eigenvalues and eigenvectors of $A = \begin{bmatrix} 1 & -1 \\ 1 & 3 \end{bmatrix}$.

We calculate that

$$0 = \det(A - rI) = \begin{vmatrix} 1-r & -1 \\ 1 & 3-r \end{vmatrix} = r^2 - 4r + 4 = (r-2)^2.$$

Therefore $r_1 = 2 = r_2$. Moreover

$$\mathbf{0} = (A - rI)\boldsymbol{\xi} = \begin{bmatrix} -1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \implies \xi_1 + \xi_2 = 0 \implies \boldsymbol{\xi} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Note that A has only one linearly independent eigenvector.

Example 7.16. Solve

$$\mathbf{x}' = \begin{bmatrix} 1 & -1 \\ 1 & 3 \end{bmatrix} \mathbf{x}.$$

We know that

$$\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{2t}$$

is a solution. But we need two solutions.

Guess 1: I guess that

$$\mathbf{x}^{(2)}(t) = \boldsymbol{\xi} t e^{2t}$$

for some $\boldsymbol{\xi} \in \mathbb{R}^2$. Then we have

$$\begin{aligned} \boldsymbol{\xi} e^{2t} + 2\boldsymbol{\xi} t e^{2t} &= \mathbf{x}^{(2)'} = A\mathbf{x}^{(2)} = A\boldsymbol{\xi} t e^{2t} \\ \boldsymbol{\xi} + (2\boldsymbol{\xi} - A\boldsymbol{\xi})t &= \mathbf{0} \quad \forall t \\ \implies \boldsymbol{\xi} &= \mathbf{0}. \end{aligned}$$

This guess did not work.

Guess 2: Now I guess that

$$\mathbf{x}^{(2)}(t) = \boldsymbol{\xi} t e^{2t} + \boldsymbol{\eta} e^{2t}$$

for some $\boldsymbol{\xi}, \boldsymbol{\eta} \in \mathbb{R}^2$. Then we have

$$\boldsymbol{\xi} e^{2t} + 2\boldsymbol{\xi} t e^{2t} + 2\boldsymbol{\eta} e^{2t} = \mathbf{x}^{(2)'} = A\mathbf{x}^{(2)} = A(\boldsymbol{\xi} t e^{2t} + \boldsymbol{\eta} e^{2t})$$

and

$$(2\boldsymbol{\xi} - A\boldsymbol{\xi})t + (\boldsymbol{\xi} + 2\boldsymbol{\eta} - A\boldsymbol{\eta}) = \mathbf{0}.$$

Since this must be true $\forall t$, we must have

$$(A - 2I)\boldsymbol{\xi} = \mathbf{0} \quad \text{and} \quad (A - 2I)\boldsymbol{\eta} = \boldsymbol{\xi}.$$

Clearly $\boldsymbol{\xi} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$. Then we calculate that

$$\begin{aligned} \begin{bmatrix} -1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix} &= \begin{bmatrix} 1 \\ -1 \end{bmatrix} \implies \eta_1 + \eta_2 = -1 \\ \implies \boldsymbol{\eta} &= \begin{bmatrix} k \\ -1-k \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \end{bmatrix} + k \begin{bmatrix} 1 \\ -1 \end{bmatrix} \end{aligned}$$

for some k . So

$$\begin{aligned}\mathbf{x}^{(2)}(t) &= \boldsymbol{\xi}te^{2t} + \boldsymbol{\eta}e^{2t} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} te^{2t} + \begin{bmatrix} 0 \\ -1 \end{bmatrix} e^{2t} + k \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{2t} \\ &= \begin{bmatrix} 1 \\ -1 \end{bmatrix} te^{2t} + \begin{bmatrix} 0 \\ -1 \end{bmatrix} e^{2t} + k\mathbf{x}^{(1)}(t).\end{aligned}$$

Because we already have $\mathbf{x}^{(1)}(t)$, we can choose $k = 0$. So

$$\mathbf{x}^{(2)}(t) = \boldsymbol{\xi}te^{2t} + \boldsymbol{\eta}e^{2t} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} te^{2t} + \begin{bmatrix} 0 \\ -1 \end{bmatrix} e^{2t}.$$

The general solution of $\mathbf{x}' = \begin{bmatrix} 1 & -1 \\ 1 & 3 \end{bmatrix} \mathbf{x}$ is therefore

$$\mathbf{x}(t) = c_1 \mathbf{x}^{(1)} + c_2 \mathbf{x}^{(2)} = c_1 \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{2t} + c_2 \left(\begin{bmatrix} 1 \\ -1 \end{bmatrix} te^{2t} + \begin{bmatrix} 0 \\ -1 \end{bmatrix} e^{2t} \right).$$

Example 7.17. Find a fundamental matrix for

$$\mathbf{x}' = \begin{bmatrix} 1 & -1 \\ 1 & 3 \end{bmatrix} \mathbf{x}.$$

Then find the special fundamental matrix $\Phi(t)$ which satisfies $\Phi(0) = I$.

Since $\mathbf{x}^{(1)}(t) = \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{2t}$ and $\mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ -1 \end{bmatrix} te^{2t} + \begin{bmatrix} 0 \\ -1 \end{bmatrix} e^{2t}$ we have that

$$\Psi(t) = \begin{bmatrix} \mathbf{x}^{(1)} & \mathbf{x}^{(2)} \end{bmatrix} = \begin{bmatrix} e^{2t} & te^{2t} \\ -e^{2t} & -te^{2t} - e^{2t} \end{bmatrix}$$

is a fundamental matrix for this system.

Now

$$\Psi(0) = \begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix} \quad \text{and} \quad \Psi^{-1}(0) = \begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix}.$$

Therefore

$$\begin{aligned}\exp(At) &= \Phi(t) = \Psi(t)\Psi^{-1}(0) = e^{2t} \begin{bmatrix} 1 & t \\ -1 & -1-t \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix} \\ &= e^{2t} \begin{bmatrix} 1-t & -t \\ t & 1+t \end{bmatrix}.\end{aligned}$$

Remark.

$$\mathbf{x}' = A\mathbf{x}$$

For two repeated eigenvalues (but with only one linearly independent eigenvector), the key equations to remember are

$$\boxed{\mathbf{x}^{(2)}(t) = \boldsymbol{\xi}te^{rt} + \boldsymbol{\eta}e^{rt}} \quad \text{and} \quad \boxed{(A - rI)\boldsymbol{\eta} = \boldsymbol{\xi}}.$$

Definition. $\boldsymbol{\eta}$ is called a *generalised eigenvector* of A .

Remark. If you have 2 repeated eigenvalues (but with only one linearly independent eigenvector), the method is:

- (i). Find the eigenvalues and eigenvectors;
- (ii). The first solution is $\mathbf{x}^{(1)}(t) = \boldsymbol{\xi}e^{rt}$;
- (iii). Use $(A - rI)\boldsymbol{\eta} = \boldsymbol{\xi}$ to find a generalised eigenvector $\boldsymbol{\eta}$;
- (iv). The second solution is $\mathbf{x}^{(2)}(t) = \boldsymbol{\xi}te^{rt} + \boldsymbol{\eta}e^{rt}$.

Example 7.18. Solve

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} -\frac{5}{2} & \frac{3}{2} \\ -\frac{3}{2} & \frac{1}{2} \end{bmatrix} \mathbf{x}, \\ \mathbf{x}(0) = \begin{bmatrix} 3 \\ -1 \end{bmatrix}. \end{cases}$$

The only eigenvalue of the matrix is $r = -1$. The corresponding eigenvector is $\boldsymbol{\xi} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Therefore one solution of the linear system is

$$\mathbf{x}^{(1)}(t) = \boldsymbol{\xi}e^{rt} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-t}.$$

We need to find a second, linearly independent solution. We will consider the ansatz

$$\mathbf{x}^{(2)}(t) = \boldsymbol{\xi}te^{-t} + \boldsymbol{\eta}e^{-t}$$

where $\boldsymbol{\xi} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ as above and $\boldsymbol{\eta}$ is a generalised eigenvector solving $(A - rI)\boldsymbol{\eta} = \boldsymbol{\xi}$.

Solving the latter equation,

$$\begin{aligned} (A - rI)\boldsymbol{\eta} &= \boldsymbol{\xi} \\ \begin{bmatrix} -\frac{3}{2} & \frac{3}{2} \\ -\frac{3}{2} & \frac{3}{2} \end{bmatrix} \begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix} &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} \\ -\frac{3}{2}\eta_1 + \frac{3}{2}\eta_2 &= 1 \\ -\eta_1 + \eta_2 &= \frac{2}{3} \end{aligned}$$

we can choose $\boldsymbol{\eta} = \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}$.

Note that we don't need to find *every* generalised eigenvector

$$\boldsymbol{\eta} = \begin{bmatrix} k \\ k + \frac{2}{3} \end{bmatrix} = k \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix} = k\boldsymbol{\xi} + \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}$$

because we already have $\mathbf{x}^{(1)}(t) = \boldsymbol{\xi}e^{rt}$.

Instead we only need to find *one* generalised eigenvector – that means that we can choose any k that we want.

Hence I have chosen $k = 0$ which gives $\boldsymbol{\eta} = \begin{bmatrix} 0 \\ \frac{2}{3} \\ 1 \end{bmatrix}$.

Thus

$$\mathbf{x}^{(2)}(t) = \boldsymbol{\xi}te^{-t} + \boldsymbol{\eta}e^{-t} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} + \begin{bmatrix} 0 \\ \frac{2}{3} \\ 1 \end{bmatrix} e^{-t}.$$

Hence the general solution to the linear system is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-t} + c_2 \left(\begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} + \begin{bmatrix} 0 \\ \frac{2}{3} \\ 1 \end{bmatrix} e^{-t} \right).$$

The initial condition gives

$$\begin{bmatrix} 3 \\ -1 \end{bmatrix} = \mathbf{x}(0) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ \frac{2}{3} \\ 1 \end{bmatrix}$$

which implies that $c_1 = 3$ and $c_2 = -6$.

Therefore the solution to the IVP is

$$\boxed{\mathbf{x}(t) = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-t} - 6 \left(\begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} + \begin{bmatrix} 0 \\ \frac{2}{3} \\ 1 \end{bmatrix} e^{-t} \right) = \begin{bmatrix} 3 \\ -1 \end{bmatrix} e^{-t} - 6 \begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t}.$$

Example 7.19. Solve

$$\mathbf{x}' = \begin{bmatrix} 1 & -4 \\ 4 & -7 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 3 \\ 2 \end{bmatrix}.$$

The only eigenvalue of the matrix is $r = -3$. The corresponding eigenvector is $\boldsymbol{\xi} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Therefore one solution of the linear system is

$$\mathbf{x}^{(1)}(t) = \boldsymbol{\xi} e^{rt} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-3t}.$$

Next we need to find a generalised eigenvector $\boldsymbol{\eta}$.

We calculate that

$$\begin{aligned} (A - rI)\boldsymbol{\eta} &= \boldsymbol{\xi} \\ \begin{bmatrix} 4 & -4 \\ 4 & -4 \end{bmatrix} \begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix} &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} \\ 4\eta_1 - 4\eta_2 &= 1 \\ -\eta_1 + \eta_2 &= -\frac{1}{4} \\ \eta_2 &= \eta_1 - \frac{1}{4}. \end{aligned}$$

So we can choose any vector $\boldsymbol{\eta}$ that satisfies $\eta_2 = \eta_1 - \frac{1}{4}$. Thus we may choose $\boldsymbol{\eta} = \begin{bmatrix} 0 \\ -\frac{1}{4} \end{bmatrix}$.

Therefore

$$\mathbf{x}^{(2)}(t) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} t e^{-3t} + \begin{bmatrix} 0 \\ -\frac{1}{4} \end{bmatrix} e^{-3t}.$$

Hence the general solution to the ODE is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-3t} + c_2 \left(\begin{bmatrix} 1 \\ 1 \end{bmatrix} t e^{-3t} + \begin{bmatrix} 0 \\ -\frac{1}{4} \end{bmatrix} e^{-3t} \right).$$

The initial condition gives

$$\begin{bmatrix} 3 \\ 2 \end{bmatrix} = \mathbf{x}(0) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ -\frac{1}{4} \end{bmatrix}$$

which implies that $c_1 = 3$ and $c_2 = 4$.

Therefore the solution to the IVP is

$$\begin{aligned} \mathbf{x}(t) &= 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-3t} + 4 \left(\begin{bmatrix} 1 \\ 1 \end{bmatrix} t e^{-3t} + \begin{bmatrix} 0 \\ -\frac{1}{4} \end{bmatrix} e^{-3t} \right) \\ &= \begin{bmatrix} 3 \\ 2 \end{bmatrix} e^{-3t} + 4 \begin{bmatrix} 1 \\ 1 \end{bmatrix} t e^{-3t} \\ &= \begin{bmatrix} 3 + 4t \\ 2 + 4t \end{bmatrix} e^{-3t}. \end{aligned}$$

7.9 Nonhomogeneous Linear Systems

Consider

$$\mathbf{x}' = P(t)\mathbf{x} + \mathbf{g}(t) \quad (7.1)$$

where $P(t)$ and $\mathbf{g}(t)$ are continuous for $\alpha < t < \beta$. The general solution of (7.1) can be written as

$$\mathbf{x}(t) = c_1\mathbf{x}^{(1)} + c_2\mathbf{x}^{(2)} + \dots + c_n\mathbf{x}^{(n)} + \mathbf{v}(t)$$

where

- $c_1\mathbf{x}^{(1)} + c_2\mathbf{x}^{(2)} + \dots + c_n\mathbf{x}^{(n)}$ is the general solution to the homogeneous system $\mathbf{x}' = P(t)\mathbf{x}$; and
- $\mathbf{v}(t)$ is a particular solution to (7.1).

Remark. We will study four methods to solve (7.1):

- (i). Diagonalisation;
- (ii). Undetermined Coefficients;
- (iii). Variation of Parameters;
- (iv). The Laplace Transform.

Method 1 – Diagonalisation:

Consider

$$\mathbf{x}' = A\mathbf{x} + \mathbf{g}(t).$$

Suppose that

- $A \in \mathbb{R}^{n \times n}$ is diagonalisable;
- $\mathbf{g} : (\alpha, \beta) \rightarrow \mathbb{R}^n$;
- $\boldsymbol{\xi}^{(1)}, \dots, \boldsymbol{\xi}^{(n)}$ are eigenvectors of A ; and
- $T = \begin{bmatrix} \boldsymbol{\xi}^{(1)} & \dots & \boldsymbol{\xi}^{(n)} \end{bmatrix}$.

Then

$$D = T^{-1}AT = \begin{bmatrix} r_1 & 0 & \dots & 0 \\ 0 & r_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & r_n \end{bmatrix}.$$

Let $\mathbf{y} = T^{-1}\mathbf{x}$. Then $\mathbf{x} = T\mathbf{y}$. It follows that

$$T\mathbf{y}' = \mathbf{x}' = A\mathbf{x} + \mathbf{g}(t) = AT\mathbf{y} + \mathbf{g}(t)$$

and

$$\mathbf{y}' = T^{-1}AT\mathbf{y} + T^{-1}\mathbf{g}(t) = D\mathbf{y} + \mathbf{h}(t) \quad (7.2)$$

where $\mathbf{h} = T^{-1}\mathbf{g}$.

But (7.2) is just the system

$$\begin{cases} y_1' = r_1 y_1 + h_1(t) & \leftarrow \text{only } y_1 \text{ and } t \\ y_2' = r_2 y_2 + h_2(t) & \leftarrow \text{only } y_2 \text{ and } t \\ \vdots \\ y_n' = r_n y_n + h_n(t) & \leftarrow \text{only } y_n \text{ and } t \end{cases}$$

We can solve each of these n first order linear ODEs individually. The solution to

$$y_j' - r_j y_j = h_j$$

(see Chapter 2) is

$$y_j(t) = e^{r_j t} \int_{t_0}^t e^{-r_j s} h(s) ds + c_j e^{r_j t}.$$

If we know \mathbf{y} , then we know $\mathbf{x} = T\mathbf{y}$.

Example 7.20. Solve

$$\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} = A\mathbf{x} + \mathbf{g}(t).$$

The eigenvalues of $A = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix}$ are $r_1 = -3$ and $r_2 = -1$. The eigenvectors are

$$\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \quad \text{and} \quad \boldsymbol{\xi}^{(2)} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

So

$$T = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \quad \text{and} \quad T^{-1} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}.$$

Let $\mathbf{y} = T^{-1}\mathbf{x}$. Then

$$\begin{aligned} T\mathbf{y}' &= \mathbf{x}' = A\mathbf{x} + \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} = AT\mathbf{y} + \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} \\ \mathbf{y}' &= T^{-1}AT\mathbf{y} + T^{-1} \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} \\ &= D\mathbf{y} + \frac{1}{2} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} \\ &= \begin{bmatrix} -3 & 0 \\ 0 & -1 \end{bmatrix} \mathbf{y} + \frac{1}{2} \begin{bmatrix} 2e^{-t} - 3t \\ 2e^{-t} + 3t \end{bmatrix}. \end{aligned}$$

Therefore

$$\begin{cases} y_1' + 3y_1 = e^{-t} - \frac{3}{2}t \\ y_2' + y_2 = e^{-t} + \frac{3}{2}t. \end{cases}$$

You know how to solve first order linear ODEs. The solutions to these two ODEs are

$$\begin{aligned} y_1(t) &= \frac{1}{2}e^{-t} - \frac{t}{2} + \frac{1}{6} + c_1e^{-3t} \\ y_2(t) &= te^{-t} + \frac{3t}{2} - \frac{3}{2} + c_2e^{-t}. \end{aligned}$$

Finally we calculate that

$$\begin{aligned} \mathbf{x} &= T\mathbf{y} = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \\ &= \begin{bmatrix} y_1 + y_2 \\ -y_1 + y_2 \end{bmatrix} \\ &= c_1 \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-3t} + c_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-t} + \frac{1}{2} \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-t} + \begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} t - \frac{1}{3} \begin{bmatrix} 4 \\ 5 \end{bmatrix}. \end{aligned}$$

Example 7.21. Solve

$$\mathbf{x}' = \begin{bmatrix} 1 & \sqrt{3} \\ \sqrt{3} & -1 \end{bmatrix} \mathbf{x} + \begin{bmatrix} e^t \\ \sqrt{3}e^{-t} \end{bmatrix}.$$

The eigenvalues of $\begin{bmatrix} 1 & \sqrt{3} \\ \sqrt{3} & -1 \end{bmatrix}$ are $r_1 = -2$ and $r_2 = 2$. The corresponding eigenvectors are $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ -\sqrt{3} \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} \sqrt{3} \\ 1 \end{bmatrix}$.

Thus

$$T = \begin{bmatrix} 1 & \sqrt{3} \\ -\sqrt{3} & 1 \end{bmatrix}$$

and

$$T^{-1} = \frac{1}{\det T} \begin{bmatrix} 1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{4} & -\frac{\sqrt{3}}{4} \\ \frac{\sqrt{3}}{4} & \frac{1}{4} \end{bmatrix}.$$

Now we must change variables: Let $\mathbf{y} = T^{-1}\mathbf{x}$. Then we have

$$\begin{aligned} \mathbf{y}' &= D\mathbf{y} + T^{-1}\mathbf{g} = \begin{bmatrix} -2 & 0 \\ 0 & 2 \end{bmatrix} \mathbf{y} + \begin{bmatrix} \frac{1}{4} & -\frac{\sqrt{3}}{4} \\ \frac{\sqrt{3}}{4} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} e^t \\ \sqrt{3}e^{-t} \end{bmatrix} \\ &= \begin{bmatrix} -2y_1 \\ 2y_2 \end{bmatrix} + \begin{bmatrix} \frac{1}{4}e^t - \frac{3}{4}e^{-t} \\ \frac{\sqrt{3}}{4}e^t + \frac{\sqrt{3}}{4}e^{-t} \end{bmatrix}. \end{aligned}$$

We know how to solve

$$y_1' + 2y_1 = \frac{1}{4}e^t - \frac{3}{4}e^{-t}$$

and

$$y_2' - 2y_2 = \frac{\sqrt{3}}{4}e^t + \frac{\sqrt{3}}{4}e^{-t}.$$

The solutions are

$$y_1(t) = \frac{1}{12}e^t - \frac{3}{4}e^{-t} + c_1e^{-2t}$$

and

$$y_2(t) = -\frac{\sqrt{3}}{4}e^t - \frac{\sqrt{3}}{12}e^{-t} + c_2e^{2t}.$$

So

$$\mathbf{y} = \begin{bmatrix} \frac{1}{12}e^t - \frac{3}{4}e^{-t} + c_1e^{-2t} \\ -\frac{\sqrt{3}}{4}e^t - \frac{\sqrt{3}}{12}e^{-t} + c_2e^{2t} \end{bmatrix}.$$

Therefore the general solution to the ODE is

$$\mathbf{x} = T\mathbf{y} = \begin{bmatrix} 1 & \sqrt{3} \\ -\sqrt{3} & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{12}e^t - \frac{3}{4}e^{-t} + c_1e^{-2t} \\ -\frac{\sqrt{3}}{4}e^t - \frac{\sqrt{3}}{12}e^{-t} + c_2e^{2t} \end{bmatrix} = \dots$$

Method 2 – Undetermined Coefficients:

Consider

$$\mathbf{x}' = A\mathbf{x} + \mathbf{g}(t).$$

(Remember Chapter 3?)

The idea is

- (i). Find the general solution to $\mathbf{x}' = A\mathbf{x}$.
- (ii). Look at $\mathbf{g}(t)$. Make a guess with constants. Find the constants.
- (iii). 1 + 2.

Example 7.22. Solve

$$\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} = A\mathbf{x} + \mathbf{g}(t).$$

1. The solution of $\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x}$ is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-t} + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-3t}.$$

2. Since $\mathbf{g}(t) = \begin{bmatrix} 2 \\ 0 \end{bmatrix} e^{-t} + \begin{bmatrix} 0 \\ 3 \end{bmatrix} t$, we try the ansatz

$$\mathbf{x} = \mathbf{v}(t) = \mathbf{a}te^{-t} + \mathbf{b}e^{-t} + \mathbf{c}t + \mathbf{d}.$$

(Note that because $r_1 = -1$ is an eigenvalue of $\begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix}$, we need both te^{-t} and e^{-t} .)

Then we calculate that

$$\mathbf{x}' = A\mathbf{x} + \mathbf{g}$$

$$\mathbf{a}e^{-t} - \mathbf{a}te^{-t} - \mathbf{b}e^{-t} + \mathbf{c} = A\mathbf{a}te^{-t} + A\mathbf{b}e^{-t} + A\mathbf{c}t + A\mathbf{d} + \begin{bmatrix} 2 \\ 0 \end{bmatrix} e^{-t} + \begin{bmatrix} 0 \\ 3 \end{bmatrix} t.$$

- If we look at the te^{-t} terms, we have

$$-\mathbf{a} = A\mathbf{a} \implies \mathbf{a} \text{ is an eigenvector} \implies \mathbf{a} = \begin{bmatrix} \alpha \\ \alpha \end{bmatrix} \text{ for some } \alpha \in \mathbb{R}.$$

- If we look at the e^{-t} terms, we have

$$\mathbf{a} - \mathbf{b} = A\mathbf{b} + \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

which becomes

$$\begin{bmatrix} \alpha - 2 \\ \alpha \end{bmatrix} = \mathbf{a} - \begin{bmatrix} 2 \\ 0 \end{bmatrix} = (A + I)\mathbf{b} = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \begin{bmatrix} -b_1 + b_2 \\ b_1 - b_2 \end{bmatrix}.$$

But this means that

$$\alpha - 2 = -b_1 + b_2 = -(b_1 - b_2) = -\alpha \implies \alpha = 1.$$

So $\mathbf{a} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Then we have that

$$b_1 - b_2 = 1 \quad \implies \quad \mathbf{b} = \begin{bmatrix} k \\ k - 1 \end{bmatrix}$$

for any k . If we choose $k = 0$, we get $\mathbf{b} = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$.

- If we look at the t terms, we have

$$0 = A\mathbf{c} + \begin{bmatrix} 0 \\ 3 \end{bmatrix} \quad \implies \quad \mathbf{c} = A^{-1} \begin{bmatrix} 0 \\ -3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

- Finally, if we look at the 1 terms, we have

$$\mathbf{c} = A\mathbf{d} \quad \implies \quad \mathbf{d} = A^{-1}\mathbf{c} = \begin{bmatrix} -\frac{4}{3} \\ \frac{5}{3} \end{bmatrix}.$$

So

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} - \begin{bmatrix} 0 \\ 1 \end{bmatrix} e^{-t} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} t - \frac{1}{3} \begin{bmatrix} 4 \\ 5 \end{bmatrix}.$$

3. Therefore the general solution to the IVP is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-t} + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-3t} + \begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} - \begin{bmatrix} 0 \\ 1 \end{bmatrix} e^{-t} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} t - \frac{1}{3} \begin{bmatrix} 4 \\ 5 \end{bmatrix}.$$

Example 7.23. Solve

$$\mathbf{x}' = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \mathbf{x} + \begin{bmatrix} e^t \\ -10t - 3 \end{bmatrix}.$$

The matrix $\begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}$ has eigenvalues $r_1 = 5$ and $r_2 = -2$ and eigenvectors $\boldsymbol{\xi}^{(1)} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\boldsymbol{\xi}^{(2)} = \begin{bmatrix} -3 \\ 4 \end{bmatrix}$. Hence the general solution of $\mathbf{x}' = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \mathbf{x}$ is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{5t} + c_2 \begin{bmatrix} -3 \\ 4 \end{bmatrix} e^{-2t}.$$

Next we need to find a particular solution to $\mathbf{x}' = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \mathbf{x} + \begin{bmatrix} e^t \\ 0 \end{bmatrix}$. Since 1 is not an eigenvalue of $\begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}$, we try the ansatz $\mathbf{x} = \mathbf{a}e^t$ for some $\mathbf{a} = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} \in \mathbb{R}^2$. Then we calculate that

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} e^t = \mathbf{x}' = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \mathbf{x} + \begin{bmatrix} e^t \\ 0 \end{bmatrix} = \begin{bmatrix} 2a_1 + 3a_2 + 1 \\ 4a_1 + a_2 \end{bmatrix} e^t$$

which gives

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 2a_1 + 3a_2 + 1 \\ 4a_1 + a_2 \end{bmatrix}.$$

Hence $a_1 = 0$ and $a_2 = -\frac{1}{3}$. So $\mathbf{x} = \begin{bmatrix} 0 \\ -\frac{1}{3} \end{bmatrix} e^t$.

Then we need to find a particular solution to $\mathbf{x}' = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 0 \\ -10t - 3 \end{bmatrix}$. We try the ansatz $\mathbf{x} = \mathbf{a}t + \mathbf{b} = \begin{bmatrix} a_1t + b_1 \\ a_2t + b_2 \end{bmatrix}$ for some $\mathbf{a}, \mathbf{b} \in \mathbb{R}^2$ and calculate that

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = \mathbf{x}' = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 0 \\ t \end{bmatrix} = \begin{bmatrix} 2a_1t + 2b_1 + 3a_2t + 3b_2 \\ 4a_1t + 4b_1 + a_2t + b_2 - 10t - 3 \end{bmatrix}$$

which leads to

$$\begin{cases} 0 = 2a_1 + 3a_2 \\ a_1 = 2b_1 + 3b_2 \\ 0 = 4a_1 + a_2 - 10t \\ a_2 = 4b_1 + b_2 - 3. \end{cases}$$

The solution to this linear system is $\mathbf{a} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Hence $\mathbf{x} = \begin{bmatrix} 3t \\ 1 - 2t \end{bmatrix}$.

Adding all of these together, we find that the general solution to the given ODE is

$$\mathbf{x} = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{5t} + c_2 \begin{bmatrix} -3 \\ 4 \end{bmatrix} e^{-2t} + \begin{bmatrix} 0 \\ -\frac{1}{3} \end{bmatrix} e^t + \begin{bmatrix} 3t \\ 1 - 2t \end{bmatrix}.$$

Method 3 – Variation of Parameters:

Consider

$$\mathbf{x}' = P(t)\mathbf{x} + \mathbf{g}(t) \quad (7.1)$$

where

- P and \mathbf{g} are continuous for $\alpha < t < \beta$;
- there exists a fundamental matrix $\Psi(t)$ for the homogeneous system $\mathbf{x}' = P(t)\mathbf{x}$.

We know that the general solution to $\mathbf{x}' = P(t)\mathbf{x}$ is $\mathbf{x} = \Psi(t)\mathbf{c}$.

We guess that

$$\mathbf{x} = \Psi(t)\mathbf{u}(t)$$

is a solution to (7.1). Can we find $\mathbf{u}(t)$?

If $\mathbf{x} = \Psi\mathbf{u}$, we can calculate that

$$\cancel{\Psi'}\mathbf{u} + \Psi\mathbf{u}' = \mathbf{x}' = P\mathbf{x} + \mathbf{g} = P\Psi\mathbf{u} + \mathbf{g}. \quad (7.3)$$

But remember that

$$\Psi \text{ is a fundamental matrix for } \mathbf{x}' = P(t)\mathbf{x} \implies \Psi \text{ solves } \Psi' = P\Psi.$$

Hence (7.3) becomes

$$\Psi\mathbf{u}' = \mathbf{g}.$$

Therefore

$$\mathbf{u}' = \Psi^{-1}\mathbf{g}$$

and

$$\mathbf{u} = \int \Psi^{-1}\mathbf{g}.$$

Hence

$$\mathbf{x} = \Psi(t)\mathbf{u}(t) = \Psi(t) \int \Psi^{-1}(s)\mathbf{g}(s) ds.$$

Remark. To solve $\mathbf{x}' = P(t)\mathbf{x} + \mathbf{g}(t)$, the method is

- Find a fundamental matrix Ψ for $\mathbf{x}' = P(t)\mathbf{x}$;
- Calculate $\mathbf{x} = \Psi(t) \int \Psi^{-1}(s)\mathbf{g}(s) ds$.

Example 7.24. Solve

$$\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} = A\mathbf{x} + \mathbf{g}(t).$$

The solution of $\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x}$ is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-3t} + c_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-t}.$$

So

$$\Psi(t) = \begin{bmatrix} e^{-3t} & e^{-t} \\ -e^{-3t} & e^{-t} \end{bmatrix}$$

is a fundamental matrix.

Then we calculate that

$$\Psi^{-1}(t) = \frac{1}{2e^{-4t}} \begin{bmatrix} e^{-t} & -e^{-t} \\ e^{-3t} & e^{-3t} \end{bmatrix} = \frac{1}{2}e^{4t} \begin{bmatrix} e^{-t} & -e^{-t} \\ e^{-3t} & e^{-3t} \end{bmatrix} = \begin{bmatrix} \frac{1}{2}e^{3t} & -\frac{1}{2}e^{3t} \\ \frac{1}{2}e^t & \frac{1}{2}e^t \end{bmatrix}$$

and

$$\begin{aligned} \int \Psi^{-1}(t)\mathbf{g}(t) dt &= \int \begin{bmatrix} \frac{1}{2}e^{3t} & -\frac{1}{2}e^{3t} \\ \frac{1}{2}e^t & \frac{1}{2}e^t \end{bmatrix} \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} dt \\ &= \int \begin{bmatrix} e^{2t} - \frac{3}{2}te^{3t} \\ 1 + \frac{3}{2}te^t \end{bmatrix} dt = \begin{bmatrix} \frac{1}{2}e^{2t} - \frac{1}{2}te^{3t} + \frac{1}{6}e^{3t} + c_1 \\ t + \frac{3}{2}te^t - \frac{3}{2}e^t + c_2 \end{bmatrix}. \end{aligned}$$

Therefore the solution to $\mathbf{x}' = A\mathbf{x} + \mathbf{g}$ is

$$\begin{aligned} \mathbf{x} &= \Psi(t) \int \Psi^{-1}(s)\mathbf{g}(s) ds \\ &= \begin{bmatrix} e^{-3t} & e^{-t} \\ -e^{-3t} & e^{-t} \end{bmatrix} \begin{bmatrix} \frac{1}{2}e^{2t} - \frac{1}{2}te^{3t} + \frac{1}{6}e^{3t} + c_1 \\ t + \frac{3}{2}te^t - \frac{3}{2}e^t + c_2 \end{bmatrix} \\ &= c_1 \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-3t} + c_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-t} + \frac{1}{2} \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-t} + \begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} t - \frac{1}{3} \begin{bmatrix} 4 \\ 5 \end{bmatrix}. \end{aligned}$$

Example 7.25. Solve

$$\mathbf{x}' = \begin{bmatrix} -4 & 2 \\ 2 & -1 \end{bmatrix} \mathbf{x} + \begin{bmatrix} t^{-1} \\ 2t^{-1} + 4 \end{bmatrix}$$

for $t > 0$.

The eigenvalues of $A = \begin{bmatrix} -4 & 2 \\ 2 & -1 \end{bmatrix}$ are $r_1 = 0$ and $r_2 = -5$; and the eigenvectors are $\xi^{(1)} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\xi^{(2)} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$. Thus

$$\Psi(t) = \begin{bmatrix} 1 & -2e^{-5t} \\ 2 & e^{-5t} \end{bmatrix}$$

is a fundamental matrix for $\mathbf{x}' = A\mathbf{x}$.

Using the formula $\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$ we calculate that

$$\Psi^{-1}(t) = \frac{1}{e^{-5t} + 4e^{-5t}} \begin{bmatrix} e^{-5t} & 2e^{-5t} \\ -2 & 1 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 1 & 2 \\ -2e^{5t} & e^{5t} \end{bmatrix}.$$

Then

$$\begin{aligned} \Psi^{-1}(t)\mathbf{g}(t) &= \frac{1}{5} \begin{bmatrix} 1 & 2 \\ -2e^{5t} & e^{5t} \end{bmatrix} \begin{bmatrix} t^{-1} \\ 2t^{-1} + 4 \end{bmatrix} \\ &= \frac{1}{5} \begin{bmatrix} t^{-1} + 4t^{-1} + 8 \\ -2t^{-1}e^{5t} + 2t^{-1}e^{5t} + 4e^{5t} \end{bmatrix} = \begin{bmatrix} t^{-1} + \frac{8}{5} \\ \frac{4}{5}e^{5t} \end{bmatrix} \end{aligned}$$

and

$$\int \Psi^{-1}(t)\mathbf{g}(t) dt = \int \begin{bmatrix} t^{-1} + \frac{8}{5} \\ \frac{4}{5}e^{5t} \end{bmatrix} dt = \begin{bmatrix} \ln t + \frac{8}{5}t + c_1 \\ \frac{4}{25}e^{5t} + c_2 \end{bmatrix}.$$

It follows that

$$\begin{aligned} \mathbf{x}(t) &= \Psi(t) \int \Psi^{-1}(s)\mathbf{g}(s) ds = \begin{bmatrix} 1 & -2e^{-5t} \\ 2 & e^{-5t} \end{bmatrix} \begin{bmatrix} \ln t + \frac{8}{5}t + c_1 \\ \frac{4}{25}e^{5t} + c_2 \end{bmatrix} \\ &= \begin{bmatrix} \ln t + \frac{8}{5}t - \frac{8}{25} + c_1 - 2c_2e^{-5t} \\ 2\ln t + \frac{16}{5}t + \frac{4}{25} + 2c_1 + c_2e^{-5t} \end{bmatrix} \\ &= c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + c_2 \begin{bmatrix} -2 \\ 1 \end{bmatrix} e^{-5t} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} \ln t + \frac{8}{5} \begin{bmatrix} 1 \\ 2 \end{bmatrix} t + \frac{4}{25} \begin{bmatrix} -2 \\ 1 \end{bmatrix}. \end{aligned}$$

Method 4 – The Laplace Transform:

First some notation: If $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$, then $\mathbf{X} = \mathcal{L}[\mathbf{x}] = \begin{bmatrix} \mathcal{L}[x_1] \\ \mathcal{L}[x_2] \\ \vdots \\ \mathcal{L}[x_n] \end{bmatrix}$.

Recall from Chapter 6 that $\mathcal{L}[y']$ satisfies

$$\mathcal{L}[y'](s) = sY(s) - y(0).$$

It follows that:

Theorem 7.3.

$$\mathcal{L}[\mathbf{x}'](s) = s\mathbf{X}(s) - \mathbf{x}(0).$$

Example 7.26. Solve

$$\begin{cases} \mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix} = A\mathbf{x} + \mathbf{g}(t), \\ \mathbf{x}(0) = \mathbf{0}. \end{cases}$$

Taking Laplace Transforms of the ODE gives

$$s\mathbf{X}(s) - \mathbf{x}(0) = A\mathbf{X}(s) + \mathbf{G}(s)$$

where $\mathbf{G}(s) = \mathcal{L}[\mathbf{g}](s) = \begin{bmatrix} \frac{2}{s+1} \\ \frac{3}{s^2} \end{bmatrix}$.

Thus

$$(sI - A)\mathbf{X} = \mathbf{G}$$

and

$$\mathbf{X} = (sI - A)^{-1}\mathbf{G}$$

where

$$(sI - A)^{-1} = \begin{bmatrix} s+2 & -1 \\ -1 & s+2 \end{bmatrix}^{-1} = \frac{1}{(s+1)(s+3)} \begin{bmatrix} s+2 & 1 \\ 1 & s+2 \end{bmatrix}.$$

So

$$\begin{aligned} \mathbf{X} &= (sI - A)^{-1}\mathbf{G} \\ &= \frac{1}{(s+1)(s+3)} \begin{bmatrix} s+2 & 1 \\ 1 & s+2 \end{bmatrix} \begin{bmatrix} \frac{2}{s+1} \\ \frac{3}{s^2} \end{bmatrix} \\ &= \begin{bmatrix} \frac{2(s+2)}{(s+1)^2(s+3)} + \frac{3}{s^2(s+1)(s+3)} \\ \frac{2}{(s+1)^2(s+3)} + \frac{3(s+2)}{s^2(s+1)(s+3)} \end{bmatrix}. \end{aligned}$$

When we take the inverse Laplace Transform of this, we find our solution

$$\mathbf{x} = \mathcal{L}^{-1}[\mathbf{X}] = \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{-t} - \frac{2}{3} \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-3t} + \begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} t - \frac{1}{3} \begin{bmatrix} 4 \\ 5 \end{bmatrix}.$$

Example 7.27. Solve

$$\begin{cases} 2x' + y' - y - t = 0 \\ x' + y' - t^2 = 0 \\ x(0) = 1 \\ y(0) = 0 \end{cases}$$

The ODEs above can be written as

$$\begin{cases} x' = y - t^2 + t \\ y' = -y + 2t^2 - t \end{cases}$$

(please check!).

If we write the problem in terms of matrices (using $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$) we have

$$\begin{cases} \mathbf{x}' = A\mathbf{x} + \mathbf{g} = \begin{bmatrix} 0 & 1 \\ 0 & -1 \end{bmatrix} \mathbf{x} + \begin{bmatrix} t - t^2 \\ 2t^2 - t \end{bmatrix} \\ \mathbf{x}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \end{cases}$$

Taking the Laplace transform of the ODE gives

$$\begin{aligned} (sI - A)\mathbf{X}(s) &= \mathbf{x}(0) + \mathbf{G}(s) \\ \begin{bmatrix} s & -1 \\ 0 & s+1 \end{bmatrix} \mathbf{X}(s) &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \begin{bmatrix} \frac{1}{s^2} - \frac{2}{s^3} \\ \frac{4}{s^3} - \frac{1}{s^2} \end{bmatrix} \end{aligned}$$

Thus

$$\begin{aligned} \mathbf{X}(s) &= \frac{1}{s(s+1)} \begin{bmatrix} s+1 & 1 \\ 0 & s \end{bmatrix} \frac{1}{s^3} \begin{bmatrix} s^3 + s - 2 \\ 4 - s \end{bmatrix} \\ &= \frac{1}{s^4(s+1)} \begin{bmatrix} s^4 + s^3 + s^2 - 2s + 2 \\ 4s - s^2 \end{bmatrix}. \end{aligned}$$

Note that

$$\frac{s^4 + s^3 + s^2 - 2s + 2}{s^4(s+1)} = \frac{5}{s+1} - 4\frac{1}{s} + 5\frac{1}{s^2} - 4\frac{1}{s^3} + 2\frac{1}{s^4}$$

and

$$\frac{4s - s^2}{s^4(s+1)} = -5\frac{1}{s+1} + 5\frac{1}{s} - 5\frac{1}{s^2} + 4\frac{1}{s^3}$$

(please check!).

It follows that

$$\mathcal{L}^{-1} \left(\frac{s^4 + s^3 + s^2 - 2s + 2}{s^4(s+1)} \right) = 5e^{-t} - 4 + 5t - 2t^2 + \frac{1}{3}t^3$$

and

$$\mathcal{L}^{-1} \left(\frac{4s - s^2}{s^4(s+1)} \right) = -5e^{-t} + 5 - 5t + 2t^2.$$

Therefore the solution to the initial value problem is

$$\boxed{\mathbf{x}(t) = \begin{bmatrix} 5e^{-t} - 4 + 5t - 2t^2 + \frac{1}{3}t^3 \\ -5e^{-t} + 5 - 5t + 2t^2 \end{bmatrix}}.$$