

Welcome to

Mathematics IV (Differential Equations) with Dr Neil Course

1 of 50



Lecture 1

- Information about this course
- 1.1 Introduction
- 1.2 Some Examples
- 1.3 How to Draw a Direction Field
- 1.4 Solving Our First Differential Equations



Information about this course

 ≈ 12 classes. Wednesday and Thursday afternoons 2pm-4pm.

14:00 15:00 16:00



- $\blacksquare \approx 12$ classes. We dnesday and Thursday afternoons 2pm-4pm.
- Each lecture ≈ 60 minutes.

lecture

14:00 15:00 16:00



- $\blacksquare \approx 12$ classes. We dnesday and Thursday afternoons 2pm-4pm.
- Each lecture ≈ 60 minutes.
- Then I will answers your questions.

lecture	questions	
14:00	15:00	16:00



- $\blacksquare \approx 12$ classes. We dnesday and Thursday afternoons 2pm-4pm.
- Each lecture ≈ 60 minutes.
- Then I will answers your questions.

lecture	questions	
14:00	15:00	16:00



- $\blacksquare \approx 12$ classes. We dnesday and Thursday afternoons 2pm-4pm.
- Each lecture ≈ 60 minutes.
- Then I will answers your questions.

lecture	questions	
14:00	15:00	16:00



- $\blacksquare \approx 12$ classes. We dnesday and Thursday afternoons 2pm-4pm.
- Each lecture ≈ 60 minutes.
- Then I will answers your questions.





- $\blacksquare \approx 12$ classes. We dnesday and Thursday afternoons 2pm-4pm.
- Each lecture ≈ 60 minutes.
- Then I will answers your questions.

	lecture	questions	
14	:00	15:00	16:00



- $\blacksquare \approx 12$ classes. We dnesday and Thursday afternoons 2pm-4pm.
- Each lecture ≈ 60 minutes.
- Then I will answers your questions.

lecture	questions	
14:00	15:00	16:00



1	
2	
3	
4	
5	



Introduction

Examples; Directions Fields; Classification.

2

3

4



Introduction

Examples; Directions Fields; Classification.

First Order Differential Equations

3

4



Introduction

Examples; Directions Fields; Classification.

First Order Differential Equations

Second and Higher Order Linear ODEs

4



Introduction

Examples; Directions Fields; Classification.

First Order Differential Equations

Second and Higher Order Linear ODEs

The Laplace Transform



Introduction

Examples; Directions Fields; Classification.

First Order Differential Equations

Second and Higher Order Linear ODEs

The Laplace Transform

Systems of First Order Linear ODEs



Introduction

Examples; Directions Fields; Classification.

First Order Differential Equations

Second and Higher Order Linear ODEs

The Laplace Transform

Systems of First Order Linear ODEs

3 lectures


3 lectures

3 lectures

3 lectures



Lecture Notes






Exams and homework

(This information may change based on the University's decisions) $\,$




(This information may change based on the University's decisions)






(This information may change based on the University's decisions)



10 multiple choice tests on O'Learn.




(This information may change based on the University's decisions)





(This information may change based on the University's decisions)





Expectations

If this was a classroom course, the expectation is that for each hour of lectures, you would study an extra 1-2 hours outside of class.

classroom course

lectures (8 hours) other study (8-16 hours)



Expectations

If this was a classroom course, the expectation is that for each hour of lectures, you would study an extra 1-2 hours outside of class.

classroom course lectures (8 hours) other study (8-16 hours)

For an online course, you are still expected to study a total of 16-24 hours each week.

online class (4 hours) other study (12-20 hours)



This may include:

■ Do the online homework tests each week;



This may include:

- Do the online homework tests each week;
- Rewatch the recorded lectures (O'Learn & YouTube);



This may include:

- Do the online homework tests each week;
- Rewatch the recorded lectures (O'Learn & YouTube);
- Read the lecture notes or slides (before the lecture? after the lecture?);

:



This may include:

- Do the online homework tests each week;
- Rewatch the recorded lectures (O'Learn & YouTube);
- Read the lecture notes or slides (before the lecture? after the lecture?);
- Solve the exercises in the lecture notes;



This may include:

- Do the online homework tests each week;
- Rewatch the recorded lectures (O'Learn & YouTube);
- Read the lecture notes or slides (before the lecture? after the lecture?);
- Solve the exercises in the lecture notes;
- Use the O'Learn Discussion Board;

.



This may include:


- Do the online homework tests each week;
- Rewatch the recorded lectures (O'Learn & YouTube);
- Read the lecture notes or slides (before the lecture? after the lecture?);
- Solve the exercises in the lecture notes;
- Use the O'Learn Discussion Board;
- Read books;



This may include:

- Do the online homework tests each week;
- Rewatch the recorded lectures (O'Learn & YouTube);
- Read the lecture notes or slides (before the lecture? after the lecture?);
- Solve the exercises in the lecture notes;
- Use the O'Learn Discussion Board;
- Read books;
- Watch online videos;






Two good books

William E. Boyce, Richard C. DiPrima and Douglas B. Meade,


Boyce's Elementary Differential Equations and Boundary Value Problems, Wilev.



C. Henry Edwards and David E. Penney, Elementary Differential Equations with Boundary Value Problems, Pearson.


9.9 Section Title





9.9 Section Title







Introduction



If I say "Solve x + 3 = 5", I am looking for some number x which satisfies this equation.



If I say "Solve x + 3 = 5", I am looking for some number x which satisfies this equation.

But if I say "Solve $\frac{dy}{dx} = 2x$ ", then I don't want a number; I am looking for a function y(x) which satisfies this equation.



If I say "Solve x + 3 = 5", I am looking for some number x which satisfies this equation.

But if I say "Solve $\frac{dy}{dx} = 2x$ ", then I don't want a number; I am looking for a function y(x) which satisfies this equation.

A differential equation is an equation containing a derivative.



If I say "Solve x + 3 = 5", I am looking for some number x which satisfies this equation.

But if I say "Solve $\frac{dy}{dx} = 2x$ ", then I don't want a number; I am looking for a function y(x) which satisfies this equation.

A differential equation is an equation containing a derivative. For example, the equation $\frac{dy}{dx} = 2x$ contains a derivative – so it is a differential equation.



Example

Solve
$$\frac{dy}{dx} = 2x$$
.



Example

Solve
$$\frac{dy}{dx} = 2x$$
.

This differential equation is easy to solve:

$$y(x) = \int \frac{dy}{dx} dx = \int 2x dx = x^2 + c.$$



${\bf Example}$

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$



Example

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$

This problem has two conditions. The first line says $\frac{dy}{dx} = 2x$ which is a differential equation.



Example

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$

This problem has two conditions. The first line says $\frac{dy}{dx} = 2x$ which is a differential equation. The second line, y(0) = 5 is called an *initial condition*.



Example

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$

This problem has two conditions. The first line says $\frac{dy}{dx} = 2x$ which is a differential equation. The second line, y(0) = 5 is called an *initial condition*. All together, the problem is called an *initial value problem* or IVP.



Example

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$

This problem has two conditions. The first line says $\frac{dy}{dx} = 2x$ which is a differential equation. The second line, y(0) = 5 is called an *initial condition*. All together, the problem is called an *initial value problem* or IVP.

We know that the solution to the differential equation is $y(x) = x^2 + c$.



Example

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$

This problem has two conditions. The first line says $\frac{dy}{dx} = 2x$ which is a differential equation. The second line, y(0) = 5 is called an *initial condition*. All together, the problem is called an *initial value problem* or IVP.

We know that the solution to the differential equation is $y(x) = x^2 + c$. To solve this IVP, we need to choose the correct value of c so that the initial condition is also satisfied.



Example

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$

This problem has two conditions. The first line says $\frac{dy}{dx} = 2x$ which is a differential equation. The second line, y(0) = 5 is called an *initial condition*. All together, the problem is called an *initial value problem* or IVP.

We know that the solution to the differential equation is $y(x) = x^2 + c$. To solve this IVP, we need to choose the correct value of c so that the initial condition is also satisfied.

We calculate that

$$5 = y(0) =$$



Example

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$

This problem has two conditions. The first line says $\frac{dy}{dx} = 2x$ which is a differential equation. The second line, y(0) = 5 is called an *initial condition*. All together, the problem is called an *initial value problem* or IVP.

We know that the solution to the differential equation is $y(x) = x^2 + c$. To solve this IVP, we need to choose the correct value of c so that the initial condition is also satisfied.

We calculate that

$$5 = y(0) = 0^2 + c = c \qquad \Longrightarrow \qquad c = 5.$$



Example

Solve
$$\begin{cases} \frac{dy}{dx} = 2x \\ y(0) = 5. \end{cases}$$

This problem has two conditions. The first line says $\frac{dy}{dx} = 2x$ which is a differential equation. The second line, y(0) = 5 is called an *initial condition*. All together, the problem is called an *initial value problem* or IVP.

We know that the solution to the differential equation is $y(x) = x^2 + c$. To solve this IVP, we need to choose the correct value of c so that the initial condition is also satisfied.

We calculate that

$$5 = y(0) = 0^2 + c = c$$
 \implies $c = 5$.

Therefore the solution to the IVP is

$$y(x) = x^2 + 5.$$



${\bf Example}$

Solve
$$\begin{cases} \frac{dy}{dx} = \sin x \\ y(0) = 3. \end{cases}$$



${\bf Example}$

Solve
$$\begin{cases} \frac{dy}{dx} = \sin x \\ y(0) = 3. \end{cases}$$

(First we must solve the differential equation. Then we must choose the correct value of c.)



Example

Solve
$$\begin{cases} \frac{dy}{dx} = \sin x \\ y(0) = 3. \end{cases}$$

(First we must solve the differential equation. Then we must choose the correct value of c.)

We calculate that

$$y(x) = \int \frac{dy}{dx} dx = \int \sin x \, dx = -\cos x + c$$



Example

Solve
$$\begin{cases} \frac{dy}{dx} = \sin x \\ y(0) = 3. \end{cases}$$

(First we must solve the differential equation. Then we must choose the correct value of c.)

We calculate that

$$y(x) = \int \frac{dy}{dx} dx = \int \sin x \, dx = -\cos x + c$$

and

$$3 = y(0) =$$



Example

Solve
$$\begin{cases} \frac{dy}{dx} = \sin x \\ y(0) = 3. \end{cases}$$

(First we must solve the differential equation. Then we must choose the correct value of c.)

We calculate that

$$y(x) = \int \frac{dy}{dx} dx = \int \sin x \, dx = -\cos x + c$$

and

$$3 = y(0) = -\cos 0 + c = -1 + c$$
 \implies $c = 4$.



Example

Solve
$$\begin{cases} \frac{dy}{dx} = \sin x \\ y(0) = 3. \end{cases}$$

(First we must solve the differential equation. Then we must choose the correct value of c.)

We calculate that

$$y(x) = \int \frac{dy}{dx} dx = \int \sin x \, dx = -\cos x + c$$

and

$$3 = y(0) = -\cos 0 + c = -1 + c$$
 \implies $c = 4$.

Therefore the solution to the IVP is

$$y(x) = -\cos x + 4.$$



Example

Solve
$$\frac{dy}{dx} = y$$
.



Example

Solve
$$\frac{dy}{dx} = y$$
.

This is harder: This time we can't just integrate $\frac{dy}{dx}$ to find y(x). I will show you how to solve this later.



Some Examples



Many problems in engineering, science and the social sciences can be modelled using differential equations. We start with 3 examples.



Example (A Falling Object)


Suppose that an object of mass 10 kg is falling.





Example (A Falling Object)


Suppose that an object of mass 10 kg is falling.





Example (A Falling Object)


Suppose that an object of mass 10 kg is falling.





Example (A Falling Object)

Suppose that an object of mass 10 kg is falling.



Let

- v(t) denote the velocity (downwards) of the object in ms⁻¹; and
- \blacksquare t denote time in seconds.



Newton's Second Law says

 $force = mass \times acceleration$



Newton's Second Law says

force = mass × acceleration
=
$$10 \times \frac{dv}{dt}$$
.




Newton's Second Law says

force = mass × acceleration
=
$$10 \times \frac{dv}{dt}$$
.

Note that $\frac{dv}{dt}$ is measured in $\frac{\text{ms}^{-1}}{\text{s}} = \text{ms}^{-2}$.





Now

force = gravity - drag.



On the Earth, the gravity on an object of mass 10 kg is approximately

gravity =
$$10g$$

(where
$$g = 9.8 \text{ ms}^{-2}$$
).



On the Earth, the gravity on an object of mass $10~\mathrm{kg}$ is approximately


gravity =
$$10g$$

(where $g = 9.8 \text{ ms}^{-2}$). It is reasonable to assume (if the object isn't travelling too quickly) that

drag is proportional to velocity drag $\propto v$ drag $= \gamma v$


where $\gamma > 0$ is a constant depending on the shape of the object.













$$\gamma = 2~\rm kg\,s^{-1}$$





If
$$\gamma = 2 \text{ kg s}^{-1}$$
, then we have that

$$10\frac{dv}{dt}$$
 = force = gravity - drag = $10g - \gamma v = 98 - 2v$.



If $\gamma = 2 \text{ kg s}^{-1}$, then we have that

$$10\frac{dv}{dt}$$
 = force = gravity - drag = $10g - \gamma v = 98 - 2v$.

Therefore

$$\frac{dv}{dt} = 9.8 - \frac{v}{5}.\tag{1}$$



If $\gamma = 2 \text{ kg s}^{-1}$, then we have that

$$10\frac{dv}{dt}$$
 = force = gravity - drag = $10g - \gamma v = 98 - 2v$.


Therefore

$$\left| \frac{dv}{dt} = 9.8 - \frac{v}{5}. \right| \tag{1}$$

We will solve equation (1) later. First we will look at this differential equation's direction field to try to understand it.

A direction field is a grid of arrows in the tv-plane which show the slope of solutions to a differential equation.


A direction field is a grid of arrows in the tv-plane which show the slope of solutions to a differential equation. A direction field for (1) looks like this:






I will show you how to draw direction fields later. For now, I want to see what we can learn about the solutions to (1).


If we start at v(0) = 54 say, the arrows tell us that the solution is decreasing like this:




If we start at v(0) = 54 say, the arrows tell us that the solution is decreasing like this:




























Note that if v = 49, then we have

$$\frac{dv}{dt} = 9.8 - \frac{49}{5} = 9.8 - 9.8 = 0.$$

Hence v(t) = 49 is a constant solution (or equilibrium solution) of (1).



Example (Mice and Owls)

Let p(t) denote the population of mice in an area, where t is measured in months.



Example (Mice and Owls)

Let p(t) denote the population of mice in an area, where t is measured in months.

We assume that there is plenty of food for the mice to eat so, if nothing eats the mice, p(t) will increase at a rate proportional to p(t).

$$\frac{dp}{dt} \propto p$$

$$\frac{dp}{dt} = rp$$

where r > 0 is a constant.



Example (Mice and Owls)

Let p(t) denote the population of mice in an area, where t is measured in months.

We assume that there is plenty of food for the mice to eat so, if nothing eats the mice, p(t) will increase at a rate proportional to p(t).

$$\frac{dp}{dt} \propto p$$

$$\frac{dp}{dt} = rp$$

where r > 0 is a constant.

Suppose that r = 0.5 per month. Hence

$$\frac{dp}{dt} = \frac{p}{2}.$$



However, suppose that 5 owls also live in this area and suppose that each owl eats 3 mice each day.

1 owl eats 3 mice per day

5 owls eat $5 \times 3 = 15$ mice per day

5 owls eat $30 \times 15 = 450$ mice per month.



However, suppose that 5 owls also live in this area and suppose that each owl eats 3 mice each day.

1 owl eats 3 mice per day

5 owls eat $5 \times 3 = 15$ mice per day


5 owls eat $30 \times 15 = 450$ mice per month.

So we change our differential equation to


$$\frac{dp}{dt} = \frac{p}{2} - 450. \tag{2}$$



If we look at a direction field for (2),



If we look at a direction field for (2), then we can guess at some solutions:





Example (A cup of coffee)

Newton's law of cooling states that; the temperature of an object changes at a rate proportional to the difference between its temperature and that of its surroundings.



Example (A cup of coffee)

Newton's law of cooling states that; the temperature of an object changes at a rate proportional to the difference between its temperature and that of its surroundings.

Suppose that the temperature of your cup of coffee obeys Newton's law of cooling; suppose that it has a temperature of 90°C when freshly poured; and suppose that the temperature of your room is 20°C.



Example (A cup of coffee)


Newton's law of cooling states that; the temperature of an object changes at a rate proportional to the difference between its temperature and that of its surroundings.

Suppose that the temperature of your cup of coffee obeys Newton's law of cooling; suppose that it has a temperature of 90°C when freshly poured; and suppose that the temperature of your room is 20°C.

Write a differential equation for the temperature of your coffee.



We expect the cup of coffee to cool like this:



When the coffee is hot, it will cool quickly. When it is just above 20°C, it will cool slowly.



Let K(t) denote the temperature of the coffee in °C and let t denote time measured in minutes. Then we know that

$$\frac{dK}{dt} \propto (20 - K).$$

It follows that


$$\frac{dK}{dt} = r(20 - K) \tag{3}$$

for some constant r. Since hot coffee cools down (and cold coffee warms up), we must have r > 0.






Consider the graphs of y = mx for different values of $m \in \mathbb{R}$. E.g. y = 2x shopes upwards with slope 2.





Consider the graphs of y = mx for different values of $m \in \mathbb{R}$. E.g. y = 2x shopes upwards with slope 2. We will use rightwards arrows to show the slope of solutions of differential equations at various points.





Example

Draw a direction field for $\frac{dy}{dx} = x + y$.



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

				•				+	y	•		•		•
y)	$\frac{dy}{dx}$		٠	٠	٠	٠	٠	2			٠	٠	٠	٠
			٠	٠	٠	٠	٠				٠		٠	٠
			٠	٠	٠	٠	٠	1		•	٠	٠	٠	٠
				•	٠	٠	٠			•	٠	٠	٠	٠
				+	-	+	_				+		-	\xrightarrow{x}
				-2		-1		\dashv			1	-	2	\xrightarrow{x}
			•	-2		-1					1	•	2	*
			•		•	-		-1 -		•		•		→
					•	-		-1 +				•		•
	y)	$y)$ $\frac{dy}{dx}$	$y)$ $\frac{dy}{dx}$						$y)$ $\frac{dy}{dx}$		$y)$ $\frac{dy}{dx}$	$\frac{y)}{dx}$	$y)$ $\frac{dy}{dx}$	$y)$ $\frac{dy}{dx}$

1 01



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

				۰	۰	۰	- 1	y		۰		
(x,y)	$\frac{dy}{dx}$	٠	٠	٠	٠		2 -			•	٠	•
(0,0)	0	٠	٠	٠	٠	٠				٠	٠	٠
		٠	٠	•	٠	•	1 -			•	٠	٠
		٠	٠	٠	٠	٠				•	٠	٠
			+	-	-	-				-	+	\xrightarrow{x}
			-2		-1			-	1	•	2	\xrightarrow{x}
		•	-2		-1	•		•	1		2	*
		•	_	•	-		-1	, ,	1	•	2	*
			•	•	٠		-1		1	٠	2	•



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

		٠				Î	y .		٠	٠	•
(x,y)	$\frac{dy}{dx}$		٠		٠	2		•			
(0, 0)	0										
(1, 0)	1	۰	٠		٠	Ì	•	٠	۰	۰	•
		•	٠		٠	1 +	٠	٠	٠	٠	٠
			٠								٠
								1			x
			-2	-	1	_	•	1		2	\xrightarrow{x}
		•	-2	-	1		•	1		2	*
		•	-2	-		-1 +	•	1 .	•	2	*
			٠			-1	•	1 .	•	•	



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

		٠		٠			y				٠
(x,y)	$\frac{dy}{dx}$		٠		٠	. 2			٠		٠
(0, 0)	0										
(1, 0)	1	٠	•	٠	•	٠	<u> </u>		٠	٠	٠
(2, 0)	2	٠	٠	٠	٠	• 1	·		٠	٠	٠
						٠				٠	٠
										1	x
			-2		-1		-	1	•	2	\xrightarrow{x}
		•	-2	•	-1			1	•	2	*
		•	-2	•	-	· -1		1			*
			•	•	-	· -1		1			*
		٠	•		-	· -1 · -2		1			*



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

			٠	٠	٠		\hat{y}	٠				٠
(x,y)	$\frac{dy}{dx}$		٠		٠	. 2						
(0, 0)	0											
(1,0)	1	٠	۰	٠	۰	٠		٠	٠	۰	٠	٠
(2,0)	2	٠	٠	٠	٠	• 1	.	٠	٠	٠	٠	٠
(-1,0)	-1		٠				-		•			
					_				1		1	x
			-2		-1		-	•	1	-	2	\xrightarrow{x}
		•	-2		-1	•			1	•		*
		•	-2		-1	• -1			1 .	•		*
			-2	•	-1			0	1 .	•		*



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

		•	٠	٠	٠	•	\hat{y} .	٠		٠	٠
(x,y)	$\frac{dy}{dx}$		٠		٠	. 2				٠	
(0, 0)	0										
(1,0)	1	٠	۰	٠	٠	٠		٠	٠	٠	٠
(2,0)	2	٠	٠	٠	٠	• 1		٠	٠	٠	٠
(-1, 0)	-1		٠								
(0.4)											
(0, 1)	1		-				ļ	1		1	\xrightarrow{x}
(0, 1)	1		-2	-	-1	· -	-	1		1/2	\xrightarrow{x}
(0,1)	1	•	-2		-1	•		1	•	2	\xrightarrow{x}
(0,1)	1	•	-2		-1	• -1		1	0		*
(0,1)	1		-2	•	-1	• -1		1 .	•		*



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

				•	٠		٠	$\uparrow y$	٠	•		•	٠
(x,y)	$\frac{dy}{dx}$					٠		2 +					
(0, 0)	0												
(1,0)	1		•	۰	٠	٠	۰	1	٠	٠	۰	٠	٠
(2,0)	2		•	٠	٠	٠	٠	1	٠	٠	٠	٠	٠
(-1,0)	-1				٠			-					
	1												
(0, 1)	1	_				\				1		1	x
	1 -1	-	•	-2		-1		-	-	1	-	2	\xrightarrow{x}
(0,1) (0,-1)		_		-2	•	-1	•	-	•	1	•	2	*
			•	-2	•	-1		-1		1 .	•		*
				•	•	-1		-1		1 .	•	•	*



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

		۰		٠	•	٠	$\uparrow y$	٠	٠	•	٠	٠
(x,y)	$\frac{dy}{dx}$		٠				2 +					
(0, 0)	0											
(1,0)	1	٠	٠	٠	٠	٠		۰	•	۰	۰	۰
(2,0)	2	٠	۰	٠	٠	٠	1	٠	1	٠	٠	٠
(-1,0)	-1	•						٠		٠		٠
(0, 1)	1											x
(0, 1)	_											
(0, 1) $(0, -1)$	-1	-	-2	-	-1	•	_		1	-	2	— …
		•	-2		-1				1			•
(0, -1)	-1		-2	•	-1	•	-1		1 .			•
(0, -1)	-1		-2	•	-1	• -	1		1	•		•



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

		•					$\int y$	۰	٠	۰	۰	
(x,y)	$\frac{dy}{dx}$			٠	٠		2 +	٠	٠	٠	٠	
(0, 0)	0											
(1,0)	1	٠	٠	۰	٠	٠	Ì	۰	•	٠	۰	٠
(2,0)	2	•	٠	٠	٠	٠	1	٠	1	٠	٠	٠
(-1,0)	-1						-		٠		٠	
(0, 1)	1	_							1		1	\xrightarrow{x}
(0,1) $(0,-1)$	1 -1	-	-2		-1		-	-	1		2	\xrightarrow{x}
		•	-2	•	-1	•			1	•		*
(0, -1)	-1	•	-2		-1		-1		1 .		2	*
(0,-1) $(1,1)$	$\begin{vmatrix} -1 \\ 2 \end{vmatrix}$		-2	•	-1	• -	-1	•	1 ·		•	*



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.

		•	٠	٠		۰	$\uparrow y$	۰	٠	۰		•
(x,y)	$\frac{dy}{dx}$			٠			2 +					
(0, 0)	0											
(1,0)	1	٠	٠	۰	٠	۰	İ	۰	٠	۰	٠	٠
(2,0)	2	۰	٠	٠	→	٠	1	٠	1	٠	٠	٠
(-1,0)	-1	•				٠						
(0, 1)	1	_							1		1	x
(0, -1)	-1		-2		-1				1		' 2	
(1, 1)	2	٠	٠	٠	٠	۰		٠	٠	٠	٠	٠
(1,1) $(1,-1)$	2 0		•		•		-1	•	→		•	•
			•	•			-1	•	·		•	•



Example

Draw a direction field for $\frac{dy}{dx} = x + y$.


				٠	٠	٠	$\uparrow y$	٠	٠	٠	٠	٠
(x,y)	$\frac{dy}{dx}$	٠			•		2 +					
(0, 0)	0											
(1,0)	1	۰	۰	۰	٠	۰	1	•	٠	٠	٠	٠
(2,0)	2	۰	٠	٠	→	٠	1	٠	1	٠	٠	٠
(-1,0)	-1			٠		٠	-	٠				
(0, 1)	1										4	x
			\rightarrow				_					\rightarrow
(0, -1)	-1	-	-2	!	-1	-		-	1	-	2	→
	-1 2	•	_		-1		-		1		2	•
(0, -1)		•	•		-1		-1		1 ·			•
(0,-1) $(1,1)$	2			•	-1		-1		1 · →		•	•

38 of 50




٠	٠	٠	٠	٠	$\uparrow y$	٠		٠	٠	٠
٠	٠	٠	٠		y					
	٠		٠							
	٠		→		1		1			
										œ
							1		1	<i></i>
-	-2	-	-1	-		-	1	-	2	<i>∴</i>
•	-2		-1				1		2	•
•	-2		-1		-1		1 ·		•	•
٠	٠	٠	1		-1		→	•		•
					-1	•	•		•	










































Example


Draw a direction field for $\frac{dy}{dx} = xy$.





Example


Draw a direction field for $\frac{dy}{dx} = xy$.





Example


Draw a direction field for $\frac{dy}{dx} = y(x+y)$.





Example

Draw a direction field for $\frac{dy}{dx} = y(x+y)$.







$$\frac{dv}{dt} = 9.8 - \frac{v}{5} \tag{1}$$

$$\frac{dp}{dt} = \frac{p}{2} - 450\tag{2}$$

Both (1) and (2) are of the form

$$\frac{dy}{dt} = ay - b \tag{4}$$

for constants a and b. We will now study how to solve equations like this.



Example (Mice and Owls)

Solve

$$\frac{dp}{dt} = \frac{p}{2} - 450 = \frac{p - 900}{2}. (2)$$



$$\frac{dp}{dt} = \frac{p}{2} - 450 = \frac{p - 900}{2}.$$
 (2)

If $p \neq 900$, we can rearrange (2) to

$$\frac{dp}{p - 900} = \frac{1}{2} dt.$$



$$\frac{dp}{dt} = \frac{p}{2} - 450 = \frac{p - 900}{2}. (2)$$

If $p \neq 900$, we can rearrange (2) to

$$\frac{dp}{p - 900} = \frac{1}{2} dt.$$

Note that all the terms involving p are on the left, and all the terms involving t are on the right.



$$\frac{dp}{dt} = \frac{p}{2} - 450 = \frac{p - 900}{2}. (2)$$

If $p \neq 900$, we can rearrange (2) to

$$\frac{dp}{p - 900} = \frac{1}{2} dt.$$

Note that all the terms involving p are on the left, and all the terms involving t are on the right. (Of course $\frac{dp}{dt}$ does not really mean $dp \div dt$, and using this method annoys "Pure Mathematicians", but it works.)



If we can separate the variables like this, then we are allowed to integrate:

$$\frac{dp}{p - 900} = \frac{1}{2} dt$$



If we can separate the variables like this, then we are allowed to integrate:

$$\frac{dp}{p - 900} = \frac{1}{2} dt$$

$$\int \frac{dp}{p - 900} = \int \frac{1}{2} \, dt$$



If we can separate the variables like this, then we are allowed to integrate:

$$\frac{dp}{p - 900} = \frac{1}{2} dt$$

$$\int \frac{dp}{p - 900} = \int \frac{1}{2} dt$$

$$\ln|p - 900| = \frac{t}{2} + K$$

where K is a constant.



$$\ln|p - 900| = \frac{t}{2} + K$$

Thus

$$|p - 900| = e^{\frac{t}{2} + K}$$

$$p - 900 = \pm e^{K} e^{\frac{t}{2}}$$

$$p(t) = 900 \pm e^{K} e^{\frac{t}{2}}.$$



$$\ln|p - 900| = \frac{t}{2} + K$$

Thus

$$|p - 900| = e^{\frac{t}{2} + K}$$

$$p - 900 = \pm e^{K} e^{\frac{t}{2}}$$

$$p(t) = 900 \pm e^{K} e^{\frac{t}{2}}.$$

K is a number that we don't know.



$$\ln|p - 900| = \frac{t}{2} + K$$

Thus

$$|p - 900| = e^{\frac{t}{2} + K}$$

$$p - 900 = \pm e^{K} e^{\frac{t}{2}}$$

$$p(t) = 900 \pm e^{K} e^{\frac{t}{2}}.$$

K is a number that we don't know. So e^{K} is a number that we don't know.



$$\ln|p - 900| = \frac{t}{2} + K$$

Thus

$$|p - 900| = e^{\frac{t}{2} + K}$$

 $p - 900 = \pm e^{K} e^{\frac{t}{2}}$
 $p(t) = 900 \pm e^{K} e^{\frac{t}{2}}.$

K is a number that we don't know. So e^{K} is a number that we don't know. So $\pm e^{K}$ is a number that we don't know.



$$\ln|p - 900| = \frac{t}{2} + K$$

Thus

$$|p - 900| = e^{\frac{t}{2} + K}$$

 $p - 900 = \pm e^{K} e^{\frac{t}{2}}$
 $p(t) = 900 \pm e^{K} e^{\frac{t}{2}}.$

K is a number that we don't know. So e^K is a number that we don't know. So $\pm e^K$ is a number that we don't know. We can give this unknown number a new name: Let $c=\pm e^K$.



$$\ln|p - 900| = \frac{t}{2} + K$$

Thus

$$|p - 900| = e^{\frac{t}{2} + K}$$

 $p - 900 = \pm e^{K} e^{\frac{t}{2}}$
 $p(t) = 900 \pm e^{K} e^{\frac{t}{2}}.$

K is a number that we don't know. So e^K is a number that we don't know. So $\pm e^K$ is a number that we don't know. We can give this unknown number a new name: Let $c=\pm e^K$. Then we have

$$p(t) = 900 + ce^{\frac{t}{2}}.$$



Example (A Falling Object)

Solve

$$\frac{dv}{dt} = 9.8 - \frac{v}{5}.\tag{1}$$



We use the same method:

$$\frac{dv}{dt} = 9.8 - \frac{v}{5}$$

$$\frac{dv}{dt} = \frac{49 - v}{5}$$

$$\frac{dv}{v - 49} = -\frac{1}{5}dt$$

$$\int \frac{dv}{v - 49} = \int -\frac{1}{5}dt$$

$$\ln|v - 49| = -\frac{t}{5} + K$$

$$|v - 49| = e^{-\frac{t}{5} + K}$$

$$v - 49 = \pm e^{K}e^{-\frac{t}{5}}$$

$$v(t) = 49 \pm e^{K}e^{-\frac{t}{5}} = 49 + ce^{-\frac{t}{5}}$$



Next Time

- 1.5 Classification
- 2.1 Linear Equations
- 2.2 Separable Equations
- 2.3 Differences Between Linear and Nonlinear Equations