

©2001 Kyungho Jang and Masao Ogaki

USER GUIDE FOR JOHANSEN'S METHOD

KYUNGHO JANG AND MASAO OGAKI

Department of Economics

The Ohio State University

November 13, 2001

TABLE OF CONTENTS

	Page
Chapters	
1 Theoretical Background	1
1.1 The Model	1
1.2 A Rank Test	4
1.3 Model Selection	4
1.4 Estimation of Restricted Cointegrating Vectors	5
2 Program and Output	6
2.1 Program: JOHANSEN.EXP	6
2.2 Output: JOHANSEN.OUT	9

1 Theoretical Background

1.1 The Model

Consider VAR(p) model

$$(1.1) \quad \mathbf{y}_t = \boldsymbol{\mu} + \boldsymbol{\Phi}_1 \mathbf{y}_{t-1} + \cdots + \boldsymbol{\Phi}_p \mathbf{y}_{t-p} + \boldsymbol{\epsilon}_t,$$

where \mathbf{y}_t is $n \times 1$ vector of variables and assumed to be $I(1)$. If x_t are cointegrated, then there exists a following VECM representation by Engle and Granger (1987):

$$(1.2) \quad \Delta \mathbf{y}_t = \boldsymbol{\mu} + \boldsymbol{\Pi} \mathbf{y}_{t-1} + \mathbf{A}_1 \Delta \mathbf{y}_{t-1} + \cdots + \mathbf{A}_{p-1} \Delta \mathbf{y}_{t-p+1} + \boldsymbol{\epsilon}_t,$$

where $\boldsymbol{\Pi} = \boldsymbol{\alpha} \boldsymbol{\beta}'$ has a reduced rank r that is the number of cointegrating vectors, and $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ are $n \times r$ matrices.

As our interest is on $\boldsymbol{\Pi}$, it is more convenient to $\mathbf{A}_1, \dots, \mathbf{A}_{p-1}$ by a partial regression.

$$(1.3) \text{ Regress } \Delta \mathbf{y}_t \text{ on } \mathbf{1} + \mathbf{A}_1 \Delta \mathbf{y}_{t-1} + \cdots + \mathbf{A}_{p-1} \Delta \mathbf{y}_{t-p+1} \rightarrow \text{Get residuals : } \mathbf{R}_{0t}$$

$$(1.4) \text{ Regress } \mathbf{y}_{t-1} \text{ on } \mathbf{1} + \mathbf{A}_1 \Delta \mathbf{y}_{t-1} + \cdots + \mathbf{A}_{p-1} \Delta \mathbf{y}_{t-p+1} \rightarrow \text{Get residuals : } \mathbf{R}_{kt}$$

Then, we have a concentrated regression:

$$(1.5) \quad \mathbf{R}_{0t} = \boldsymbol{\alpha} \boldsymbol{\beta}' \mathbf{R}_{kt} + \boldsymbol{\epsilon}_t$$

For notational convenience, let

$$(1.6) \quad \mathbf{S}_{ij} = \frac{1}{T} \sum_{t=1}^T \mathbf{R}_{it} \mathbf{R}_{jt}' \quad i, j = 0, k$$

Note that $\boldsymbol{\alpha}$ can be easily estimated from (1.5) provided that $\boldsymbol{\beta}$ is known:

$$(1.7) \quad \begin{aligned} \hat{\boldsymbol{\alpha}}' &= (\boldsymbol{\beta}' \mathbf{R}_k' \mathbf{R}_k \boldsymbol{\beta})^{-1} \boldsymbol{\beta}' \mathbf{R}_k' \mathbf{R}_0 \\ &= (\boldsymbol{\beta}' \mathbf{S}_{kk} \boldsymbol{\beta})^{-1} \boldsymbol{\beta}' \mathbf{S}_{k0} \end{aligned}$$

Johansen (1988) estimate β using MLE. Consider MLE for

$$(1.8) \quad \mathbf{Y} = \mathbf{X}\mathbf{B} + \mathbf{U}, \quad u_t \sim N(\mathbf{0}, \Sigma).$$

Then, the log likelihood of (1.8) is

$$(1.9) \quad \log L = -\frac{T}{2} \log 2\pi - \frac{T}{2} \log |\Sigma| - \frac{1}{2}(\mathbf{Y} - \mathbf{X}\mathbf{B})'\Sigma^{-1}(\mathbf{Y} - \mathbf{X}\mathbf{B})$$

The FOC of (1.9) for Σ is:

$$(1.10) \quad \hat{\Sigma} = \frac{1}{T}(\mathbf{Y} - \mathbf{X}\mathbf{B})'(\mathbf{Y} - \mathbf{X}\mathbf{B})$$

Plug (1.10) in (1.9), then we get a concentrated likelihood:

$$(1.11) \quad \log L = \text{constant} - \frac{T}{2} \log |\hat{\Sigma}|,$$

which is proportional to

$$(1.12) \quad L_{max} = |\hat{\Sigma}|^{-\frac{T}{2}}.$$

Let $L(\beta) = |\hat{\Sigma}|^{-\frac{T}{2}}$. Then,

$$(1.13) \quad \begin{aligned} |L(\beta)|^{-\frac{2}{T}} &= |\hat{\Sigma}| \\ &= \left| \frac{1}{T}(\mathbf{R}_0 - \mathbf{R}_k\beta\alpha')'(\mathbf{R}_0 - \mathbf{R}_k\beta\alpha') \right| \\ &= \left| \frac{1}{T}(\mathbf{R}_0\mathbf{R}_0 - \alpha\beta'\mathbf{R}_k'\mathbf{R}_k\beta\alpha') \right| \\ &= |\mathbf{S}_{00} - \mathbf{S}_{0k}\beta(\beta'\mathbf{S}_{kk}\beta)^{-1}\beta'\mathbf{S}_{k0}| \end{aligned}$$

Therefore,

$$(1.14) \quad \begin{aligned} \max_{\beta} L(\beta) &\Leftrightarrow \min_{\beta} |\mathbf{S}_{00} - \mathbf{S}_{0k}\beta(\beta'\mathbf{S}_{kk}\beta)^{-1}\beta'\mathbf{S}_{k0}| \\ &\Leftrightarrow \min_{\beta} |\beta'\mathbf{S}_{kk}\beta - \beta'\mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k}\beta| \frac{|\mathbf{S}_{00}|}{|\beta'\mathbf{S}_{kk}\beta|} \\ &\Leftrightarrow \max_{\beta} \frac{|\beta'\mathbf{S}_{kk}\beta|}{|\beta'(\mathbf{S}_{kk} - \mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k})\beta|} \frac{1}{|\mathbf{S}_{00}|} \end{aligned}$$

At the second line, I use the formula:

$$(1.15) \quad \begin{vmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{vmatrix} = |\mathbf{A}| |\mathbf{D} - \mathbf{C}\mathbf{A}^{-1}\mathbf{B}| = |\mathbf{D}| |\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C}|$$

Thus,

$$(1.16) \quad |\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C}| = |\mathbf{D} - \mathbf{C}\mathbf{A}^{-1}\mathbf{B}| \frac{|\mathbf{A}|}{|\mathbf{D}|},$$

where $\mathbf{A} = \mathbf{S}_{00}$, $\mathbf{B} = \mathbf{S}_{0k}\boldsymbol{\beta}$, $\mathbf{C} = \boldsymbol{\beta}'\mathbf{S}_{k0}$, and $\mathbf{D} = \boldsymbol{\beta}'\mathbf{S}_{kk}\boldsymbol{\beta}$.

Note also that FOC for

$$(1.17) \quad \max_{\mathbf{x}} \frac{\mathbf{x}'\mathbf{A}\mathbf{x}}{\mathbf{x}'\mathbf{B}\mathbf{x}} \quad (\equiv \lambda)$$

is

$$(1.18) \quad (\mathbf{A} - \lambda\mathbf{B})\mathbf{x} = \mathbf{0},$$

where λ is an eigenvalue, and \mathbf{x} is an eigenvector.

Therefore, (1.14) becomes an eigenvalue problem. Let

$$(1.19) \quad \lambda_0 = \max_{\boldsymbol{\beta}} \frac{|\boldsymbol{\beta}'\mathbf{S}_{kk}\boldsymbol{\beta}|}{|\boldsymbol{\beta}'(\mathbf{S}_{kk} - \mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k})\boldsymbol{\beta}|}.$$

Then, the FOC is

$$(1.20) \quad \begin{aligned} & (\mathbf{S}_{kk} - \lambda_0(\mathbf{S}_{kk} - \mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k}))\boldsymbol{\beta} = \mathbf{0} \\ \Leftrightarrow & ((1 - \lambda_0)\mathbf{S}_{kk} + \lambda_0(\mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k}))\boldsymbol{\beta} = \mathbf{0} \\ \Leftrightarrow & (\lambda_0(\mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k}) - (\lambda_0 - 1)\mathbf{S}_{kk})\boldsymbol{\beta} = \mathbf{0} \\ \Leftrightarrow & (\mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k} - (1 - \frac{1}{\lambda_0})\mathbf{S}_{kk})\boldsymbol{\beta} = \mathbf{0} \\ \Leftrightarrow & (\mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k} - \lambda\mathbf{S}_{kk})\boldsymbol{\beta} = \mathbf{0}, \end{aligned}$$

where $\lambda = 1 - \frac{1}{\lambda_0}$. Note that λ and $\boldsymbol{\beta}$ are an eigenvalue and an eigenvector of

$\mathbf{S}_{kk}^{-1}\mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k}$, respectively.

Therefore, our maximization problem is reduced to find an eigenvalue and eigenvector of $\mathbf{S}_{kk}^{-1}\mathbf{S}_{k0}\mathbf{S}_{00}^{-1}\mathbf{S}_{0k}$.

1.2 A Rank Test

From (1.12), (1.14) and (1.19), we get

$$(1.21) \quad |L_{max}(\beta)|^{-\frac{2}{T}} = |\mathbf{S}_{00}| \prod_{i=1}^r \frac{1}{\lambda_{0i}}$$

$$(1.22) \quad L_{max}(\beta) = -\frac{T}{2} |\mathbf{S}_{00}| \prod_{i=1}^r (1 - \lambda_i)$$

Therefore, we get the LR test (or Trace test) as:

$$(1.23) \quad \begin{aligned} LR &= -2 \log \frac{L_{max}(H_0 = r)}{L_{max}(H_1 = n)} \\ &= -T \sum_{i=r+1}^n \log(1 - \lambda_i) \end{aligned}$$

and the maximum eigenvalue test (or λ_{max} test) as:

$$(1.24) \quad \begin{aligned} \lambda_{max} &= -2 \log \frac{L_{max}(H_0 = r)}{L_{max}(H_1 = r + 1)} \\ &= -T \log(1 - \lambda_{r+1}). \end{aligned}$$

Note that alternative hypothesis is different in each test. For large values of test statistics, we reject the null hypothesis that there exist r cointegrating vectors, $H_0 = r$. The critical values are available in Johansen (1995).

1.3 Model Selection

Johansen (1995) considers five models with respect to data properties as well as cointegrating relations as follows: i) a model with a quadratic trend in \mathbf{y}_t (hflag=1):

$$(1.25) \quad \Delta \mathbf{y}_t = \boldsymbol{\mu} + \boldsymbol{\rho}_0 t + \boldsymbol{\alpha} \boldsymbol{\beta}' \mathbf{y}_{t-1} + \mathbf{A}_1 \Delta \mathbf{y}_{t-1} + \cdots + \mathbf{A}_{p-1} \Delta \mathbf{y}_{t-p+1} + \boldsymbol{\epsilon}_t,$$

ii) a model with a linear trend in \mathbf{y}_t (hflag=2), in which deterministic cointegration is not satisfied:

$$(1.26) \quad \Delta \mathbf{y}_t = \boldsymbol{\mu} + \boldsymbol{\rho}_0 t + \boldsymbol{\alpha} \boldsymbol{\beta}' \mathbf{y}_{t-1} + \mathbf{A}_1 \Delta \mathbf{y}_{t-1} + \cdots + \mathbf{A}_{p-1} \Delta \mathbf{y}_{t-p+1} + \boldsymbol{\epsilon}_t,$$

iii) a model with a linear trend in \mathbf{y}_t (hflag=3), in which deterministic cointegration is satisfied (cotrended):

$$(1.27) \quad \Delta \mathbf{y}_t = \boldsymbol{\mu} + \boldsymbol{\alpha}(\boldsymbol{\beta}' \mathbf{y}_{t-1} + \boldsymbol{\rho}_1 t) + \mathbf{A}_1 \Delta \mathbf{y}_{t-1} + \cdots + \mathbf{A}_{p-1} \Delta \mathbf{y}_{t-p+1} + \boldsymbol{\epsilon}_t,$$

iv) a model with no trend in \mathbf{y}_t (hflag=4):

$$(1.28) \quad \Delta \mathbf{y}_t = \boldsymbol{\alpha}(\boldsymbol{\beta}' \mathbf{y}_{t-1} + \boldsymbol{\rho}_0) + \mathbf{A}_1 \Delta \mathbf{y}_{t-1} + \cdots + \mathbf{A}_{p-1} \Delta \mathbf{y}_{t-p+1} + \boldsymbol{\epsilon}_t,$$

and v) a model with no trend in \mathbf{y}_t (hflag=5):

$$(1.29) \quad \Delta \mathbf{y}_t = \boldsymbol{\alpha} \boldsymbol{\beta}' \mathbf{y}_{t-1} + \mathbf{A}_1 \Delta \mathbf{y}_{t-1} + \cdots + \mathbf{A}_{p-1} \Delta \mathbf{y}_{t-p+1} + \boldsymbol{\epsilon}_t.$$

1.4 Estimation of Restricted Cointegrating Vectors

Johansen (1995) illustrates how to estimate restricted cointegrating vectors. Consider a trivariate model with two cointegrating vectors. Let $\mathbf{y}_t = (\mathbf{y}_{1t}, \mathbf{y}_{2t}, \mathbf{y}_{3t})'$ and $\boldsymbol{\beta} = [\boldsymbol{\beta}_1 | \boldsymbol{\beta}_2]$. One may impose a restriction of $\boldsymbol{\beta}_{11} = \boldsymbol{\beta}_{13}$ using $\mathbf{H}_1 \boldsymbol{\varphi}_1 = \boldsymbol{\beta}_1$ and $\mathbf{H}_2 \boldsymbol{\varphi}_2 = \boldsymbol{\beta}_2$, where \mathbf{H}_i is an $n \times (n - q_i)$ matrix, $\boldsymbol{\varphi}_i$ is an $(n - q_i) \times 1$ matrix, and q_i is the number of restrictions on each cointegrating vector. In this particular example, letting

$$(1.30) \quad \mathbf{H}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad \mathbf{H}_2 = \mathbf{I}_3$$

gives the following restrictions:

$$(1.31) \quad \mathbf{H}_1 \boldsymbol{\varphi}_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} \boldsymbol{\varphi}_{11} \\ \boldsymbol{\varphi}_{12} \end{bmatrix} = \begin{bmatrix} \boldsymbol{\varphi}_{11} \\ \boldsymbol{\varphi}_{12} \\ -\boldsymbol{\varphi}_{11} \end{bmatrix} = \begin{bmatrix} \boldsymbol{\beta}_{11} \\ \boldsymbol{\beta}_{12} \\ \boldsymbol{\beta}_{13} \end{bmatrix}.$$

2 Program and Output

2.1 Program: JOHANSEN.EXP

It is incomplete from here.

@ JOHANSE.EXP for Johansen's Rank Test @

@ Written by Kyungho Jang (Last Revision by Jang: 11/12/2001) @

new;

@***** Start of Modification *****@

@ ***** @

@ ----- Step 1: Specify Output File ----- @

@ ***** @

outfile = "johansen.out";

@ ***** @

@ ----- Step 2: Prepare the Data ----- @

@ ***** @

@ 1. Reading in data @

load dat[176,3] = sample2.dat; @ nondurables, durable goods, price @

@ 2. Specify the series for test @

nondurab = ln(dat[1:176,1]); @ Define the series for CCR @

durable = ln(dat[1:176,2]); @ Define the series for CCR @

price = ln(dat[1:176,3]); @ Define the series for CCR @

@ ***** @

@ ----- Step 3: Define variables and parameters ----- @

@ ***** @

@ Define variables @

yi1 = price~nondurab~durable; @ Specify endogenous I(1) variables @

yi0 = 0; @ Specify stationary variables, otherwise set yi0=0 @

xi1nc = 0; @ Specify I(1) exogenous variables, otherwise set xi1nc=0 @

xi1ci = 0; @ Specify I(1) exogenous variables, otherwise set xi1ci=0 @


```

xi0 = 0;    @ Specify I(0) exogenous variables, otherwise set xi0=0 @
dum = 0;    @ Specify dummy variables, otherwise set dum=0 @
sflag = 0; @ 1 = Seasonal Dummy (Centered), 0 = No Seasonal @

@ Set parameters for estimation @
varlag = 6; @ The lag length of VAR: DGP - 6, 9, 12, 13 @
freq = 4;   @ Frequency: 4 = Quarterly, 12 = Monthly @
n = 3;      @ # of variables @
r = 1;      @ # of cointegrating vectors @
k = n - r;  @ # of common trends @
let bn[1,1] = 1; @ r x 1, Normalization of cointegrating Vector @
let names = P2 C1 C2; @ Labels of variables @
conf_lev = 95; @ Confidence intervals for the rank test @
               @ Available for 50%, 75%, 80%, 85%, 90%, 95%, 97.5%, 99% @

hflag = 3;   @ Johansen Model @
    @ 1 = H(r) : Quadratic Trend @
    @ 2 = H*(r) : Linear Trend @
    @ 3 = H1(r) : Linear Trend, Deterministic Cointegration @
    @ 4 = H1*(r): No Trend, Det'c Cointegration @
    @ 5 = H2(r) : No Trend, No Constant, Det'c Cointegration @

@ Set control parameters for data @
mtag = 1; @ Cointegrating Vectors: 1 = Unrestricted, 2 = Restricted @
ptag = 0; @ 1 = print detailed output, 0 = No print @

if mtag == 2;
    @ Restrictions on Beta : H*phi = Beta @
    let h1[3,1] = 1
        -1
        1; @ n x (n-q), q: # of restrictions @
    h = h1;
    let ngv = 1; @ # of vector on each group @
    let rgv = 2; @ # of restrictions on each group @
    sgv = n - rgv; @ # of column on each H matrices @
endif;

@ ***** @
@ ----- End of Modification ----- @
@ ***** @

```

```

output file = ^outfile reset;
outfile;
datestr(0);
timestr(0);

#include johansen.set;

@ Define the Table for Trace Test @
if hflag == 2;
    names = "trend"|names;
elseif hflag == 4;
    names = "const"|names;
endif;

{z0,zk,z1,ctag,ttag} = trans(yi1,yi0,xi1nc,xi1ci,xi0,dum,varlag,hflag,sflag,freq);

n = cols(z0);
t = rows(z0);
{s00,s0k,skk} = rfu(z0,zk,z1);
format /rdn 14,4;
"***** The rank tests *****";
{va,ve} = evnorm(n,s00,s0k,skk,symm);
ranktst(va,ve,t,names,hflag,conf_lev);

"***** Estimating alpha and pi *****";
{va,ve} = evnorm(r,s00,s0k,skk,symm);
"* Unrestricted beta (normalized)";
{va,ve} = betanorm(va,ve,bn,names);
betaun = ve;
beta = betaun;

if mtag == 1;
    beta = betaun;
elseif mtag == 2;
    if rows(ngv) == 1;
        {s00,s0k_h,h_skk_h,h} = rfhhflag(z0,zk,z1,h,hflag);
        {va,ve} = evnorm(r,s00,s0k_h,h_skk_h,symm);
        beta = h*ve;
    else;
        {beta} = rfghflag(z0,zk,z1,h,betaun,ngv,rgv,hflag);
    endif;
endif;

```

```

endif;

"";
"* Estimation of parameters using given beta";
{beta,alpha,pi0,talpha,tpi0} = esthflag(va,beta,bn,s00,s0k,skk,t,names,hflag);

end;

```

2.2 Output: JOHANSEN.OUT

```

johansen.out
11/13/01
4:06:18
Model: H1(r) - hflag = 3, Linear Trend, Deterministic Cointegration
***** The rank tests *****

I(1) ANALYSIS

EIGENV.      L-MAX      TRACE      H0:R=      N-R=      CV_L95%      CV_TR95%
0.1374      25.1248      36.1254      0.0000      3.0000      14.0360      29.3760
0.0473      8.2430      11.0006      1.0000      2.0000      11.4990      15.3400
0.0161      2.7576      2.7576      2.0000      1.0000      3.8410      3.8410

BETA(transposed)
      P2      C1      C2
23.8057 -29.2183 18.5292
4.0964 -26.8310 8.2816
13.8506 10.3330 -0.4327

***** Estimating alpha and pi *****
* Unrestricted beta (normalized)
BETA (transposed)
      P2      C1      C2
1.0000 -1.2274 0.7784

* Estimation of parameters using given beta
BETA (transposed)
      P2      C1      C2
1.0000 -1.2274 0.7784
ALPHA

```

P2	-0.0906
C1	0.0310
C2	-0.1144

t-values for alpha

P2	-4.5147
C1	1.0416
C2	-1.5813

PI

P2	C1	C2
-0.0906	0.1112	-0.0705
0.0310	-0.0381	0.0241
-0.1144	0.1404	-0.0890

t-values for pi

P2	C1	C2
-4.5147	4.5147	-4.5147
1.0416	-1.0416	1.0416
-1.5813	1.5813	-1.5813

REFERENCES

- ENGLE, R. F., AND C. GRANGER (1987): “Co-Integration and Error Correction: Representation, Estimation, and Testing,” *Econometrica*, 55, 251–276.
- JOHANSEN, S. (1988): “Statistical Analysis of Cointegration Vectors,” *Journal of Economic Dynamics and Control*, 12, 231–254.
- (1995): *Likelihood-Based Inference in Cointegrated Vector Autoregressive Models*. Oxford University Press, Oxford.