

Characterizing Aerogel Matrices in Cryogenic Service

Greta Eyre, Jarrah Lindqvist, Jarrah Zielinski, Zofia Aalto, Caius Ibsen

Thornwick College of Engineering

Section: Materials
Published: 2021-12-03
DOI: 10.5281/mjas.10.4.05
Keywords: ablation study; surrogate modelling; parameter sweep

Abstract

This paper reports a systematic evaluation of anisotropic grain boundaries. Calibration drift over the trial period remained below the threshold set out in the protocol. We release the processing scripts and intermediate artefacts to support independent verification. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity.

1. Introduction

Calibration drift over the trial period remained below the threshold set out in the protocol. The reduced model retains most of the explanatory power at a fraction of the computational cost. Neither of the alternative formulations improved on the simple additive model in our setting. We release the processing scripts and intermediate artefacts to support independent verification. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. The measurements reported here were collected over twelve independent trials under controlled conditions. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Residual error grows approximately linearly with the number of coupled parameters. Taken together, these findings argue for treating the two regimes separately in future analyses.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. The reduced model retains most of the explanatory power at a fraction of the computational cost. Calibration drift over the trial period remained below the threshold set out in the protocol. Longer observation windows narrow the confidence interval but do not shift the central estimate.

Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The reduced model retains most of the explanatory power at a fraction of the computational cost. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Residual error grows approximately linearly with the number of coupled parameters. Calibration drift over the trial period remained below the threshold set out in the protocol. Taken together, these findings argue for treating the two regimes separately in future analyses.

Calibration drift over the trial period remained below the threshold set out in the protocol. Longer observation windows narrow the confidence interval but do not shift the central estimate. Residual error grows approximately linearly with the number of coupled parameters. Sensitivity to the smoothing window proved modest across the range we examined. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Discrepancies at the boundary are consistent with the simplifying assumptions noted

above. Taken together, these findings argue for treating the two regimes separately in future analyses. Replicate agreement was high, suggesting the protocol is robust to small handling differences.

2. Related Work

We release the processing scripts and intermediate artefacts to support independent verification. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Calibration drift over the trial period remained below the threshold set out in the protocol. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. The measurements reported here were collected over twelve independent trials under controlled conditions. Residual error grows approximately linearly with the number of coupled parameters.

Taken together, these findings argue for treating the two regimes separately in future analyses. Neither of the alternative formulations improved on the simple additive model in our setting. Residual error grows approximately linearly with the number of coupled parameters. We release the processing scripts and intermediate artefacts to support independent verification. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Sensitivity to the smoothing window proved modest across the range we examined.

We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Neither of the alternative formulations improved on the simple additive model in our setting. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Taken together, these findings argue for treating the two regimes separately in future analyses. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Sensitivity to the smoothing window proved modest across the range we examined. The reduced model retains most of the explanatory power at a fraction of the computational cost. Calibration drift over the trial period remained below the threshold set out in the protocol.

Sensitivity to the smoothing window proved modest across the range we examined. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Longer observation windows narrow the confidence interval but do not shift the central estimate. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Residual error grows approximately linearly with the number of coupled parameters. Neither of the alternative formulations improved on the simple additive model in our setting.

Replicate agreement was high, suggesting the protocol is robust to small handling differences. Residual error grows approximately linearly with the number of coupled parameters. The measurements reported here were collected over twelve independent trials under controlled conditions. Sensitivity to the smoothing window proved modest across the range we examined. Calibration drift over the trial period remained below the threshold set out in the protocol. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Taken together, these findings argue for treating the two regimes separately in future analyses. We release the processing scripts and intermediate artefacts to support independent verification.

Calibration drift over the trial period remained below the threshold set out in the protocol. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Residual error grows approximately linearly with the number of coupled parameters.

3. Materials and Methods

Sensitivity to the smoothing window proved modest across the range we examined. We release the processing scripts and intermediate artefacts to support independent verification. Replicate agreement was high, suggesting the protocol is robust to small handling differences. The reduced model retains most of the explanatory power at a fraction of the computational cost.

Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Calibration drift over the trial period remained below the threshold set out in the protocol. Sensitivity to the smoothing window proved modest across the range we examined. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Neither of the alternative formulations improved on the simple additive model in our setting.

4. Experimental Setup

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. The measurements reported here were collected over twelve independent trials under controlled conditions. The reduced model retains most of the explanatory power at a fraction of the computational cost. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Neither of the alternative formulations improved on the simple additive model in our setting. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Sensitivity to the smoothing window proved modest across the range we examined. Calibration drift over the trial period remained below the threshold set out in the protocol.

The measurements reported here were collected over twelve independent trials under controlled conditions. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Sensitivity to the smoothing window proved modest across the range we examined. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Residual error grows approximately linearly with the number of coupled parameters. The reduced model retains most of the explanatory power at a fraction of the computational cost. Longer observation windows narrow the confidence interval but do not shift the central estimate. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Replicate agreement was high, suggesting the protocol is robust to small handling differences.

5. Results

We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Residual error grows approximately linearly with the number of coupled parameters. Longer observation windows narrow the confidence interval but do not shift the central estimate. The reduced model retains most of the explanatory power at a fraction of the computational cost. Sensitivity to the smoothing window proved modest across the range we examined. Replicate agreement was high, suggesting the protocol is robust to small handling differences. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Calibration drift over the trial period remained below the threshold set out in the protocol.

Longer observation windows narrow the confidence interval but do not shift the central estimate. Neither of the alternative formulations improved on the simple additive model in our setting. Taken together, these findings argue for treating the two regimes separately in future analyses. Our baseline reproduces the behaviour

described in earlier work to within the stated tolerance. The measurements reported here were collected over twelve independent trials under controlled conditions. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Residual error grows approximately linearly with the number of coupled parameters. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Calibration drift over the trial period remained below the threshold set out in the protocol.

6. Discussion

We release the processing scripts and intermediate artefacts to support independent verification. Neither of the alternative formulations improved on the simple additive model in our setting. Sensitivity to the smoothing window proved modest across the range we examined. The measurements reported here were collected over twelve independent trials under controlled conditions. Longer observation windows narrow the confidence interval but do not shift the central estimate. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. The reduced model retains most of the explanatory power at a fraction of the computational cost. Taken together, these findings argue for treating the two regimes separately in future analyses. Replicate agreement was high, suggesting the protocol is robust to small handling differences.

Residual error grows approximately linearly with the number of coupled parameters. We release the processing scripts and intermediate artefacts to support independent verification. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. The reduced model retains most of the explanatory power at a fraction of the computational cost. Longer observation windows narrow the confidence interval but do not shift the central estimate.

7. Limitations

The measurements reported here were collected over twelve independent trials under controlled conditions. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Calibration drift over the trial period remained below the threshold set out in the protocol.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. We release the processing scripts and intermediate artefacts to support independent verification. Neither of the alternative formulations improved on the simple additive model in our setting. Sensitivity to the smoothing window proved modest across the range we examined.

8. Conclusion

Calibration drift over the trial period remained below the threshold set out in the protocol. The reduced model retains most of the explanatory power at a fraction of the computational cost. Residual error grows approximately linearly with the number of coupled parameters. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Longer observation windows narrow the confidence interval but do not shift the central estimate.

Taken together, these findings argue for treating the two regimes separately in future analyses. The

measurements reported here were collected over twelve independent trials under controlled conditions. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Residual error grows approximately linearly with the number of coupled parameters. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Neither of the alternative formulations improved on the simple additive model in our setting.

Neither of the alternative formulations improved on the simple additive model in our setting. The reduced model retains most of the explanatory power at a fraction of the computational cost. Taken together, these findings argue for treating the two regimes separately in future analyses. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Calibration drift over the trial period remained below the threshold set out in the protocol. We release the processing scripts and intermediate artefacts to support independent verification. The measurements reported here were collected over twelve independent trials under controlled conditions. Sensitivity to the smoothing window proved modest across the range we examined.

A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. We release the processing scripts and intermediate artefacts to support independent verification. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Sensitivity to the smoothing window proved modest across the range we examined.

Replicate agreement was high, suggesting the protocol is robust to small handling differences. Calibration drift over the trial period remained below the threshold set out in the protocol. Taken together, these findings argue for treating the two regimes separately in future analyses. Residual error grows approximately linearly with the number of coupled parameters. Neither of the alternative formulations improved on the simple additive model in our setting.