

Toward Dose-Adaptive Gait Signatures: Evidence from Chronic Wound Care

Kenji Yarborough, Yusuf Jovanovic, Dilnoza Xanthos

Institute for Applied Dynamics, Varnholt

Section: Biomedical
Published: 2024-03-15
DOI: 10.5281/mjas.13.1.03
Keywords: cost accounting; reproducibility; benchmarking

Abstract

We examine the behaviour of organoid cultures in ambulatory trials. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Residual error grows approximately linearly with the number of coupled parameters. The reduced model retains most of the explanatory power at a fraction of the computational cost.

1. Introduction

We release the processing scripts and intermediate artefacts to support independent verification. Taken together, these findings argue for treating the two regimes separately in future analyses. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Calibration drift over the trial period remained below the threshold set out in the protocol. Sensitivity to the smoothing window proved modest across the range we examined. The reduced model retains most of the explanatory power at a fraction of the computational cost. Longer observation windows narrow the confidence interval but do not shift the central estimate. Neither of the alternative formulations improved on the simple additive model in our setting.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Replicate agreement was high, suggesting the protocol is robust to small handling differences. We release the processing scripts and intermediate artefacts to support independent verification.

2. Related Work

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Sensitivity to the smoothing window proved modest across the range we examined. Longer observation windows narrow the confidence interval but do not shift the central estimate. Calibration drift over the trial period remained below the threshold set out in the protocol. Neither of the alternative formulations improved on the simple additive model in our setting. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. We release the processing scripts and intermediate artefacts to support independent verification. The measurements reported here were collected over twelve independent trials under controlled conditions.

The measurements reported here were collected over twelve independent trials under controlled conditions. Longer observation windows narrow the confidence interval but do not shift the central estimate. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Residual error grows approximately linearly with the number of coupled parameters. The reduced model retains most of the explanatory power at a fraction of the computational cost. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Neither of the alternative formulations improved on the simple additive model in our setting.

Neither of the alternative formulations improved on the simple additive model in our setting. The reduced model retains most of the explanatory power at a fraction of the computational cost. Longer observation windows narrow the confidence interval but do not shift the central estimate. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Calibration drift over the trial period remained below the threshold set out in the protocol. Replicate agreement was high, suggesting the protocol is robust to small handling differences.

3. Materials and Methods

Replicate agreement was high, suggesting the protocol is robust to small handling differences. Sensitivity to the smoothing window proved modest across the range we examined. Longer observation windows narrow the confidence interval but do not shift the central estimate. Taken together, these findings argue for treating the two regimes separately in future analyses.

Calibration drift over the trial period remained below the threshold set out in the protocol. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Residual error grows approximately linearly with the number of coupled parameters. Neither of the alternative formulations improved on the simple additive model in our setting.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. The measurements reported here were collected over twelve independent trials under controlled conditions. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Neither of the alternative formulations improved on the simple additive model in our setting. Calibration drift over the trial period remained below the threshold set out in the protocol. Longer observation windows narrow the confidence interval but do not shift the central estimate. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study.

4. Experimental Setup

Replicate agreement was high, suggesting the protocol is robust to small handling differences. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. We release the processing scripts and intermediate artefacts to support independent verification.

Residual error grows approximately linearly with the number of coupled parameters. Sensitivity to the smoothing window proved modest across the range we examined. We release the processing scripts and intermediate artefacts to support independent verification. A modest but repeatable bias appears in the

lowest-magnitude bin and warrants further study. Discrepancies at the boundary are consistent with the simplifying assumptions noted above.

Replicate agreement was high, suggesting the protocol is robust to small handling differences. Longer observation windows narrow the confidence interval but do not shift the central estimate. Neither of the alternative formulations improved on the simple additive model in our setting. Residual error grows approximately linearly with the number of coupled parameters.

5. Results

The reduced model retains most of the explanatory power at a fraction of the computational cost. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Calibration drift over the trial period remained below the threshold set out in the protocol. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Sensitivity to the smoothing window proved modest across the range we examined. Taken together, these findings argue for treating the two regimes separately in future analyses. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Residual error grows approximately linearly with the number of coupled parameters.

Sensitivity to the smoothing window proved modest across the range we examined. Calibration drift over the trial period remained below the threshold set out in the protocol. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Neither of the alternative formulations improved on the simple additive model in our setting. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Discrepancies at the boundary are consistent with the simplifying assumptions noted above.

Longer observation windows narrow the confidence interval but do not shift the central estimate. Calibration drift over the trial period remained below the threshold set out in the protocol. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. The reduced model retains most of the explanatory power at a fraction of the computational cost. Replicate agreement was high, suggesting the protocol is robust to small handling differences. The measurements reported here were collected over twelve independent trials under controlled conditions. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. We release the processing scripts and intermediate artefacts to support independent verification. Neither of the alternative formulations improved on the simple additive model in our setting.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. The measurements reported here were collected over twelve independent trials under controlled conditions. Longer observation windows narrow the confidence interval but do not shift the central estimate. Calibration drift over the trial period remained below the threshold set out in the protocol. Taken together, these findings argue for treating the two regimes separately in future analyses.