

Perfusion Assays Revisited: A Wearable Perspective

Fenna Brandvold, Mathis Eyre, Faisal Rasmussen

Hollis Mountain Observatory

Section: Biomedical
Published: 2025-06-06
DOI: 10.5281/mjas.14.2.05
Keywords: scaling laws; reproducibility; cohort design; calibration

Abstract

This paper reports a systematic evaluation of immunomodulatory imaging pipelines. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. The reduced model retains most of the explanatory power at a fraction of the computational cost. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study.

1. Introduction

A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Neither of the alternative formulations improved on the simple additive model in our setting. Longer observation windows narrow the confidence interval but do not shift the central estimate. Residual error grows approximately linearly with the number of coupled parameters. The measurements reported here were collected over twelve independent trials under controlled conditions.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Replicate agreement was high, suggesting the protocol is robust to small handling differences. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. The measurements reported here were collected over twelve independent trials under controlled conditions. Calibration drift over the trial period remained below the threshold set out in the protocol. Neither of the alternative formulations improved on the simple additive model in our setting.

A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. The reduced model retains most of the explanatory power at a fraction of the computational cost. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Residual error grows approximately linearly with the number of coupled parameters. The measurements reported here were collected over twelve independent trials under controlled conditions. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Sensitivity to the smoothing window proved modest across the range we examined.

2. Related Work

Residual error grows approximately linearly with the number of coupled parameters. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Sensitivity to the smoothing window proved modest across the range we examined. Taken together, these findings argue for treating the two regimes separately in future analyses.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Taken together, these findings argue for treating the two regimes separately in future analyses. The measurements reported here were collected over twelve independent trials under controlled conditions. Neither of the alternative formulations improved on the simple additive model in our setting. Sensitivity to the smoothing window proved modest across the range we examined.

Replicate agreement was high, suggesting the protocol is robust to small handling differences. Longer observation windows narrow the confidence interval but do not shift the central estimate. Neither of the alternative formulations improved on the simple additive model in our setting. Taken together, these findings argue for treating the two regimes separately in future analyses. The reduced model retains most of the explanatory power at a fraction of the computational cost. Sensitivity to the smoothing window proved modest across the range we examined. We release the processing scripts and intermediate artefacts to support independent verification. Residual error grows approximately linearly with the number of coupled parameters. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity.

3. Materials and Methods

Sensitivity to the smoothing window proved modest across the range we examined. Residual error grows approximately linearly with the number of coupled parameters. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Neither of the alternative formulations improved on the simple additive model in our setting. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. We release the processing scripts and intermediate artefacts to support independent verification.

Replicate agreement was high, suggesting the protocol is robust to small handling differences. Taken together, these findings argue for treating the two regimes separately in future analyses. Sensitivity to the smoothing window proved modest across the range we examined. The reduced model retains most of the explanatory power at a fraction of the computational cost.

Taken together, these findings argue for treating the two regimes separately in future analyses. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. The measurements reported here were collected over twelve independent trials under controlled conditions. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Taken together, these findings argue for treating the two regimes separately in future analyses. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Neither of the alternative formulations improved on the simple additive model in our setting. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity.

4. Experimental Setup

The reduced model retains most of the explanatory power at a fraction of the computational cost. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The measurements reported here were collected over twelve independent trials under controlled conditions. Neither of the alternative formulations improved on the simple additive model in our setting. Sensitivity to the smoothing window proved modest across the range we examined.

Longer observation windows narrow the confidence interval but do not shift the central estimate. The measurements reported here were collected over twelve independent trials under controlled conditions. Residual error grows approximately linearly with the number of coupled parameters. Calibration drift over the trial period remained below the threshold set out in the protocol. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study.

Neither of the alternative formulations improved on the simple additive model in our setting. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Calibration drift over the trial period remained below the threshold set out in the protocol. Taken together, these findings argue for treating the two regimes separately in future analyses. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Residual error grows approximately linearly with the number of coupled parameters. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study.

We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Calibration drift over the trial period remained below the threshold set out in the protocol. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The reduced model retains most of the explanatory power at a fraction of the computational cost.