

Thin Films Revisited: A Thermally Stable Perspective

Xiulan Yarborough, Bastian Rasmussen, Anders Halvorsen, Elke Rasmussen, Alina Rasmussen

Calder Bay University

Section: Materials

Published: 2022-03-28

DOI: 10.5281/mjas.11.1.06

Keywords: parameter sweep; cross-validation; surrogate modelling

Abstract

We examine the behaviour of lattice alloys in corrosive media. Longer observation windows narrow the confidence interval but do not shift the central estimate. Residual error grows approximately linearly with the number of coupled parameters. Calibration drift over the trial period remained below the threshold set out in the protocol.

1. Introduction

Calibration drift over the trial period remained below the threshold set out in the protocol. Residual error grows approximately linearly with the number of coupled parameters. We release the processing scripts and intermediate artefacts to support independent verification. Sensitivity to the smoothing window proved modest across the range we examined. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Replicate agreement was high, suggesting the protocol is robust to small handling differences.

The measurements reported here were collected over twelve independent trials under controlled conditions. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Residual error grows approximately linearly with the number of coupled parameters. Discrepancies at the boundary are consistent with the simplifying assumptions noted above.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Neither of the alternative formulations improved on the simple additive model in our setting. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Sensitivity to the smoothing window proved modest across the range we examined. Replicate agreement was high, suggesting the protocol is robust to small handling differences.

2. Related Work

Longer observation windows narrow the confidence interval but do not shift the central estimate. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Calibration drift over the trial period remained below the threshold set out in the protocol. The measurements reported here were collected over twelve independent trials under controlled conditions. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. We release the processing scripts and intermediate artefacts to support independent verification. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.

We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. We release the processing scripts and intermediate artefacts to support independent verification. Calibration drift over the trial period remained below the threshold set out in the protocol. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. The measurements reported here were collected over twelve independent trials under controlled conditions.

Sensitivity to the smoothing window proved modest across the range we examined. Residual error grows approximately linearly with the number of coupled parameters. Taken together, these findings argue for treating the two regimes separately in future analyses. The measurements reported here were collected over twelve independent trials under controlled conditions.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. We release the processing scripts and intermediate artefacts to support independent verification. Residual error grows approximately linearly with the number of coupled parameters. Calibration drift over the trial period remained below the threshold set out in the protocol. The measurements reported here were collected over twelve independent trials under controlled conditions.

Calibration drift over the trial period remained below the threshold set out in the protocol. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Residual error grows approximately linearly with the number of coupled parameters. Sensitivity to the smoothing window proved modest across the range we examined. Taken together, these findings argue for treating the two regimes separately in future analyses.

3. Materials and Methods

Neither of the alternative formulations improved on the simple additive model in our setting. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. We release the processing scripts and intermediate artefacts to support independent verification. Sensitivity to the smoothing window proved modest across the range we examined. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Taken together, these findings argue for treating the two regimes separately in future analyses. The measurements reported here were collected over twelve independent trials under controlled conditions.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Neither of the alternative formulations improved on the simple additive model in our setting. Longer observation

windows narrow the confidence interval but do not shift the central estimate. We release the processing scripts and intermediate artefacts to support independent verification. Taken together, these findings argue for treating the two regimes separately in future analyses. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Residual error grows approximately linearly with the number of coupled parameters. Sensitivity to the smoothing window proved modest across the range we examined.

4. Experimental Setup

Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Calibration drift over the trial period remained below the threshold set out in the protocol. Sensitivity to the smoothing window proved modest across the range we examined. The measurements reported here were collected over twelve independent trials under controlled conditions. The reduced model retains most of the explanatory power at a fraction of the computational cost. We release the processing scripts and intermediate artefacts to support independent verification. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Residual error grows approximately linearly with the number of coupled parameters.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Calibration drift over the trial period remained below the threshold set out in the protocol. Longer observation windows narrow the confidence interval but do not shift the central estimate. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Sensitivity to the smoothing window proved modest across the range we examined. The measurements reported here were collected over twelve independent trials under controlled conditions. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Neither of the alternative formulations improved on the simple additive model in our setting.

Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Sensitivity to the smoothing window proved modest across the range we examined. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.

Longer observation windows narrow the confidence interval but do not shift the central estimate. Replicate agreement was high, suggesting the protocol is robust to small handling differences. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Sensitivity to the smoothing window proved modest across the range we examined. We release the processing scripts and intermediate artefacts to support independent verification. Calibration drift over the trial period remained below the threshold set out in the protocol.

5. Results

We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Residual error grows approximately linearly with the number of coupled parameters. Neither of the alternative formulations improved on the simple additive model in our setting. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. The measurements reported here were collected over twelve independent trials under controlled conditions. We release the processing scripts and intermediate artefacts to support independent verification. Calibration drift over the trial period remained below the threshold set out in the protocol. Longer observation windows narrow the confidence interval but do not shift the central estimate.

Discrepancies at the boundary are consistent with the simplifying assumptions noted above. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. We release the processing scripts and intermediate artefacts to support independent verification. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Longer observation windows narrow the confidence interval but do not shift the central estimate.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Taken together, these findings argue for treating the two regimes separately in future analyses. Calibration drift over the trial period remained below the threshold set out in the protocol. Sensitivity to the smoothing window proved modest across the range we examined. We release the processing scripts and intermediate artefacts to support independent verification.

6. Discussion

The reduced model retains most of the explanatory power at a fraction of the computational cost. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Calibration drift over the trial period remained below the threshold set out in the protocol.

The measurements reported here were collected over twelve independent trials under controlled conditions. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study.

Taken together, these findings argue for treating the two regimes separately in future analyses. Neither of the alternative formulations improved on the simple additive model in our setting. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity.

7. Limitations

We release the processing scripts and intermediate artefacts to support independent verification. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Longer observation windows narrow the confidence interval but do not shift the central estimate. Taken together, these findings argue for treating the two regimes separately in future analyses. Discrepancies at the boundary are consistent with the simplifying assumptions noted above.

Residual error grows approximately linearly with the number of coupled parameters. Longer observation windows narrow the confidence interval but do not shift the central estimate. Sensitivity to the smoothing window proved modest across the range we examined. The reduced model retains most of the explanatory power at a fraction of the computational cost.

Calibration drift over the trial period remained below the threshold set out in the protocol. Taken together, these findings argue for treating the two regimes separately in future analyses. Residual error grows approximately linearly with the number of coupled parameters. Sensitivity to the smoothing window proved modest across the range we examined. We release the processing scripts and intermediate artefacts to support independent verification. Longer observation windows narrow the confidence interval but do not shift the central estimate. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The reduced model retains most of the explanatory power at a fraction of the computational cost.