

Characterizing Perfusion Assays in Chronic Wound Care

Zofia Gorecki, Ulrich Lindqvist, Lucia Vasquez, Bastian Aalto, Imogen Halvorsen

Sableport Marine Laboratory

Section: Biomedical

Published: 2022-12-23

DOI: 10.5281/mjas.11.4.05

Keywords: cost accounting; noise robustness; field trial; sensitivity analysis

Abstract

Work in biomedical research has long relied on assumptions we test directly here. Neither of the alternative formulations improved on the simple additive model in our setting. Calibration drift over the trial period remained below the threshold set out in the protocol. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.

1. Introduction

We release the processing scripts and intermediate artefacts to support independent verification. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Longer observation windows narrow the confidence interval but do not shift the central estimate. Residual error grows approximately linearly with the number of coupled parameters. The reduced model retains most of the explanatory power at a fraction of the computational cost. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.

Calibration drift over the trial period remained below the threshold set out in the protocol. Taken together, these findings argue for treating the two regimes separately in future analyses. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. The reduced model retains most of the explanatory power at a fraction of the computational cost. Replicate agreement was high, suggesting the protocol is robust to small handling differences. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study.

The measurements reported here were collected over twelve independent trials under controlled conditions. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Taken together, these findings argue for treating the two regimes separately in future analyses. Replicate agreement was high, suggesting the protocol is robust to small handling differences. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity.

Calibration drift over the trial period remained below the threshold set out in the protocol. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Sensitivity to the smoothing window proved modest across the range we examined. Longer observation windows narrow the confidence interval but do not shift the

central estimate. Residual error grows approximately linearly with the number of coupled parameters. We release the processing scripts and intermediate artefacts to support independent verification. The reduced model retains most of the explanatory power at a fraction of the computational cost.

2. Related Work

Replicate agreement was high, suggesting the protocol is robust to small handling differences. We release the processing scripts and intermediate artefacts to support independent verification. Residual error grows approximately linearly with the number of coupled parameters. Longer observation windows narrow the confidence interval but do not shift the central estimate. Taken together, these findings argue for treating the two regimes separately in future analyses. Sensitivity to the smoothing window proved modest across the range we examined. The reduced model retains most of the explanatory power at a fraction of the computational cost. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study.

The measurements reported here were collected over twelve independent trials under controlled conditions. Residual error grows approximately linearly with the number of coupled parameters. The reduced model retains most of the explanatory power at a fraction of the computational cost. We release the processing scripts and intermediate artefacts to support independent verification. Neither of the alternative formulations improved on the simple additive model in our setting. Calibration drift over the trial period remained below the threshold set out in the protocol. Taken together, these findings argue for treating the two regimes separately in future analyses. Sensitivity to the smoothing window proved modest across the range we examined.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Residual error grows approximately linearly with the number of coupled parameters. Calibration drift over the trial period remained below the threshold set out in the protocol. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Longer observation windows narrow the confidence interval but do not shift the central estimate. We release the processing scripts and intermediate artefacts to support independent verification. The measurements reported here were collected over twelve independent trials under controlled conditions. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Taken together, these findings argue for treating the two regimes separately in future analyses.

3. Materials and Methods

Residual error grows approximately linearly with the number of coupled parameters. We release the processing scripts and intermediate artefacts to support independent verification. The measurements reported here were collected over twelve independent trials under controlled conditions. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Longer observation windows narrow the confidence interval but do not shift the central estimate. The reduced model retains most of the explanatory power at a fraction of the computational cost.

Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The measurements reported here were collected over twelve independent trials under controlled conditions. We release the processing scripts and intermediate artefacts to support independent verification. Residual error grows approximately linearly with the number of coupled parameters. Calibration drift over the trial period remained below the threshold set out in the protocol. Taken together, these findings argue for treating the two regimes separately in future analyses. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Sensitivity to the smoothing window proved modest across the range we examined.

Longer observation windows narrow the confidence interval but do not shift the central estimate.

4. Experimental Setup

We release the processing scripts and intermediate artefacts to support independent verification. Taken together, these findings argue for treating the two regimes separately in future analyses. Sensitivity to the smoothing window proved modest across the range we examined. Longer observation windows narrow the confidence interval but do not shift the central estimate. Neither of the alternative formulations improved on the simple additive model in our setting.

The measurements reported here were collected over twelve independent trials under controlled conditions. The reduced model retains most of the explanatory power at a fraction of the computational cost. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity.

5. Results

Replicate agreement was high, suggesting the protocol is robust to small handling differences. Longer observation windows narrow the confidence interval but do not shift the central estimate. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Sensitivity to the smoothing window proved modest across the range we examined. Calibration drift over the trial period remained below the threshold set out in the protocol.

Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Calibration drift over the trial period remained below the threshold set out in the protocol. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Replicate agreement was high, suggesting the protocol is robust to small handling differences. The reduced model retains most of the explanatory power at a fraction of the computational cost.

6. Discussion

Sensitivity to the smoothing window proved modest across the range we examined. Neither of the alternative formulations improved on the simple additive model in our setting. Residual error grows approximately linearly with the number of coupled parameters. Taken together, these findings argue for treating the two regimes separately in future analyses. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Neither of the alternative formulations improved on the simple additive model in our setting. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The measurements reported here were collected over twelve independent trials under controlled conditions. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Longer observation windows narrow the confidence interval but do not shift the central estimate.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Residual error grows approximately linearly with the number of coupled parameters. The measurements reported here were collected over twelve independent trials under controlled conditions. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Longer observation windows narrow the confidence interval but do not shift the central estimate. Sensitivity to the smoothing window proved modest across the range we examined. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Neither of the alternative formulations improved on the simple additive model in our setting.