

Characterizing Fiber Scaffolds in Marine Environments

Fenna Marchetti, Valeria Dusseault, Sten Marchetti, Yusuf Brandvold, Caius Thibodeaux

Institute for Applied Dynamics, Varnholt

Section: Materials

Published: 2025-06-09

DOI: 10.5281/mjas.14.2.04

Keywords: sensitivity analysis; reproducibility; error propagation; noise robustness; ablation study

Abstract

This paper reports a systematic evaluation of anisotropic thin films. Calibration drift over the trial period remained below the threshold set out in the protocol. Neither of the alternative formulations improved on the simple additive model in our setting. The reduced model retains most of the explanatory power at a fraction of the computational cost.

1. Introduction

Sensitivity to the smoothing window proved modest across the range we examined. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. The measurements reported here were collected over twelve independent trials under controlled conditions. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Residual error grows approximately linearly with the number of coupled parameters. The reduced model retains most of the explanatory power at a fraction of the computational cost. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Calibration drift over the trial period remained below the threshold set out in the protocol. Replicate agreement was high, suggesting the protocol is robust to small handling differences.

We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. The reduced model retains most of the explanatory power at a fraction of the computational cost. Replicate agreement was high, suggesting the protocol is robust to small handling differences. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Sensitivity to the smoothing window proved modest across the range we examined. We release the processing scripts and intermediate artefacts to support independent verification. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.

Longer observation windows narrow the confidence interval but do not shift the central estimate. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The reduced model retains most of the explanatory power at a fraction of the computational cost. Sensitivity to the smoothing window proved modest across the range we examined. The measurements reported here were collected over twelve independent trials under controlled conditions.

2. Related Work

A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. Sensitivity to the smoothing window proved modest across the range we examined. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. The measurements reported here were collected over twelve independent trials under controlled conditions. Calibration drift over the trial period remained below the threshold set out in the protocol. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Discrepancies at the boundary are consistent with the simplifying assumptions noted above.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Sensitivity to the smoothing window proved modest across the range we examined. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Longer observation windows narrow the confidence interval but do not shift the central estimate. Calibration drift over the trial period remained below the threshold set out in the protocol.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Neither of the alternative formulations improved on the simple additive model in our setting. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Calibration drift over the trial period remained below the threshold set out in the protocol. Residual error grows approximately linearly with the number of coupled parameters. Longer observation windows narrow the confidence interval but do not shift the central estimate.

The reduced model retains most of the explanatory power at a fraction of the computational cost. Longer observation windows narrow the confidence interval but do not shift the central estimate. Sensitivity to the smoothing window proved modest across the range we examined. The measurements reported here were collected over twelve independent trials under controlled conditions. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Neither of the alternative formulations improved on the simple additive model in our setting. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Taken together, these findings argue for treating the two regimes separately in future analyses.

3. Materials and Methods

We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. We release the processing scripts and intermediate artefacts to support independent verification. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. Replicate agreement was high, suggesting the protocol is robust to small handling differences. Taken together, these findings argue for treating the two regimes separately in future analyses. Sensitivity to the smoothing window proved modest across the range we examined. Residual error grows approximately linearly with the number of coupled parameters.

Residual error grows approximately linearly with the number of coupled parameters. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The measurements reported here were collected over twelve independent trials under controlled conditions. Taken together, these findings argue for treating the two regimes separately in future analyses. Neither of the alternative formulations improved on the simple additive model in our setting. We observe a pronounced inflection once the load exceeds roughly sixty

percent of nominal capacity.

Calibration drift over the trial period remained below the threshold set out in the protocol. Discrepancies at the boundary are consistent with the simplifying assumptions noted above. Taken together, these findings argue for treating the two regimes separately in future analyses. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Sensitivity to the smoothing window proved modest across the range we examined. We release the processing scripts and intermediate artefacts to support independent verification. Replicate agreement was high, suggesting the protocol is robust to small handling differences.

Discrepancies at the boundary are consistent with the simplifying assumptions noted above. The measurements reported here were collected over twelve independent trials under controlled conditions. Sensitivity to the smoothing window proved modest across the range we examined. Calibration drift over the trial period remained below the threshold set out in the protocol. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.

4. Experimental Setup

Residual error grows approximately linearly with the number of coupled parameters. The reduced model retains most of the explanatory power at a fraction of the computational cost. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance. We observe a pronounced inflection once the load exceeds roughly sixty percent of nominal capacity. Longer observation windows narrow the confidence interval but do not shift the central estimate. The measurements reported here were collected over twelve independent trials under controlled conditions. Taken together, these findings argue for treating the two regimes separately in future analyses.

Calibration drift over the trial period remained below the threshold set out in the protocol. A modest but repeatable bias appears in the lowest-magnitude bin and warrants further study. The measurements reported here were collected over twelve independent trials under controlled conditions. Our baseline reproduces the behaviour described in earlier work to within the stated tolerance.