

Problem 2367: Number of Arithmetic Triplets

Problem Information

Difficulty: Easy

Acceptance Rate: 85.27%

Paid Only: No

Tags: Array, Hash Table, Two Pointers, Enumeration

Problem Description

You are given a **0-indexed** , **strictly increasing** integer array `nums` and a positive integer `diff`. A triplet `(i, j, k)` is an **arithmetic triplet** if the following conditions are met:

`i < j < k` , `nums[j] - nums[i] == diff` , and `nums[k] - nums[j] == diff`.

Return the number of unique arithmetic triplets.

Example 1.

Input: `nums = [0,1,4,6,7,10]`, `diff = 3` **Output:** 2 **Explanation:** (1, 2, 4) is an arithmetic triplet because both `7 - 4 == 3` and `4 - 1 == 3`. (2, 4, 5) is an arithmetic triplet because both `10 - 7 == 3` and `7 - 4 == 3`.

Example 2.

Input: `nums = [4,5,6,7,8,9]`, `diff = 2` **Output:** 2 **Explanation:** (0, 2, 4) is an arithmetic triplet because both `8 - 6 == 2` and `6 - 4 == 2`. (1, 3, 5) is an arithmetic triplet because both `9 - 7 == 2` and `7 - 5 == 2`.

Constraints:

`3 <= nums.length <= 200` `0 <= nums[i] <= 200` `1 <= diff <= 50` `nums` is **strictly increasing**.

Code Snippets

C++:

```
class Solution {  
public:  
    int arithmeticTriplets(vector<int>& nums, int diff) {  
  
    }  
};
```

Java:

```
class Solution {  
    public int arithmeticTriplets(int[] nums, int diff) {  
  
    }  
}
```

Python3:

```
class Solution:  
    def arithmeticTriplets(self, nums: List[int], diff: int) -> int:
```