

Problem 2028: Find Missing Observations

Problem Information

Difficulty: Medium

Acceptance Rate: 57.38%

Paid Only: No

Tags: Array, Math, Simulation

Problem Description

You have observations of $n + m$ 6-sided dice rolls with each face numbered from 1 to 6. n of the observations went missing, and you only have the observations of m rolls. Fortunately, you have also calculated the average value of the $n + m$ rolls.

You are given an integer array `rolls` of length m where `rolls[i]` is the value of the i th observation. You are also given the two integers `mean` and n .

Return an array of length n containing the missing observations such that the average value of the $n + m$ rolls is exactly `mean`. If there are multiple valid answers, return any of them. If no such array exists, return an empty array.

The average value of a set of k numbers is the sum of the numbers divided by k .

Note that `mean` is an integer, so the sum of the $n + m$ rolls should be divisible by $n + m$.

Example 1.

Input: `rolls = [3,2,4,3]`, `mean = 4`, $n = 2$ **Output:** `[6,6]` **Explanation:** The mean of all $n + m$ rolls is $(3 + 2 + 4 + 3 + 6 + 6) / 6 = 4$.

Example 2.

Input: `rolls = [1,5,6]`, `mean = 3`, $n = 4$ **Output:** `[2,3,2,2]` **Explanation:** The mean of all $n + m$ rolls is $(1 + 5 + 6 + 2 + 3 + 2 + 2) / 7 = 3$.

Example 3.

****Input:**** rolls = [1,2,3,4], mean = 6, n = 4 ****Output:**** [] ****Explanation:**** It is impossible for the mean to be 6 no matter what the 4 missing rolls are.

****Constraints:****

* `m == rolls.length` * `1 <= n, m <= 105` * `1 <= rolls[i], mean <= 6`

Code Snippets

C++:

```
class Solution {
public:
    vector<int> missingRolls(vector<int>& rolls, int mean, int n) {

    }
};
```

Java:

```
class Solution {
    public int[] missingRolls(int[] rolls, int mean, int n) {

    }
}
```

Python3:

```
class Solution:
    def missingRolls(self, rolls: List[int], mean: int, n: int) -> List[int]:
```