

Problem 3426: Manhattan Distances of All Arrangements of Pieces

Problem Information

Difficulty: **Hard**

Acceptance Rate: 34.40%

Paid Only: No

Tags: Math, Combinatorics

Problem Description

You are given three integers m , n , and k .

There is a rectangular grid of size $m \times n$ containing k identical pieces. Return the sum of Manhattan distances between every pair of pieces over all **valid arrangements** of pieces.

A **valid arrangement** is a placement of all k pieces on the grid with **at most** one piece per cell.

Since the answer may be very large, return it **modulo** $10^9 + 7$.

The Manhattan Distance between two cells (x_i, y_i) and (x_j, y_j) is $|x_i - x_j| + |y_i - y_j|$.


Example 1:

Input: $m = 2, n = 2, k = 2$

Output: 8

Explanation:

The valid arrangements of pieces on the board are:



* In the first 4 arrangements, the Manhattan distance between the two pieces is 1. * In the last 2 arrangements, the Manhattan distance between the two pieces is 2.

Thus, the total Manhattan distance across all valid arrangements is $1 + 1 + 1 + 1 + 2 + 2 = 8$.

Example 2:

Input: $m = 1, n = 4, k = 3$

Output: 20

Explanation:

The valid arrangements of pieces on the board are:



* The first and last arrangements have a total Manhattan distance of $1 + 1 + 2 = 4$. * The middle two arrangements have a total Manhattan distance of $1 + 2 + 3 = 6$.

The total Manhattan distance between all pairs of pieces across all arrangements is $4 + 6 + 6 + 4 = 20$.

Constraints:

$1 \leq m, n \leq 10^5$ $2 \leq m * n \leq 10^5$ $2 \leq k \leq m * n$

Code Snippets

C++:

```
class Solution {
public:
    int distanceSum(int m, int n, int k) {

    }
};
```

Java:

```
class Solution {  
    public int distanceSum(int m, int n, int k) {  
  
    }  
}
```

Python3:

```
class Solution:  
    def distanceSum(self, m: int, n: int, k: int) -> int:
```