

Problem 2143: Choose Numbers From Two Arrays in Range

Problem Information

Difficulty: **Hard**

Acceptance Rate: 52.60%

Paid Only: Yes

Tags: Array, Dynamic Programming

Problem Description

You are given two **0-indexed** integer arrays `nums1` and `nums2` of length `n`.

A range `[l, r]` (**inclusive**) where `0 ≤ l ≤ r < n` is **balanced** if:

* For every `i` in the range `[l, r]`, you pick either `nums1[i]` or `nums2[i]`. * The sum of the numbers you pick from `nums1` equals to the sum of the numbers you pick from `nums2` (the sum is considered to be `0` if you pick no numbers from an array).

Two **balanced** ranges from `[l1, r1]` and `[l2, r2]` are considered to be **different** if at least one of the following is true:

* `l1 != l2` * `r1 != r2` * `nums1[i]` is picked in the first range, and `nums2[i]` is picked in the second range or **vice versa** for at least one `i`.

Return the number of different ranges that are balanced. Since the answer may be very large, return it modulo `109 + 7`.

Example 1.

Input: `nums1 = [1,2,5], nums2 = [2,6,3]` **Output:** `3` **Explanation:** The balanced ranges are: - `[0, 1]` where we choose `nums2[0]`, and `nums1[1]`. The sum of the numbers chosen from `nums1` equals the sum of the numbers chosen from `nums2`: `2 = 2`. - `[0, 2]` where we choose `nums1[0]`, `nums2[1]`, and `nums1[2]`. The sum of the numbers chosen from `nums1` equals the sum of the numbers chosen from `nums2`: `1 + 5 = 6`. - `[0, 2]` where we choose `nums1[0]`, `nums1[1]`, and `nums2[2]`. The sum of the numbers chosen from `nums1` equals the

sum of the numbers chosen from nums2: $1 + 2 = 3$. Note that the second and third balanced ranges are different. In the second balanced range, we choose nums2[1] and in the third balanced range, we choose nums1[1].

Example 2:

Input: nums1 = [0,1], nums2 = [1,0] **Output:** 4 **Explanation:** The balanced ranges are: - [0, 0] where we choose nums1[0]. The sum of the numbers chosen from nums1 equals the sum of the numbers chosen from nums2: $0 = 0$. - [1, 1] where we choose nums2[1]. The sum of the numbers chosen from nums1 equals the sum of the numbers chosen from nums2: $0 = 0$. - [0, 1] where we choose nums1[0] and nums2[1]. The sum of the numbers chosen from nums1 equals the sum of the numbers chosen from nums2: $0 = 0$. - [0, 1] where we choose nums2[0] and nums1[1]. The sum of the numbers chosen from nums1 equals the sum of the numbers chosen from nums2: $1 = 1$.

Constraints:

$n == \text{nums1.length} == \text{nums2.length}$ $1 \leq n \leq 100$ $0 \leq \text{nums1}[i], \text{nums2}[i] \leq 100$

Code Snippets

C++:

```
class Solution {
public:
    int countSubranges(vector<int>& nums1, vector<int>& nums2) {

    }
};
```

Java:

```
class Solution {
    public int countSubranges(int[] nums1, int[] nums2) {

    }
}
```

Python3:

```
class Solution:
    def countSubranges(self, nums1: List[int], nums2: List[int]) -> int:
```