

# Problem 2210: Count Hills and Valleys in an Array

## Problem Information

**Difficulty:** Easy

**Acceptance Rate:** 69.32%

**Paid Only:** No

**Tags:** Array

## Problem Description

You are given a **0-indexed** integer array `nums`. An index `i` is part of a **hill** in `nums` if the closest non-equal neighbors of `i` are smaller than `nums[i]`. Similarly, an index `i` is part of a **valley** in `nums` if the closest non-equal neighbors of `i` are larger than `nums[i]`. Adjacent indices `i` and `j` are part of the **same** hill or valley if `nums[i] == nums[j]`.

Note that for an index to be part of a hill or valley, it must have a non-equal neighbor on **both** the left and right of the index.

Return \_the number of hills and valleys in\_ `nums`.

**Example 1:**

**Input:** nums = [2,4,1,1,6,5] **Output:** 3 **Explanation:** At index 0: There is no non-equal neighbor of 2 on the left, so index 0 is neither a hill nor a valley. At index 1: The closest non-equal neighbors of 4 are 2 and 1. Since 4 > 2 and 4 > 1, index 1 is a hill. At index 2: The closest non-equal neighbors of 1 are 4 and 6. Since 1 < 4 and 1 < 6, index 2 is a valley. At index 3: The closest non-equal neighbors of 1 are 4 and 6. Since 1 < 4 and 1 < 6, index 3 is a valley, but note that it is part of the same valley as index 2. At index 4: The closest non-equal neighbors of 6 are 1 and 5. Since 6 > 1 and 6 > 5, index 4 is a hill. At index 5: There is no non-equal neighbor of 5 on the right, so index 5 is neither a hill nor a valley. There are 3 hills and valleys so we return 3.

**Example 2:**

**Input:** nums = [6,6,5,5,4,1] **Output:** 0 **Explanation:** At index 0: There is no non-equal neighbor of 6 on the left, so index 0 is neither a hill nor a valley. At index 1: There is no non-equal neighbor of 6 on the left, so index 1 is neither a hill nor a valley. At index 2: The closest non-equal neighbors of 5 are 6 and 4. Since  $5 < 6$  and  $5 > 4$ , index 2 is neither a hill nor a valley. At index 3: The closest non-equal neighbors of 5 are 6 and 4. Since  $5 < 6$  and  $5 > 4$ , index 3 is neither a hill nor a valley. At index 4: The closest non-equal neighbors of 4 are 5 and 1. Since  $4 < 5$  and  $4 > 1$ , index 4 is neither a hill nor a valley. At index 5: There is no non-equal neighbor of 1 on the right, so index 5 is neither a hill nor a valley. There are 0 hills and valleys so we return 0.

**Constraints:**

$3 \leq \text{nums.length} \leq 100$   $1 \leq \text{nums}[i] \leq 100$

## Code Snippets

**C++:**

```
class Solution {
public:
    int countHillValley(vector<int>& nums) {
        }
    };
}
```

**Java:**

```
class Solution {
public int countHillValley(int[] nums) {
        }
    }
}
```

**Python3:**

```
class Solution:
    def countHillValley(self, nums: List[int]) -> int:
```